



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:47 PM UTC

PDB ID : 6PTR / pdb_00006ptr
Title : Crystal structure of a DnaN sliding clamp (DNA polymerase III subunit beta) from *Bartonella birtlesii* bound to griselimycin
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2019-07-16
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	
1	B	381	
2	X	11	
2	Y	11	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	370	Total	C	N	O	S	0	12	0
			2884	1831	491	548	14			
1	B	372	Total	C	N	O	S	0	8	0
			2902	1842	499	549	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP J11Y24
A	-6	ALA	-	expression tag	UNP J11Y24
A	-5	HIS	-	expression tag	UNP J11Y24
A	-4	HIS	-	expression tag	UNP J11Y24
A	-3	HIS	-	expression tag	UNP J11Y24
A	-2	HIS	-	expression tag	UNP J11Y24
A	-1	HIS	-	expression tag	UNP J11Y24
A	0	HIS	-	expression tag	UNP J11Y24
B	-7	MET	-	initiating methionine	UNP J11Y24
B	-6	ALA	-	expression tag	UNP J11Y24
B	-5	HIS	-	expression tag	UNP J11Y24
B	-4	HIS	-	expression tag	UNP J11Y24
B	-3	HIS	-	expression tag	UNP J11Y24
B	-2	HIS	-	expression tag	UNP J11Y24
B	-1	HIS	-	expression tag	UNP J11Y24
B	0	HIS	-	expression tag	UNP J11Y24

- Molecule 2 is a protein called ACE-MVA-MP8-NZC-LEU-MP8-LEU-MVA-PRO-MLU-GLY.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	X	11	Total	C	N	O	0	0	0
			79	57	10	12			
2	Y	11	Total	C	N	O	0	0	0
			79	57	10	12			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total I 7 7	0	0
4	B	4	Total I 4 4	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	334	Total O 339 339	0	5

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	X	12	Total O 12 12	0	0
5	B	281	Total O 282 282	0	1
5	Y	8	Total O 8 8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.95Å 93.13Å 110.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.92 – 1.75 42.92 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.3 (42.92-1.75) 98.3 (42.92-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.68 (at 1.75Å)	Xtrriage
Refinement program	PHENIX (1.16_3546: ???)	Depositor
R, R_{free}	0.178 , 0.206 0.178 , 0.206	Depositor DCC
R_{free} test set	2034 reflections (2.11%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for k,h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6620	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MP8, NZC, EDO, IOD, ACE, MLU, MVA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2966	0.60	0/4023
1	B	0.34	0/2972	0.60	0/4028
2	X	0.29	0/24	0.55	0/26
2	Y	0.51	0/24	0.61	0/26
All	All	0.35	0/5986	0.60	0/8103

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	360	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2884	0	2907	26	0
1	B	2902	0	2949	18	0
2	X	79	0	96	2	0
2	Y	79	0	96	1	0
3	A	20	0	30	1	0
3	B	4	0	6	0	0
4	A	7	0	0	0	0
4	B	4	0	0	1	0
5	A	339	0	0	9	0
5	B	282	0	0	4	0
5	X	12	0	0	0	0
5	Y	8	0	0	0	0
All	All	6620	0	6084	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:363:GLU:HG2	1:B:364:VAL:HG23	1.60	0.83
1:B:361:ASN:ND2	5:B:601:HOH:O	2.13	0.74
1:A:50:ASP:OD2	5:A:601:HOH:O	2.12	0.67
1:B:167:ASP:OD2	1:B:171:LYS:NZ	2.28	0.66
1:B:218:ASP:OD1	5:B:602:HOH:O	2.14	0.64
1:A:252:GLN:NE2	5:A:608:HOH:O	2.36	0.57
1:A:300:VAL:HG22	1:A:310:GLU:HG2	1.85	0.57
1:B:189:ALA:HB1	1:B:193:VAL:HG21	1.87	0.56
1:A:82:PRO:HG2	1:A:85:SER:HB3	1.88	0.55
1:A:208:GLU:OE1	5:A:602:HOH:O	2.18	0.55
1:B:338:LEU:HB3	1:B:358:ASP:OD1	2.06	0.54
1:B:16:ARG:HD2	1:B:55:GLU:OE2	2.08	0.54
1:A:128[B]:CYS:SG	1:A:196:MET:HA	2.48	0.54
1:A:189:ALA:HB1	1:A:193:VAL:HG21	1.89	0.54
1:A:208:GLU:HB2	1:A:239[A]:PHE:CZ	2.43	0.53
1:A:51:LEU:HD11	1:A:239[A]:PHE:CZ	2.43	0.53
1:A:76:ASP:OD1	1:A:79[B]:ARG:NH2	2.40	0.52
1:A:185:VAL:HG23	1:A:364:VAL:HG22	1.92	0.51
1:A:142:ASP:HB3	3:A:502:EDO:H12	1.94	0.50
1:B:20:VAL:HG21	1:B:53[A]:VAL:CG2	2.41	0.50
1:B:116:ILE:HG22	4:B:503:IOD:I	2.82	0.50
1:A:212:LEU:HD23	1:A:234:LEU:HD21	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:PHE:HD2	1:A:181[A]:ARG:HH11	1.60	0.49
2:Y:7:LEU:HA	2:Y:8:MVA:HN1	1.59	0.48
1:A:51:LEU:HD21	1:A:239[A]:PHE:CZ	2.49	0.48
1:B:185:VAL:HG23	1:B:364:VAL:HG22	1.95	0.48
1:A:8:SER:HB2	5:A:793:HOH:O	2.12	0.47
1:A:154[B]:ARG:NH2	5:A:616:HOH:O	2.47	0.47
1:A:264:ILE:HB	1:A:316:THR:HB	1.95	0.47
1:B:363:GLU:HG3	5:B:729:HOH:O	2.14	0.47
1:B:8:SER:OG	1:B:83:ASP:OD1	2.33	0.46
1:A:79[A]:ARG:HD2	5:A:611:HOH:O	2.17	0.45
1:A:107:HIS:NE2	1:A:109:GLN:OE1	2.43	0.45
1:A:51:LEU:HD11	1:A:239[A]:PHE:CE1	2.52	0.45
1:B:53[A]:VAL:HG22	1:B:239:PHE:CD1	2.52	0.45
1:B:37:ALA:HB1	1:B:61[B]:ILE:HD12	1.99	0.44
1:A:181[B]:ARG:NH1	5:A:620:HOH:O	2.50	0.43
1:A:156:TYR:CD1	1:A:246:GLY:HA3	2.53	0.43
1:A:203:ARG:HD2	5:A:868:HOH:O	2.18	0.43
1:B:268[A]:GLN:NE2	5:B:615:HOH:O	2.50	0.43
1:B:338:LEU:HD22	1:B:358:ASP:N	2.34	0.42
1:B:20:VAL:HG21	1:B:53[A]:VAL:HG23	2.00	0.42
2:X:7:LEU:HA	2:X:8:MVA:HN1	1.63	0.42
2:X:3:MP8:HA	2:X:4:NZC:H40	1.69	0.41
1:A:60:ASN:O	5:A:603:HOH:O	2.21	0.41
1:A:208:GLU:HB2	1:A:239[A]:PHE:HZ	1.84	0.41
1:B:100:SER:HB2	1:B:109:GLN:NE2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	378/381 (99%)	365 (97%)	12 (3%)	1 (0%)	36 21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	376/381 (99%)	364 (97%)	12 (3%)	0	100	100
2	X	3/11 (27%)	3 (100%)	0	0	100	100
2	Y	3/11 (27%)	3 (100%)	0	0	100	100
All	All	760/784 (97%)	735 (97%)	24 (3%)	1 (0%)	48	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	217	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/335 (97%)	321 (99%)	3 (1%)	70	60
1	B	327/335 (98%)	323 (99%)	4 (1%)	63	49
2	X	3/3 (100%)	3 (100%)	0	100	100
2	Y	3/3 (100%)	3 (100%)	0	100	100
All	All	657/676 (97%)	650 (99%)	7 (1%)	70	52

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	VAL
1	A	42[A]	VAL
1	A	42[B]	VAL
1	B	19	ARG
1	B	97	SER
1	B	286	ARG
1	B	293	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	GLN
1	A	166	HIS
1	A	184	GLN
1	A	268	GLN
1	B	302	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLU	Y	10	2	7,8,9	0.40	0	7,9,11	0.37	0
2	MP8	X	3	2	6,8,9	0.97	0	3,10,12	1.35	0
2	MVA	Y	8	2	6,7,8	0.46	0	6,8,10	0.94	0
2	MP8	X	6	2	6,8,9	0.58	0	3,10,12	1.34	0
2	MP8	Y	6	2	6,8,9	0.88	0	3,10,12	1.07	0
2	MVA	X	8	2	6,7,8	0.32	0	6,8,10	0.90	0
2	NZC	X	4	2	6,7,8	0.40	0	6,8,10	0.67	0
2	MVA	Y	2	2	6,7,8	0.39	0	6,8,10	0.56	0
2	MLU	X	10	2	7,8,9	0.13	0	7,9,11	0.92	0
2	NZC	Y	4	2	6,7,8	0.76	0	6,8,10	0.83	0
2	MP8	Y	3	2	6,8,9	0.81	0	3,10,12	1.25	0
2	MVA	X	2	2	6,7,8	0.38	0	6,8,10	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLU	Y	10	2	-	0/5/8/10	-
2	MP8	X	3	2	-	0/0/11/13	0/1/1/1
2	MVA	Y	8	2	-	0/6/8/10	-
2	MP8	X	6	2	-	0/0/11/13	0/1/1/1
2	MP8	Y	6	2	-	0/0/11/13	0/1/1/1
2	MVA	X	8	2	-	0/6/8/10	-
2	NZC	X	4	2	-	1/6/8/10	-
2	MVA	Y	2	2	-	0/6/8/10	-
2	MLU	X	10	2	-	0/5/8/10	-
2	NZC	Y	4	2	-	1/6/8/10	-
2	MP8	Y	3	2	-	0/0/11/13	0/1/1/1
2	MVA	X	2	2	-	2/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	X	2	MVA	O-C-CA-CB
2	X	4	NZC	CB-CA-N-C40
2	Y	4	NZC	CB-CA-N-C40
2	X	2	MVA	CB-CA-N-CN

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	X	3	MP8	1	0
2	Y	8	MVA	1	0
2	X	8	MVA	1	0
2	X	4	NZC	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 11 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	503	-	3,3,3	0.44	0	2,2,2	0.65	0
3	EDO	A	504	-	3,3,3	0.49	0	2,2,2	0.25	0
3	EDO	B	501	-	3,3,3	0.43	0	2,2,2	0.42	0
3	EDO	A	505	-	3,3,3	0.42	0	2,2,2	0.35	0
3	EDO	A	502	-	3,3,3	0.33	0	2,2,2	0.57	0
3	EDO	A	501	-	3,3,3	0.36	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	503	-	-	0/1/1/1	-
3	EDO	A	504	-	-	0/1/1/1	-
3	EDO	B	501	-	-	1/1/1/1	-
3	EDO	A	505	-	-	0/1/1/1	-
3	EDO	A	502	-	-	1/1/1/1	-
3	EDO	A	501	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	501	EDO	O1-C1-C2-O2
3	A	502	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/381 (97%)	0.18	27 (7%) 21 24	13, 25, 51, 89	12 (3%)
1	B	372/381 (97%)	0.24	17 (4%) 37 43	12, 27, 50, 72	8 (2%)
2	X	4/11 (36%)	-0.49	0 100 100	21, 25, 28, 32	0
2	Y	4/11 (36%)	-0.84	0 100 100	17, 18, 18, 18	0
All	All	750/784 (95%)	0.20	44 (5%) 28 32	12, 26, 51, 89	20 (2%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	LEU	7.3
1	A	59	VAL	5.4
1	A	217	ILE	5.3
1	A	123	PRO	5.3
1	B	361	ASN	5.3
1	A	23	ARG	5.2
1	B	93	GLU	5.0
1	A	212	LEU	4.4
1	A	0	HIS	4.4
1	B	360	ASN	4.4
1	A	218	ASP	4.1
1	A	283	ASP	4.0
1	B	257	LEU	3.9
1	A	83	ASP	3.8
1	A	22	GLU	3.8
1	A	85	SER	3.5
1	B	168	ASP	3.5
1	B	92	ASP	3.5
1	B	358	ASP	3.4
1	A	93	GLU	3.3
1	B	169	VAL	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	-1	HIS	3.3
1	B	155	TYR	3.1
1	A	257	LEU	3.1
1	A	21	VAL	3.1
1	B	194	ASP	3.1
1	A	285	GLY	3.0
1	B	24	ARG	2.9
1	A	216	GLU	2.9
1	A	284	ARG	2.8
1	A	24	ARG	2.7
1	B	26	THR	2.6
1	B	153	THR	2.5
1	A	282	ASN	2.5
1	A	125	GLN	2.5
1	A	258	GLY	2.5
1	A	215	GLU	2.4
1	B	313	LEU	2.4
1	B	362	ALA	2.4
1	A	194	ASP	2.3
1	A	60	ASN	2.3
1	A	98	ALA	2.2
1	A	9	GLN	2.0
1	B	357	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MVA	X	8	8/9	0.95	0.09	24,28,31,31	0
2	MLU	X	10	9/10	0.95	0.10	28,31,38,41	0
2	MVA	X	2	8/9	0.96	0.08	22,26,28,31	0
2	NZC	X	4	8/9	0.96	0.06	16,22,24,25	0
2	MP8	X	6	8/9	0.97	0.06	20,24,25,28	0
2	MP8	X	3	8/9	0.97	0.05	20,20,22,22	0
2	NZC	Y	4	8/9	0.97	0.06	15,16,17,18	0
2	MLU	Y	10	9/10	0.97	0.06	17,19,23,24	0
2	MP8	Y	3	8/9	0.98	0.05	13,15,20,21	0
2	MVA	Y	8	8/9	0.98	0.04	14,16,21,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MVA	Y	2	8/9	0.98	0.05	14,19,20,20	0
2	MP8	Y	6	8/9	0.98	0.06	17,18,23,25	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	501	4/4	0.82	0.20	52,53,64,72	0
3	EDO	A	501	4/4	0.85	0.15	41,45,49,58	0
3	EDO	A	503	4/4	0.86	0.20	28,30,46,58	0
3	EDO	A	505	4/4	0.88	0.12	41,45,46,50	0
4	IOD	A	511	1/1	0.88	0.19	89,89,89,89	1
3	EDO	A	502	4/4	0.90	0.14	47,47,47,51	0
4	IOD	A	512	1/1	0.91	0.10	36,36,36,36	1
3	EDO	A	504	4/4	0.95	0.09	28,33,38,45	0
4	IOD	B	502	1/1	0.95	0.07	45,45,45,45	1
4	IOD	B	503	1/1	0.95	0.11	61,61,61,61	1
4	IOD	B	504	1/1	0.96	0.10	46,46,46,46	1
4	IOD	A	508	1/1	0.97	0.04	35,35,35,35	1
4	IOD	A	509	1/1	0.98	0.04	28,28,28,28	1
4	IOD	A	507	1/1	0.98	0.04	39,39,39,39	1
4	IOD	A	510	1/1	0.99	0.04	37,37,37,37	1
4	IOD	A	506	1/1	1.00	0.02	27,27,27,27	1
4	IOD	B	505	1/1	1.00	0.03	23,23,23,23	1

6.5 Other polymers [i](#)

There are no such residues in this entry.