



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 09:33 PM UTC

PDB ID : 7PXP / pdb_00007pxp
Title : Benzoylsuccinyl-CoA thiolase
Authors : Ermler, U.; Heider, H.; Weidenweber, S.; Demmer, U.
Deposited on : 2021-10-08
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	
1	B	146	
1	E	146	
1	F	146	
2	C	392	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	392	88% 10% ..
2	G	392	% 91% 7% .
2	H	392	89% 8% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Benzoylsuccinyl-CoA thiolase subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	132	Total	C	N	O	S	0	0	0
			1073	692	171	202	8			
1	B	131	Total	C	N	O	S	0	0	0
			1064	687	170	199	8			
1	E	133	Total	C	N	O	S	0	0	0
			1082	698	173	203	8			
1	F	131	Total	C	N	O	S	0	0	0
			1064	687	170	199	8			

- Molecule 2 is a protein called Benzoylsuccinyl-CoA thiolase subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	384	Total	C	N	O	S	0	0	0
			2841	1767	501	542	31			
2	D	385	Total	C	N	O	S	0	0	0
			2855	1775	506	543	31			
2	G	391	Total	C	N	O	S	0	0	0
			2892	1798	512	551	31			
2	H	387	Total	C	N	O	S	0	0	0
			2868	1785	507	545	31			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	391	GLY	-	expression tag	UNP Q39VG1
C	392	SER	-	expression tag	UNP Q39VG1
D	391	GLY	-	expression tag	UNP Q39VG1
D	392	SER	-	expression tag	UNP Q39VG1
G	391	GLY	-	expression tag	UNP Q39VG1
G	392	SER	-	expression tag	UNP Q39VG1
H	391	GLY	-	expression tag	UNP Q39VG1
H	392	SER	-	expression tag	UNP Q39VG1

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	B	1	Total Zn 1 1	0	0
3	E	1	Total Zn 1 1	0	0
3	F	1	Total Zn 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	10	Total O 10 10	0	0
4	B	16	Total O 16 16	0	0
4	C	55	Total O 55 55	0	0
4	D	36	Total O 36 36	0	0
4	E	17	Total O 17 17	0	0
4	F	17	Total O 18 18	0	1
4	G	90	Total O 91 91	0	1
4	H	90	Total O 90 90	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Benzoylsuccinyl-CoA thiolase subunit

Chain A: 




- Molecule 1: Benzoylsuccinyl-CoA thiolase subunit

Chain B: 



- Molecule 1: Benzoylsuccinyl-CoA thiolase subunit

Chain E: 




- Molecule 1: Benzoylsuccinyl-CoA thiolase subunit

Chain F: 




- Molecule 2: Benzoylsuccinyl-CoA thiolase subunit


Chain C: 




- Molecule 2: Benzoylsuccinyl-CoA thiolase subunit



- Molecule 2: Benzoylsuccinyl-CoA thiolase subunit



- Molecule 2: Benzoylsuccinyl-CoA thiolase subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.79Å 105.51Å 142.54Å 90.00° 92.27° 90.00°	Depositor
Resolution (Å)	48.41 – 2.00 48.41 – 2.00	Depositor EDS
% Data completeness (in resolution range)	93.3 (48.41-2.00) 83.6 (48.41-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.211 , 0.233 0.217 , 0.238	Depositor DCC
R_{free} test set	7666 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.513	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 25.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.078 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16076	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/1100	1.11	3/1488 (0.2%)
1	B	0.73	0/1091	1.12	5/1475 (0.3%)
1	E	0.76	0/1109	1.12	3/1499 (0.2%)
1	F	0.77	0/1091	1.12	2/1475 (0.1%)
2	C	0.86	0/2889	1.29	7/3898 (0.2%)
2	D	0.82	0/2903	1.33	13/3916 (0.3%)
2	G	0.97	2/2942 (0.1%)	1.28	9/3970 (0.2%)
2	H	0.99	7/2917 (0.2%)	1.25	5/3935 (0.1%)
All	All	0.87	9/16042 (0.1%)	1.24	47/21656 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	71	MET	SD-CE	-15.48	1.40	1.79
2	H	223	MET	SD-CE	-12.65	1.48	1.79
2	H	71	MET	SD-CE	-11.55	1.50	1.79
2	H	146	MET	SD-CE	-10.07	1.54	1.79
2	H	200	MET	SD-CE	-8.64	1.57	1.79

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	42	ASP	CA-CB-CG	8.01	120.61	112.60
2	G	268	SER	N-CA-C	-7.55	97.28	109.96
1	B	31	ASP	CA-CB-CG	7.41	120.01	112.60
2	C	357	VAL	CA-C-N	7.25	133.26	121.87
2	C	357	VAL	C-N-CA	7.25	133.26	121.87

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	82	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1073	0	1044	4	0
1	B	1064	0	1039	2	0
1	E	1082	0	1057	4	0
1	F	1064	0	1039	5	0
2	C	2841	0	2824	16	0
2	D	2855	0	2840	16	0
2	G	2892	0	2873	19	0
2	H	2868	0	2851	27	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	10	0	0	1	0
4	B	16	0	0	0	0
4	C	55	0	0	1	0
4	D	36	0	0	0	0
4	E	17	0	0	0	0
4	F	18	0	0	0	0
4	G	91	0	0	0	0
4	H	90	0	0	0	0
All	All	16076	0	15567	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:36:MET:HE1	2:H:71:MET:HE1	1.41	0.98
2:G:36:MET:HE1	2:G:71:MET:HE1	1.49	0.92
2:H:200:MET:HE3	2:H:284:PHE:HB3	1.54	0.90
2:H:106:LEU:HD22	2:H:222:MET:HE2	1.53	0.90
2:H:200:MET:CE	2:H:284:PHE:HB3	2.07	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	130/146 (89%)	128 (98%)	2 (2%)	0	100	100
1	B	129/146 (88%)	126 (98%)	3 (2%)	0	100	100
1	E	131/146 (90%)	129 (98%)	2 (2%)	0	100	100
1	F	129/146 (88%)	128 (99%)	1 (1%)	0	100	100
2	C	380/392 (97%)	368 (97%)	11 (3%)	1 (0%)	36	35
2	D	381/392 (97%)	364 (96%)	15 (4%)	2 (0%)	24	21
2	G	389/392 (99%)	373 (96%)	15 (4%)	1 (0%)	36	35
2	H	383/392 (98%)	370 (97%)	12 (3%)	1 (0%)	36	35
All	All	2052/2152 (95%)	1986 (97%)	61 (3%)	5 (0%)	43	42

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	358	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	358	GLY
2	H	353	GLY
2	D	353	GLY
2	G	353	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/132 (90%)	114 (96%)	5 (4%)	26	25
1	B	118/132 (89%)	112 (95%)	6 (5%)	21	19
1	E	120/132 (91%)	116 (97%)	4 (3%)	33	34
1	F	118/132 (89%)	114 (97%)	4 (3%)	32	33
2	C	297/302 (98%)	292 (98%)	5 (2%)	53	60
2	D	298/302 (99%)	287 (96%)	11 (4%)	30	30
2	G	301/302 (100%)	295 (98%)	6 (2%)	48	54
2	H	299/302 (99%)	294 (98%)	5 (2%)	53	60
All	All	1670/1736 (96%)	1624 (97%)	46 (3%)	38	41

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	54	MET
2	G	3	LEU
1	E	66	VAL
1	F	54	MET
2	G	42	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	57	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	115	HIS
2	H	367	HIS
2	D	118	HIS
2	D	153	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	132/146 (90%)	-0.19	0 100 100	36, 52, 76, 102	0
1	B	131/146 (89%)	-0.29	0 100 100	30, 45, 65, 95	0
1	E	133/146 (91%)	-0.31	1 (0%) 82 82	31, 43, 67, 98	0
1	F	131/146 (89%)	-0.45	0 100 100	28, 42, 64, 81	0
2	C	384/392 (97%)	-0.49	1 (0%) 90 89	27, 38, 54, 74	0
2	D	385/392 (98%)	-0.27	1 (0%) 90 89	30, 46, 65, 93	0
2	G	391/392 (99%)	-0.67	4 (1%) 79 79	20, 30, 48, 82	0
2	H	387/392 (98%)	-0.70	0 100 100	22, 33, 46, 87	0
All	All	2074/2152 (96%)	-0.48	7 (0%) 90 89	20, 38, 63, 102	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	124	ALA	4.3
1	E	146	ALA	2.7
2	G	127	ALA	2.7
2	G	123	ALA	2.3
2	D	391	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	A	201	1/1	1.00	0.02	56,56,56,56	0
3	ZN	B	201	1/1	1.00	0.01	31,31,31,31	0
3	ZN	E	201	1/1	1.00	0.01	36,36,36,36	0
3	ZN	F	201	1/1	1.00	0.01	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 201:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN E 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.