



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 10:43 AM EDT

PDB ID : 2RTC / pdb\_00002rtc  
Title : APOSTREPTAVIDIN, PH 3.60, SPACE GROUP I222  
Authors : Katz, B.A.  
Deposited on : 1997-09-11  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	135	
1	D	135	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4197 atoms, of which 2073 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STREPTAVIDIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	123	Total	C	H	N	O	45	13	0
			1943	628	939	171	205			
1	D	121	Total	C	H	N	O	73	13	0
			1902	613	918	169	202			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	1
			5	4	1		
2	B	1	Total	O	S	0	1
			5	4	1		
2	D	1	Total	O	S	0	1
			5	4	1		
2	D	1	Total	O	S	0	1
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	S	0	1
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	53	Total	H	O	0	8
			155	102	53		
3	D	58	Total	H	O	0	2
			172	114	58		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: STREPTAVIDIN



#### • Molecule 1: STREPTAVIDIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.18Å 105.39Å 47.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.50 – 1.50 7.50 – 1.50	Depositor EDS
% Data completeness (in resolution range)	60.0 (7.50-1.50) 79.3 (7.50-1.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 1.32Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.210 , 0.265 0.234 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.4	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.22 , 64.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4197	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	1.88	16/1043 (1.5%)	1.89	39/1428 (2.7%)
1	D	1.87	12/1021 (1.2%)	1.91	36/1399 (2.6%)
All	All	1.88	28/2064 (1.4%)	1.90	75/2827 (2.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3
1	D	0	3
All	All	0	6

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	127	HIS	CD2-NE2	-6.92	1.30	1.37
1	B	92	TRP	NE1-CE2	-6.66	1.30	1.37
1	B	127	HIS	CG-ND1	-6.64	1.30	1.38
1	B	73	LEU	CA-CB	6.45	1.63	1.53
1	D	127	HIS	CG-ND1	-6.39	1.31	1.38
1	D	87[A]	HIS	CD2-NE2	-6.31	1.30	1.37
1	D	87[B]	HIS	CD2-NE2	-6.31	1.30	1.37
1	D	87[A]	HIS	CG-ND1	-6.26	1.31	1.38
1	D	87[B]	HIS	CG-ND1	-6.26	1.31	1.38
1	D	61[A]	ASP	CG-OD2	-6.15	1.21	1.33
1	D	61[B]	ASP	CG-OD2	-6.15	1.21	1.33
1	B	134	LYS	N-CA	6.11	1.54	1.45
1	B	87[A]	HIS	CG-ND1	-6.09	1.31	1.38
1	B	87[B]	HIS	CG-ND1	-6.09	1.31	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	87[A]	HIS	CD2-NE2	-5.89	1.31	1.37
1	B	87[B]	HIS	CD2-NE2	-5.89	1.31	1.37
1	B	108	TRP	NE1-CE2	-5.85	1.31	1.37
1	D	45	SER	CA-CB	5.70	1.62	1.53
1	D	127	HIS	CD2-NE2	-5.64	1.31	1.37
1	B	61[A]	ASP	CG-OD2	-5.61	1.22	1.33
1	B	61[B]	ASP	CG-OD2	-5.61	1.22	1.33
1	B	29	PHE	CA-C	5.27	1.58	1.52
1	B	96	TYR	CA-CB	5.22	1.60	1.53
1	D	87[A]	HIS	N-CA	5.07	1.54	1.46
1	D	87[B]	HIS	N-CA	5.07	1.54	1.46
1	D	79	TRP	CA-CB	5.06	1.58	1.52
1	B	75	TRP	NE1-CE2	-5.04	1.31	1.37
1	B	79	TRP	CG-CD2	-5.02	1.34	1.43

All (75) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	78	ALA	N-CA-C	-10.25	92.95	109.76
1	B	78	ALA	N-CA-C	-10.03	92.92	109.07
1	D	123	THR	N-CA-C	9.11	123.37	108.52
1	D	115	THR	N-CA-C	-9.07	97.44	110.59
1	D	29	PHE	CA-CB-CG	9.03	122.83	113.80
1	D	107[A]	GLN	N-CA-C	-8.68	94.75	109.24
1	D	107[B]	GLN	N-CA-C	-8.68	94.75	109.24
1	D	95	GLN	OE1-CD-NE2	-8.46	114.14	122.60
1	B	128	ASP	CA-CB-CG	7.87	120.47	112.60
1	B	123	THR	N-CA-C	7.78	121.58	108.90
1	B	104	ILE	N-CA-C	-7.34	97.03	107.75
1	B	107	GLN	N-CA-C	-7.27	97.09	109.24
1	B	107	GLN	CA-CB-CG	-7.18	99.74	114.10
1	B	113	GLY	N-CA-C	-7.12	100.46	111.64
1	B	108	TRP	CG-CD2-CE3	-6.94	126.96	133.90
1	D	49	ASN	N-CA-C	-6.91	101.07	110.68
1	B	134	LYS	O-C-N	6.69	127.22	121.20
1	B	49	ASN	N-CA-C	-6.55	100.47	109.71
1	D	92	TRP	CG-CD1-NE1	-6.53	101.71	110.20
1	B	92	TRP	CG-CD1-NE1	-6.52	101.72	110.20
1	B	115	THR	N-CA-C	-6.49	101.79	110.55
1	D	23	ASN	OD1-CG-ND2	-6.39	116.21	122.60
1	D	108	TRP	CG-CD1-NE1	-6.38	101.91	110.20
1	B	95	GLN	OE1-CD-NE2	-6.38	116.22	122.60

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	TRP	NE1-CE2-CZ2	6.34	139.60	130.10
1	B	75	TRP	CG-CD1-NE1	-6.33	101.97	110.20
1	D	75	TRP	CG-CD1-NE1	-6.31	102.00	110.20
1	D	81	ASN	CA-CB-CG	6.16	118.76	112.60
1	B	108	TRP	CB-CG-CD2	6.10	135.34	126.80
1	D	31	VAL	CB-CA-C	-6.02	101.40	110.55
1	B	85	ASN	OD1-CG-ND2	-5.95	116.66	122.60
1	B	114	THR	OG1-CB-CG2	-5.94	97.41	109.30
1	D	106	THR	N-CA-C	5.88	118.64	109.52
1	D	83	TYR	CA-CB-CG	-5.86	103.35	113.90
1	B	21	TRP	NE1-CE2-CZ2	5.85	138.87	130.10
1	B	79	TRP	CG-CD1-NE1	-5.85	102.60	110.20
1	D	120	TRP	CG-CD1-NE1	-5.77	102.70	110.20
1	B	120	TRP	NE1-CE2-CZ2	5.74	138.70	130.10
1	D	92	TRP	NE1-CE2-CZ2	5.70	138.65	130.10
1	D	75	TRP	NE1-CE2-CZ2	5.66	138.59	130.10
1	D	120	TRP	NE1-CE2-CZ2	5.61	138.52	130.10
1	B	108	TRP	NE1-CE2-CD2	-5.56	100.17	107.40
1	B	30	ILE	N-CA-C	-5.54	99.77	107.80
1	D	26	GLY	N-CA-C	-5.54	107.36	115.72
1	B	57	THR	CA-CB-OG1	-5.53	101.31	109.60
1	D	44	GLU	N-CA-C	-5.53	99.40	108.41
1	B	120	TRP	CG-CD1-NE1	-5.52	103.02	110.20
1	B	85	ASN	CA-CB-CG	-5.50	107.09	112.60
1	D	21	TRP	CG-CD1-NE1	-5.47	103.08	110.20
1	B	81	ASN	CA-CB-CG	5.45	118.05	112.60
1	D	92	TRP	CD1-NE1-CE2	5.43	118.68	108.90
1	D	129	THR	CA-CB-OG1	-5.40	101.49	109.60
1	D	21	TRP	NE1-CE2-CZ2	5.40	138.20	130.10
1	B	106	THR	N-CA-C	5.39	117.18	109.14
1	B	21	TRP	CG-CD1-NE1	-5.34	103.26	110.20
1	B	82	ASN	OD1-CG-ND2	-5.31	117.29	122.60
1	B	29	PHE	N-CA-CB	-5.30	102.45	110.77
1	D	20	THR	CA-CB-OG1	-5.29	101.67	109.60
1	D	108	TRP	NE1-CE2-CZ2	5.28	138.02	130.10
1	D	128	ASP	CA-CB-CG	5.26	117.86	112.60
1	D	108	TRP	CD1-NE1-CE2	5.19	118.24	108.90
1	B	129	THR	CA-CB-OG1	-5.17	101.84	109.60
1	B	21	TRP	NE1-CE2-CD2	-5.15	100.71	107.40
1	B	108	TRP	CB-CG-CD1	-5.12	119.22	126.90
1	D	46	ALA	N-CA-C	-5.11	107.08	113.72
1	B	83	TYR	CA-CB-CG	-5.10	104.72	113.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	120	TRP	NE1-CE2-CD2	-5.08	100.80	107.40
1	D	79	TRP	CG-CD1-NE1	-5.07	103.61	110.20
1	B	92	TRP	CD1-NE1-CE2	5.07	118.02	108.90
1	B	108	TRP	CD1-NE1-CE2	5.06	118.00	108.90
1	D	79	TRP	NE1-CE2-CZ2	5.05	137.68	130.10
1	D	75	TRP	CD1-NE1-CE2	5.05	117.98	108.90
1	B	79	TRP	NE1-CE2-CZ2	5.03	137.65	130.10
1	B	21	TRP	CD1-NE1-CE2	5.02	117.94	108.90
1	D	75	TRP	NE1-CE2-CD2	-5.00	100.89	107.40

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	103	ARG	Sidechain
1	B	53	ARG	Sidechain
1	B	63[A]	ALA	Peptide
1	D	103	ARG	Sidechain
1	D	53	ARG	Sidechain
1	D	63[A]	ALA	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1004	939	927	15	0
1	D	984	918	906	19	0
2	B	10	0	0	0	0
2	D	15	0	0	0	0
3	B	53	102	0	0	4
3	D	58	114	0	1	3
All	All	2124	2073	1833	28	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22[B]:TYR:CE1	1:B:131:THR:HB	2.21	0.76
1:B:114:THR:H	1:D:95:GLN:HE22	1.32	0.75
1:B:95:GLN:HE22	1:D:114:THR:H	1.37	0.71
1:D:51:GLU:O	1:D:52:SER:HB2	2.01	0.60
1:B:51:GLU:O	1:B:52:SER:CB	2.52	0.58
1:B:114:THR:H	1:D:95:GLN:NE2	2.04	0.52
1:B:51:GLU:O	1:B:52:SER:HB3	2.12	0.49
1:D:24:GLN:HG2	1:D:25:LEU:N	2.28	0.49
1:D:110:LEU:C	1:D:110:LEU:HD23	2.38	0.49
1:D:51:GLU:O	1:D:52:SER:CB	2.61	0.49
1:B:22[A]:TYR:CE1	1:B:28:THR:HG23	2.48	0.48
1:D:51:GLU:HB2	1:D:84:ARG:NE	2.28	0.48
1:B:114:THR:N	1:D:95:GLN:HE22	2.05	0.47
1:D:23:ASN:HB3	1:D:130:PHE:CE1	2.49	0.47
1:B:61[A]:ASP:OD2	1:D:85:ASN:OD1	2.33	0.47
1:D:13:ALA:N	3:D:1174:HOH:O	2.48	0.47
1:B:133:VAL:HG22	1:B:134:LYS:HG3	1.97	0.45
1:B:115:THR:HA	1:D:68[B]:GLY:O	2.18	0.44
1:D:120:TRP:CZ2	1:D:121:LYS:HD2	2.53	0.43
1:D:23:ASN:OD1	1:D:23:ASN:C	2.60	0.43
1:D:30:ILE:O	1:D:41:GLY:HA3	2.18	0.42
1:B:37:GLY:HA2	1:B:60[B]:TYR:CE1	2.55	0.41
1:B:37:GLY:HA2	1:B:60[A]:TYR:CE1	2.56	0.41
1:B:19:GLY:HA2	1:B:135:PRO:HA	2.02	0.41
1:B:39:LEU:HD13	1:B:60[B]:TYR:CD1	2.56	0.41
1:D:21:TRP:CZ2	1:D:132:LYS:HE3	2.56	0.41
1:D:121:LYS:H	1:D:121:LYS:HG2	1.73	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:491[C]:HOH:O	3:B:491[C]:HOH:O[3_655]	0.51	1.69
3:D:1465[C]:HOH:O	3:D:1465[C]:HOH:O[3_556]	0.82	1.38
3:B:491[C]:HOH:O	3:D:1465[C]:HOH:O[8_555]	1.39	0.81
3:B:1460[C]:HOH:O	3:B:1460[C]:HOH:O[3_655]	1.42	0.78
3:B:1460[C]:HOH:O	3:D:1465[C]:HOH:O[8_555]	1.99	0.21

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	134/135 (99%)	123 (92%)	8 (6%)	3 (2%)	5	0
1	D	132/135 (98%)	127 (96%)	2 (2%)	3 (2%)	5	0
All	All	266/270 (98%)	250 (94%)	10 (4%)	6 (2%)	5	0

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	52	SER
1	D	15	ALA
1	D	52	SER
1	D	16	GLY
1	B	64[A]	PRO
1	B	64[B]	PRO

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	100/100 (100%)	95 (95%)	5 (5%)	22	3
1	D	98/100 (98%)	89 (91%)	9 (9%)	8	0
All	All	198/200 (99%)	184 (93%)	14 (7%)	16	1

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	39	LEU
1	B	62[A]	SER
1	B	62[B]	SER
1	B	73	LEU
1	B	133	VAL
1	D	31	VAL
1	D	36	ASP
1	D	39	LEU
1	D	52	SER
1	D	53	ARG
1	D	73[A]	LEU
1	D	73[B]	LEU
1	D	107[A]	GLN
1	D	107[B]	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN
1	B	107	GLN
1	D	95	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	403[A]	-	4,4,4	0.54	0	6,6,6	0.29	0
2	SO4	D	401[B]	-	4,4,4	0.52	0	6,6,6	0.18	0
2	SO4	D	402[A]	-	4,4,4	0.65	0	6,6,6	0.38	0
2	SO4	D	400[A]	-	4,4,4	0.38	0	6,6,6	0.25	0
2	SO4	B	404[A]	-	4,4,4	0.52	0	6,6,6	0.17	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	B	120/135 (88%)	0.08	9 (7%)	20 22	10, 23, 41, 56	24 (20%)
1	D	117/135 (86%)	0.10	5 (4%)	40 44	1, 26, 56, 71	20 (17%)
All	All	237/270 (87%)	0.09	14 (5%)	28 30	1, 24, 51, 71	44 (18%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	ALA	9.9
1	B	63[A]	ALA	7.0
1	D	65[A]	ALA	5.2
1	B	66[A]	THR	4.7
1	D	68[A]	GLY	3.9
1	B	68[A]	GLY	3.9
1	B	62[A]	SER	3.7
1	B	100	ALA	3.1
1	D	13	ALA	3.0
1	D	83	TYR	2.7
1	B	67[A]	ASP	2.4
1	D	62[A]	SER	2.3
1	B	135	PRO	2.1
1	B	99	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	D	400[A]	5/5	0.49	0.15	65,65,67,67	5
2	SO4	B	404[A]	5/5	0.73	0.20	55,56,58,59	5
2	SO4	D	401[B]	5/5	0.73	0.14	69,70,72,72	5
2	SO4	B	403[A]	5/5	0.78	0.17	61,62,63,65	5
2	SO4	D	402[A]	5/5	0.85	0.16	57,57,58,60	5

## 6.5 Other polymers

There are no such residues in this entry.