



wwPDB EM Validation Summary Report ⓘ

Sep 19, 2026 – 07:18 am BST

PDB ID : 9T35 / pdb_00009t35
EMDB ID : EMD-55484
Title : Cryo-EM structure of homo-hexameric hLRRC8A L402W
Authors : Bertelli, S.; Wang, L.; Klussendorf, M.; Pusch, M.; Dutzler, R.; Stauber, T.
Deposited on : 2025-10-24
Resolution : 2.78 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10754 (2.28 - 3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	811	
1	B	811	
1	C	811	
1	D	811	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	811	
1	F	811	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 15342 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Volume-regulated anion channel subunit LRRC8A.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	305	Total	C	N	O	S	0	0
			2557	1694	406	440	17		
1	B	305	Total	C	N	O	S	0	0
			2557	1694	406	440	17		
1	C	305	Total	C	N	O	S	0	0
			2557	1694	406	440	17		
1	D	305	Total	C	N	O	S	0	0
			2557	1694	406	440	17		
1	E	305	Total	C	N	O	S	0	0
			2557	1694	406	440	17		
1	F	305	Total	C	N	O	S	0	0
			2557	1694	406	440	17		


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q8IWT6
A	402	TRP	LEU	engineered mutation	UNP Q8IWT6
A	811	ALA	-	expression tag	UNP Q8IWT6
B	1	SER	-	expression tag	UNP Q8IWT6
B	402	TRP	LEU	engineered mutation	UNP Q8IWT6
B	811	ALA	-	expression tag	UNP Q8IWT6
C	1	SER	-	expression tag	UNP Q8IWT6
C	402	TRP	LEU	engineered mutation	UNP Q8IWT6
C	811	ALA	-	expression tag	UNP Q8IWT6
D	1	SER	-	expression tag	UNP Q8IWT6
D	402	TRP	LEU	engineered mutation	UNP Q8IWT6
D	811	ALA	-	expression tag	UNP Q8IWT6
E	1	SER	-	expression tag	UNP Q8IWT6
E	402	TRP	LEU	engineered mutation	UNP Q8IWT6
E	811	ALA	-	expression tag	UNP Q8IWT6
F	1	SER	-	expression tag	UNP Q8IWT6
F	402	TRP	LEU	engineered mutation	UNP Q8IWT6
F	811	ALA	-	expression tag	UNP Q8IWT6






- Molecule 1: Volume-regulated anion channel subunit LRRC8A



- Molecule 1: Volume-regulated anion channel subunit LRRC8A



THR	G93	D102	R103	H104	Q105	D110	V129	R148	I159	L160	F164	D165	S166	P167	W168	R171	A172	L173	S174	E175	T176	V177	V178	E179	GLU	SER	ASP	PRO	LYS	PRO	ALA	PHE	SER	LYS	MET	ASN	ASN	GLY	SER	MET	ASP	LYS	LYS	SER	THR	VAL	SER	GLU	ASP	VAL	GLU	ALA
THR	VAL	PRO	MET	GLN	ARG	LYS	SER	LYS	SER	ARG	ILE	VAL	ASP	ARG	GLU	THR	G230	V231	L232	D233	K234	K235	E236	G237	E238	Q239	E245	K246	R251	G257	D258	I259	F275	L278	Y281	R306	R309	F353	E354	S355	I356	R357	E358	S359	S360							
SER	TYR	D364	V368	F372	L385	Y386	R389	F390	L394	S395	E396	VAL	SER	SER	GLU	ASN	TRP	ARG	GLN	ASN	LEU	ASN	ASN	TRP	THR	LEU	ASP	LYS	LEU	ARG	GLN	ALA	THR	LYS	ASN	GLU	LEU	HIS	LEU	PHE	MET	LEU	SER	GLY	ILE							
PRO	ASP	THR	VAL	PHE	ASP	LEU	VAL	GLU	GLU	PRO	ILE	VAL	VAL	ASP	THR	ILE	PRO	PRO	ILE	GLN	ALA	THR	GLY	LEU	LYS	GLU	LEU	TRP	ILE	THR	ALA	ALA	ALA	PRO	ALA	ALA	PHE	LEU	ARG	GLU	GLU	ASN	ARG	ALA	LEU							
HIS	ILE	PHE	THR	ASP	ILE	GLU	TRP	ILE	TYR	SER	THR	GLU	GLU	GLU	GLU	ILE	HIS	THR	THR	ASN	LEU	THR	ASN	GLY	ARG	TYR	ILE	VAL	ILE	GLY	ARG	ALA	VAL	VAL	ARG	LEU	LYS	ASN	LEU	SER	LEU	THR	LEU	LEU	LEU	LEU						
PRO	GLN	VAL	THR	ASP	VAL	VAL	HIS	VAL	LEU	ASN	GLN	GLY	GLY	THR	LYS	LEU	VAL	ASN	SER	LEU	LYS	ASN	ALA	ASN	LEU	LEU	THR	GLU	GLU	ILE	THR	CYS	ASP	GLY	ARG	GLU	THR	ASN	GLU	THR	ASN	GLU	THR	GLN	GLU							
ILE	ASP	LYS	ASN	ASN	LEU	LYS	THR	LYS	ILE	PHE	GLN	LEU	HIS	THR	ARG	THR	THR	THR	LEU	TRP	ASN	HIS	ILE	ALA	ALA	TYR	ILE	PRO	ILE	GLN	ILE	GLY	ASN	LEU	ARG	LEU	THR	ASN	ASN	ARG	ASN	ILE	GLU	LYS	ILE	PRO						
THR	GLN	PHE	TYR	CYS	ARG	LYS	LEU	ARG	HIS	ASN	ASN	THR	THR	PHE	LEU	PRO	ALA	ILE	GLY	GLN	ASN	LEU	GLN	ASN	LEU	LEU	ALA	THR	THR	ALA	THR	GLU	PHE	GLN	CYS	ARG	LYS	LEU	ARG	ALA	PRO	HIS	LEU	GLU	VAL							
GLY	ASN	VAL	LEU	GLN	SER	PRO	SER	GLU	LEU	THR	THR	ASN	ILE	ILE	GLU	LEU	ARG	GLY	ARG	GLY	LEU	GLU	GLU	GLU	GLU	GLU	GLY	VAL	VAL	GLU	GLU	ASP	LEU	PHE	ASN	ASN	ASN	THR	LEU	PRO	PRO	GLU	VAL									
LYS	GLU	ARG	TRP	ARG	ALA	ASP	GLU	LYS	GLU	ALA	ALA	ILE	ILE	ILE	GLU	CYS	LEU	LEU	LEU	CYS	LEU	PRO	VAL	GLU	GLY	LEU	GLY	VAL	VAL	GLU	GLU	GLU	LEU	ASN	ASN	THR	LEU	ALA	PRO	GLU	GLU	VAL										

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	42551	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60.61	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOCONTINUUM (6k x 4k)	Depositor
Maximum map value	1.278	Depositor
Minimum map value	-0.476	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.035	Depositor
Recommended contour level	0.34	Depositor
Map size (Å)	333.312, 333.312, 333.312	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.302, 1.302, 1.302	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.08	0/2630	0.20	0/3561
1	B	0.08	0/2630	0.19	0/3561
1	C	0.08	0/2630	0.20	0/3561
1	D	0.08	0/2630	0.19	0/3561
1	E	0.08	0/2630	0.19	0/3561
1	F	0.08	0/2630	0.20	0/3561
All	All	0.08	0/15780	0.19	0/21366

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2557	0	2553	15	0
1	B	2557	0	2553	13	0
1	C	2557	0	2553	12	0
1	D	2557	0	2553	14	0
1	E	2557	0	2553	21	0
1	F	2557	0	2553	19	0
All	All	15342	0	15318	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:ARG:HD2	1:A:259:ILE:HG12	1.88	0.56
1:C:148:ARG:HD2	1:C:259:ILE:HG12	1.88	0.55
1:F:148:ARG:HD2	1:F:259:ILE:HG12	1.88	0.55
1:B:148:ARG:HD2	1:B:259:ILE:HG12	1.88	0.55
1:D:148:ARG:HD2	1:D:259:ILE:HG12	1.88	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/811 (37%)	289 (97%)	8 (3%)	0	100	100
1	B	297/811 (37%)	289 (97%)	8 (3%)	0	100	100
1	C	297/811 (37%)	289 (97%)	8 (3%)	0	100	100
1	D	297/811 (37%)	289 (97%)	8 (3%)	0	100	100
1	E	297/811 (37%)	289 (97%)	8 (3%)	0	100	100
1	F	297/811 (37%)	289 (97%)	8 (3%)	0	100	100
All	All	1782/4866 (37%)	1734 (97%)	48 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/749 (38%)	276 (98%)	7 (2%)	42	73
1	B	283/749 (38%)	276 (98%)	7 (2%)	42	73
1	C	283/749 (38%)	276 (98%)	7 (2%)	42	73
1	D	283/749 (38%)	276 (98%)	7 (2%)	42	73
1	E	283/749 (38%)	276 (98%)	7 (2%)	42	73
1	F	283/749 (38%)	276 (98%)	7 (2%)	42	73
All	All	1698/4494 (38%)	1656 (98%)	42 (2%)	42	73

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	25	ASP
1	F	25	ASP
1	E	49	GLN
1	E	173	LEU
1	F	51	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	105	GLN
1	F	239	GLN
1	C	105	GLN
1	C	49	GLN
1	F	288	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-55484. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 127




Y Index: 142




Z Index: 142


6.3.2 Raw map



X Index: 127



Y Index: 142




Z Index: 118


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.34. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 67 nm³; this corresponds to an approximate mass of 61 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.360 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.360 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.78	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.45	3.89	3.52

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.45 differs from the reported value 2.78 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-55484 and PDB model 9T35. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.34 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.34).

9.4 Atom inclusion [i](#)



At the recommended contour level, 72% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.34) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6320	0.5150
A	0.6320	0.5160
B	0.6300	0.5160
C	0.6330	0.5150
D	0.6350	0.5120
E	0.6330	0.5160
F	0.6310	0.5140

