



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:49 AM UTC

PDB ID : 4TKZ / pdb_00004tkz
Title : Crystal structure of phosphotransferase system component EIIA from *Streptococcus agalactiae*
Authors : Nakamichi, Y.; Maruyama, Y.; Oiki, S.; Mikami, B.; Murata, K.; Hashimoto, W.
Deposited on : 2014-05-28
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	 78% 8% 14%
1	B	152	 72% 12% 14%
1	C	152	 78% 7% 14%
1	D	152	 75% 9% 14%
1	E	152	 74% 11% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	152	 79%6%14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein gbs1890.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	130	Total	C	N	O	S	0	1	0
			1002	642	161	193	6			
1	B	130	Total	C	N	O	S	0	2	0
			1007	644	162	195	6			
1	C	130	Total	C	N	O	S	0	2	0
			1015	648	167	194	6			
1	D	130	Total	C	N	O	S	0	0	0
			994	636	160	192	6			
1	E	130	Total	C	N	O	S	0	1	0
			1002	641	161	193	7			
1	F	130	Total	C	N	O	S	0	3	0
			1018	650	165	196	7			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	145	LEU	-	expression tag	UNP Q8E371
A	146	GLU	-	expression tag	UNP Q8E371
A	147	HIS	-	expression tag	UNP Q8E371
A	148	HIS	-	expression tag	UNP Q8E371
A	149	HIS	-	expression tag	UNP Q8E371
A	150	HIS	-	expression tag	UNP Q8E371
A	151	HIS	-	expression tag	UNP Q8E371
A	152	HIS	-	expression tag	UNP Q8E371
B	145	LEU	-	expression tag	UNP Q8E371
B	146	GLU	-	expression tag	UNP Q8E371
B	147	HIS	-	expression tag	UNP Q8E371
B	148	HIS	-	expression tag	UNP Q8E371
B	149	HIS	-	expression tag	UNP Q8E371
B	150	HIS	-	expression tag	UNP Q8E371
B	151	HIS	-	expression tag	UNP Q8E371
B	152	HIS	-	expression tag	UNP Q8E371
C	145	LEU	-	expression tag	UNP Q8E371

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	146	GLU	-	expression tag	UNP Q8E371
C	147	HIS	-	expression tag	UNP Q8E371
C	148	HIS	-	expression tag	UNP Q8E371
C	149	HIS	-	expression tag	UNP Q8E371
C	150	HIS	-	expression tag	UNP Q8E371
C	151	HIS	-	expression tag	UNP Q8E371
C	152	HIS	-	expression tag	UNP Q8E371
D	145	LEU	-	expression tag	UNP Q8E371
D	146	GLU	-	expression tag	UNP Q8E371
D	147	HIS	-	expression tag	UNP Q8E371
D	148	HIS	-	expression tag	UNP Q8E371
D	149	HIS	-	expression tag	UNP Q8E371
D	150	HIS	-	expression tag	UNP Q8E371
D	151	HIS	-	expression tag	UNP Q8E371
D	152	HIS	-	expression tag	UNP Q8E371
E	145	LEU	-	expression tag	UNP Q8E371
E	146	GLU	-	expression tag	UNP Q8E371
E	147	HIS	-	expression tag	UNP Q8E371
E	148	HIS	-	expression tag	UNP Q8E371
E	149	HIS	-	expression tag	UNP Q8E371
E	150	HIS	-	expression tag	UNP Q8E371
E	151	HIS	-	expression tag	UNP Q8E371
E	152	HIS	-	expression tag	UNP Q8E371
F	145	LEU	-	expression tag	UNP Q8E371
F	146	GLU	-	expression tag	UNP Q8E371
F	147	HIS	-	expression tag	UNP Q8E371
F	148	HIS	-	expression tag	UNP Q8E371
F	149	HIS	-	expression tag	UNP Q8E371
F	150	HIS	-	expression tag	UNP Q8E371
F	151	HIS	-	expression tag	UNP Q8E371
F	152	HIS	-	expression tag	UNP Q8E371

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	51	Total	O	0	0
			51	51		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	48	Total 48	O 48	0	0
3	C	51	Total 51	O 51	0	0
3	D	41	Total 41	O 41	0	0
3	E	27	Total 27	O 27	0	0
3	F	47	Total 47	O 47	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein gbs1890

Chain A: 



- Molecule 1: Putative uncharacterized protein gbs1890

Chain B: 



- Molecule 1: Putative uncharacterized protein gbs1890

Chain C: 



- Molecule 1: Putative uncharacterized protein gbs1890

Chain D: 



- Molecule 1: Putative uncharacterized protein gbs1890

Chain E: 



- Molecule 1: Putative uncharacterized protein gbs1890


Chain F:

79%

6%

•

14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	52.29Å 53.80Å 94.89Å 91.14° 90.02° 60.97°	Depositor
Resolution (Å)	33.78 – 1.80 33.78 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.7 (33.78-1.80) 95.7 (33.78-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.206 , 0.242 0.206 , 0.241	Depositor DCC
R_{free} test set	4026 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtriage
Anisotropy	0.830	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 31.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for k,-h+k,l 0.013 for h-k,h,l 0.105 for -h+k,-h,l 0.105 for -k,h-k,l 0.438 for h,h-k,-l 0.117 for -h+k,k,-l 0.019 for -h,-k,l 0.015 for k,h,-l 0.099 for -k,-h,-l 0.018 for -h,-h+k,-l 0.017 for h-k,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6369	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1015	0.81	0/1373
1	B	0.44	0/1020	0.79	0/1380
1	C	0.41	0/1029	0.81	0/1391
1	D	0.40	0/1007	0.79	0/1362
1	E	0.42	0/1015	0.78	0/1372
1	F	0.42	0/1032	0.78	0/1395
All	All	0.42	0/6118	0.79	0/8273

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1002	0	1030	7	0
1	B	1007	0	1032	12	0
1	C	1015	0	1038	6	0
1	D	994	0	1020	9	0
1	E	1002	0	1030	11	0
1	F	1018	0	1040	5	0
2	A	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	12	0	16	3	0
2	C	6	0	8	2	0
2	D	24	0	32	2	0
2	E	12	0	16	3	0
3	A	51	0	0	0	0
3	B	48	0	0	1	0
3	C	51	0	0	0	0
3	D	41	0	0	0	0
3	E	27	0	0	0	0
3	F	47	0	0	0	0
All	All	6369	0	6278	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22[B]:LEU:HD21	1:B:37:ALA:HB2	1.74	0.70
1:D:42:ASN:HB3	2:D:203:GOL:H32	1.77	0.66
1:B:86:LYS:HE3	1:B:113[B]:VAL:HG11	1.80	0.64
1:E:71:ASN:HB3	2:E:202:GOL:H31	1.80	0.62
1:C:105:MET:HE1	1:C:115:LYS:HE3	1.83	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/152 (85%)	127 (98%)	2 (2%)	0	100	100
1	B	130/152 (86%)	128 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	130/152 (86%)	129 (99%)	1 (1%)	0	100	100
1	D	128/152 (84%)	127 (99%)	1 (1%)	0	100	100
1	E	128/152 (84%)	126 (98%)	2 (2%)	0	100	100
1	F	130/152 (86%)	129 (99%)	1 (1%)	0	100	100
All	All	775/912 (85%)	766 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/132 (88%)	114 (98%)	2 (2%)	53	45
1	B	117/132 (89%)	112 (96%)	5 (4%)	26	13
1	C	117/132 (89%)	114 (97%)	3 (3%)	40	28
1	D	115/132 (87%)	110 (96%)	5 (4%)	26	13
1	E	116/132 (88%)	113 (97%)	3 (3%)	40	28
1	F	118/132 (89%)	117 (99%)	1 (1%)	73	70
All	All	699/792 (88%)	680 (97%)	19 (3%)	38	27

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	130	CYS
1	E	130	CYS
1	F	52	VAL
1	E	28	GLU
1	C	28	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	HIS
1	B	26	HIS
1	D	42	ASN
1	E	71	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	E	201	-	5,5,5	0.38	0	5,5,5	0.47	0
2	GOL	A	201	-	5,5,5	0.33	0	5,5,5	0.35	0
2	GOL	A	202	-	5,5,5	0.35	0	5,5,5	0.25	0
2	GOL	D	204	-	5,5,5	0.46	0	5,5,5	0.28	0
2	GOL	C	201	-	5,5,5	0.45	0	5,5,5	1.09	0
2	GOL	B	202	-	5,5,5	0.37	0	5,5,5	0.30	0
2	GOL	D	203	-	5,5,5	0.36	0	5,5,5	0.32	0
2	GOL	D	201	-	5,5,5	0.36	0	5,5,5	0.39	0
2	GOL	D	202	-	5,5,5	0.48	0	5,5,5	0.16	0
2	GOL	E	202	-	5,5,5	0.50	0	5,5,5	0.46	0
2	GOL	B	201	-	5,5,5	0.29	0	5,5,5	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	E	201	-	-	0/4/4/4	-
2	GOL	A	201	-	-	0/4/4/4	-
2	GOL	A	202	-	-	0/4/4/4	-
2	GOL	D	204	-	-	2/4/4/4	-
2	GOL	C	201	-	-	2/4/4/4	-
2	GOL	B	202	-	-	2/4/4/4	-
2	GOL	D	203	-	-	2/4/4/4	-
2	GOL	D	201	-	-	2/4/4/4	-
2	GOL	D	202	-	-	2/4/4/4	-
2	GOL	E	202	-	-	4/4/4/4	-
2	GOL	B	201	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	201	GOL	C1-C2-C3-O3
2	D	201	GOL	C1-C2-C3-O3
2	D	202	GOL	O1-C1-C2-C3
2	D	203	GOL	O1-C1-C2-C3
2	D	204	GOL	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	201	GOL	1	0
2	D	204	GOL	1	0
2	C	201	GOL	2	0
2	B	202	GOL	2	0
2	D	203	GOL	1	0
2	E	202	GOL	2	0
2	B	201	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	130/152 (85%)	-0.81	0 100 100	11, 22, 38, 54	1 (0%)
1	B	130/152 (85%)	-0.88	0 100 100	10, 22, 38, 64	2 (1%)
1	C	130/152 (85%)	-0.85	0 100 100	11, 23, 41, 48	2 (1%)
1	D	130/152 (85%)	-0.74	0 100 100	16, 25, 44, 64	0
1	E	130/152 (85%)	-0.83	0 100 100	16, 24, 44, 58	1 (0%)
1	F	130/152 (85%)	-0.85	0 100 100	12, 22, 39, 47	3 (2%)
All	All	780/912 (85%)	-0.83	0 100 100	10, 23, 41, 64	9 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	C	201	6/6	0.96	0.06	33,36,45,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	D	203	6/6	0.96	0.07	36,40,49,57	0
2	GOL	B	201	6/6	0.97	0.05	23,35,39,43	0
2	GOL	D	201	6/6	0.97	0.04	23,34,37,41	0
2	GOL	B	202	6/6	0.97	0.05	35,40,43,53	0
2	GOL	E	201	6/6	0.97	0.09	24,31,33,36	0
2	GOL	E	202	6/6	0.97	0.05	34,37,41,43	0
2	GOL	A	201	6/6	0.98	0.05	29,33,33,37	0
2	GOL	D	204	6/6	0.98	0.06	35,37,44,48	0
2	GOL	A	202	6/6	0.98	0.06	24,34,39,45	0
2	GOL	D	202	6/6	0.98	0.06	33,36,42,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.