



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:41 AM UTC

PDB ID : 6TKV / pdb_00006tkv
Title : Crystal structure of the human FUT8 in complex with GDP and a biantennary complex N-glycan
Authors : Garcia-Garcia, A.; Ceballos-Laita, L.; Serna, L.; Artschwager, R.; Reichardt, N.C.; Corzana, F.; Hurtado-Guerrero, R.
Deposited on : 2019-11-29
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	 4% 85% 13%
2	B	453	 4% 85% 13%
3	C	7	 57% 43%
3	D	7	 71% 29%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8353 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-(1,6)-fucosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3810	2425	670	700	15	0	1	0


- Molecule 2 is a protein called Alpha-(1,6)-fucosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	453	3701	2359	649	679	14	0	2	0

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ASN	deletion	UNP Q9BYC5
B	?	-	SER	deletion	UNP Q9BYC5
B	?	-	ILE	deletion	UNP Q9BYC5
B	?	-	SER	deletion	UNP Q9BYC5
B	?	-	TRP	deletion	UNP Q9BYC5
B	?	-	SER	deletion	UNP Q9BYC5
B	?	-	ALA	deletion	UNP Q9BYC5
B	?	-	GLY	deletion	UNP Q9BYC5
B	?	-	LEU	deletion	UNP Q9BYC5
B	?	-	HIS	deletion	UNP Q9BYC5
B	?	-	ASN	deletion	UNP Q9BYC5
B	?	-	ARG	deletion	UNP Q9BYC5
B	?	-	TYR	deletion	UNP Q9BYC5
B	?	-	THR	deletion	UNP Q9BYC5
B	?	-	GLU	deletion	UNP Q9BYC5

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	7	90	50	4	36	0	0	0
3	D	7	90	50	4	36	0	0	0

- Molecule 4 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	28	10	5	11	2	0	0
4	B	1	28	10	5	11	2	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	243	Total	O	0	0
			243	243		
6	B	261	Total	O	0	0
			261	261		

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

71%

29%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	208.04Å 68.59Å 173.92Å 90.00° 149.90° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 20.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-1.95) 99.1 (20.00-1.95)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.185 , 0.219 0.194 , 0.227	Depositor DCC
R_{free} test set	3503 reflections (3.92%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtrriage
Anisotropy	0.539	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 65.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8353	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, GDP, GOL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.21	9/3913 (0.2%)	1.46	23/5304 (0.4%)
2	B	1.22	12/3799 (0.3%)	1.41	21/5148 (0.4%)
All	All	1.21	21/7712 (0.3%)	1.44	44/10452 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
All	All	0	3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	473	ARG	CD-NE	-7.99	1.35	1.46
1	A	483	HIS	CE1-NE2	7.12	1.39	1.32
1	A	395	ARG	CD-NE	-6.73	1.36	1.46
1	A	347	GLU	C-O	6.71	1.31	1.24
2	B	109	GLY	C-O	6.55	1.32	1.23

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	473	ARG	NE-CZ-NH2	-14.42	106.22	119.20
1	A	180	ARG	NE-CZ-NH2	-13.85	106.74	119.20
2	B	180	ARG	NE-CZ-NH2	-12.08	108.33	119.20
2	B	473	ARG	CD-NE-CZ	11.04	139.86	124.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ARG	NE-CZ-NH2	-10.76	109.52	119.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	180	ARG	Sidechain
2	B	180	ARG	Sidechain
2	B	428	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3810	0	3723	37	1
2	B	3701	0	3628	41	0
3	C	90	0	76	2	0
3	D	90	0	76	1	0
4	A	28	0	12	0	0
4	B	28	0	12	2	0
5	A	24	0	32	6	0
5	B	78	0	104	9	0
6	A	243	0	0	9	1
6	B	261	0	0	7	1
All	All	8353	0	7663	83	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 83 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:244:GLU:OE2	2:B:269:ARG:HD2	1.86	0.75
1:A:110:LYS:HB3	5:A:603:GOL:H2	1.73	0.70
2:B:365:ARG:NH1	2:B:409:ASP:OD2	2.27	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:403:ARG:HD2	1:A:427:ILE:HD11	1.77	0.67
1:A:180:ARG:HD3	6:A:723:HOH:O	1.96	0.66

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:545:ARG:O	1:A:545:ARG:O[2_656]	1.91	0.29
6:A:881:HOH:O	6:B:757:HOH:O[2_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/468 (100%)	456 (98%)	11 (2%)	0	100	100
2	B	451/453 (100%)	437 (97%)	14 (3%)	0	100	100
All	All	918/921 (100%)	893 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/410 (100%)	396 (96%)	15 (4%)	31	21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	400/398 (100%)	387 (97%)	13 (3%)	33	24
All	All	811/808 (100%)	783 (96%)	28 (4%)	33	22

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	572	GLU
2	B	572	GLU
2	B	243	LEU
2	B	352	LYS
2	B	229	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	489	ASN
2	B	148	HIS
2	B	489	ASN
1	A	357	HIS
1	A	391	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3	15,15,15	1.06	0	21,21,21	1.92	5 (23%)
3	NAG	C	2	3	14,14,15	2.37	3 (21%)	17,19,21	2.57	9 (52%)
3	BMA	C	3	3	11,11,12	2.29	3 (27%)	15,15,17	1.15	1 (6%)
3	MAN	C	4	3	11,11,12	1.98	3 (27%)	15,15,17	1.26	2 (13%)
3	NAG	C	5	3	14,14,15	0.86	0	17,19,21	1.36	2 (11%)
3	MAN	C	6	3	11,11,12	2.70	7 (63%)	15,15,17	1.23	2 (13%)
3	NAG	C	7	3	14,14,15	2.60	4 (28%)	17,19,21	2.25	6 (35%)
3	NAG	D	1	3	15,15,15	1.88	3 (20%)	21,21,21	4.44	9 (42%)
3	NAG	D	2	3	14,14,15	1.75	5 (35%)	17,19,21	1.44	3 (17%)
3	BMA	D	3	3	11,11,12	2.48	4 (36%)	15,15,17	1.71	4 (26%)
3	MAN	D	4	3	11,11,12	2.09	3 (27%)	15,15,17	1.29	2 (13%)
3	NAG	D	5	3	14,14,15	1.89	2 (14%)	17,19,21	1.29	2 (11%)
3	MAN	D	6	3	11,11,12	2.15	3 (27%)	15,15,17	1.43	3 (20%)
3	NAG	D	7	3	14,14,15	1.77	3 (21%)	17,19,21	1.75	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3	-	0/6/26/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	NAG	C	5	3	-	0/6/23/26	0/1/1/1
3	MAN	C	6	3	-	0/2/19/22	0/1/1/1
3	NAG	C	7	3	-	2/6/23/26	0/1/1/1
3	NAG	D	1	3	-	1/6/26/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	NAG	D	5	3	-	0/6/23/26	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
3	NAG	D	7	3	-	2/6/23/26	0/1/1/1

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	7	NAG	O5-C1	7.38	1.56	1.43
3	C	2	NAG	C1-C2	-6.99	1.42	1.52
3	C	3	BMA	O5-C5	6.10	1.55	1.43
3	D	5	NAG	C1-C2	5.74	1.60	1.52
3	D	3	BMA	O5-C5	5.19	1.53	1.43

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-C2-C3	-12.39	93.65	110.54
3	D	1	NAG	C1-O5-C5	-10.79	92.78	113.65
3	D	1	NAG	O5-C1-C2	8.09	117.65	109.52
3	C	2	NAG	O7-C7-N2	5.78	132.20	121.98
3	C	7	NAG	C1-O5-C5	5.64	119.75	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
3	D	7	NAG	C4-C5-C6-O6
3	C	7	NAG	O5-C5-C6-O6
3	C	7	NAG	C4-C5-C6-O6


There are no ring outliers.

5 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	NAG	1	0
3	C	6	MAN	1	0
3	D	2	NAG	1	0
3	D	7	NAG	1	0
3	C	7	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	605	-	5,5,5	0.22	0	5,5,5	0.48	0
5	GOL	A	602	-	5,5,5	0.13	0	5,5,5	0.21	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	612	-	5,5,5	0.16	0	5,5,5	0.65	0
5	GOL	B	607	-	5,5,5	0.17	0	5,5,5	0.43	0
5	GOL	B	608	-	5,5,5	0.20	0	5,5,5	0.53	0
5	GOL	B	606	-	5,5,5	0.16	0	5,5,5	0.38	0
5	GOL	B	602	-	5,5,5	0.16	0	5,5,5	0.38	0
5	GOL	B	610	-	5,5,5	0.18	0	5,5,5	0.39	0
5	GOL	B	605	-	5,5,5	0.10	0	5,5,5	0.30	0
4	GDP	B	601	-	29,30,30	2.30	7 (24%)	45,47,47	1.93	10 (22%)
5	GOL	B	609	-	5,5,5	0.11	0	5,5,5	0.27	0
5	GOL	A	603	-	5,5,5	0.18	0	5,5,5	0.52	0
5	GOL	B	611	-	5,5,5	0.17	0	5,5,5	0.39	0
5	GOL	B	614	-	5,5,5	0.19	0	5,5,5	0.40	0
4	GDP	A	601	-	29,30,30	1.90	7 (24%)	45,47,47	1.69	12 (26%)
5	GOL	B	603	-	5,5,5	0.10	0	5,5,5	0.34	0
5	GOL	B	604	-	5,5,5	0.17	0	5,5,5	0.41	0
5	GOL	A	604	-	5,5,5	0.14	0	5,5,5	0.14	0
5	GOL	B	613	-	5,5,5	0.11	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	605	-	-	4/4/4/4	-
5	GOL	A	602	-	-	2/4/4/4	-
5	GOL	B	612	-	-	3/4/4/4	-
5	GOL	B	607	-	-	4/4/4/4	-
5	GOL	B	608	-	-	2/4/4/4	-
5	GOL	B	606	-	-	4/4/4/4	-
5	GOL	B	602	-	-	1/4/4/4	-
5	GOL	B	610	-	-	2/4/4/4	-
5	GOL	B	605	-	-	2/4/4/4	-
4	GDP	B	601	-	-	7/16/32/32	0/3/3/3
5	GOL	B	609	-	-	2/4/4/4	-
5	GOL	A	603	-	-	2/4/4/4	-
5	GOL	B	611	-	-	3/4/4/4	-
5	GOL	B	614	-	-	3/4/4/4	-
4	GDP	A	601	-	-	5/16/32/32	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	603	-	-	2/4/4/4	-
5	GOL	B	604	-	-	0/4/4/4	-
5	GOL	A	604	-	-	0/4/4/4	-
5	GOL	B	613	-	-	4/4/4/4	-

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	601	GDP	PA-O3A	9.43	1.69	1.59
4	A	601	GDP	PA-O3A	7.26	1.67	1.59
4	B	601	GDP	C6-N1	-4.91	1.29	1.38
4	A	601	GDP	O6-C6	2.99	1.29	1.23
4	B	601	GDP	C2-N1	-2.62	1.31	1.37

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	601	GDP	C5-C4-N3	-5.90	118.99	128.39
4	B	601	GDP	C2-N3-C4	5.15	121.17	112.30
4	B	601	GDP	C6-C5-N7	4.16	137.86	130.29
4	A	601	GDP	N9-C4-N3	3.76	133.46	125.95
4	A	601	GDP	O6-C6-C5	-3.57	117.12	126.53

There are no chirality outliers.

5 of 52 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	601	GDP	PB-O3A-PA-O5'
4	B	601	GDP	PA-O3A-PB-O3B
5	A	602	GOL	O1-C1-C2-C3
5	A	603	GOL	O1-C1-C2-O2
5	A	603	GOL	O1-C1-C2-C3

There are no ring outliers.

10 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	605	GOL	1	0
5	A	602	GOL	2	0
5	B	606	GOL	1	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	610	GOL	3	0
4	B	601	GDP	2	0
5	A	603	GOL	2	0
5	B	603	GOL	2	0
5	B	604	GOL	2	0
5	A	604	GOL	1	0
5	B	613	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	429:ASP	C	445:ASN	N	11.79

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	468/468 (100%)	0.07	7 (1%) 72 78	17, 37, 55, 69	1 (0%)
2	B	453/453 (100%)	0.08	17 (3%) 44 50	17, 32, 62, 97	2 (0%)
All	All	921/921 (100%)	0.07	24 (2%) 57 64	17, 35, 59, 97	3 (0%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	370	VAL	6.4
2	B	108	LEU	5.2
2	B	369	LYS	3.7
1	A	434	TRP	3.6
2	B	252	THR	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	7	14/15	0.78	0.14	52,57,61,69	0
3	NAG	C	7	14/15	0.79	0.15	67,73,78,79	0
3	MAN	C	6	11/12	0.87	0.10	47,50,55,62	0
3	NAG	D	1	15/15	0.88	0.10	37,42,45,47	0
3	NAG	C	2	14/15	0.88	0.11	44,48,51,55	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MAN	D	6	11/12	0.89	0.09	38,41,43,45	0
3	NAG	C	1	15/15	0.90	0.09	44,47,49,57	0
3	NAG	C	5	14/15	0.93	0.07	35,36,38,42	0
3	MAN	C	4	11/12	0.93	0.07	39,41,44,45	0
3	BMA	D	3	11/12	0.94	0.07	32,33,34,36	0
3	MAN	D	4	11/12	0.95	0.06	33,35,37,38	0
3	NAG	D	2	14/15	0.95	0.07	34,36,38,41	0
3	BMA	C	3	11/12	0.95	0.07	39,40,41,46	0
3	NAG	D	5	14/15	0.96	0.06	29,30,33,34	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	602	6/6	0.79	0.14	52,59,61,61	0
5	GOL	B	602	6/6	0.80	0.16	67,71,72,74	0
5	GOL	A	603	6/6	0.81	0.15	45,56,58,63	0
5	GOL	A	605	6/6	0.82	0.15	57,66,71,73	0
5	GOL	B	608	6/6	0.84	0.14	63,67,70,72	0
4	GDP	B	601	28/28	0.85	0.14	49,62,67,68	0
5	GOL	B	610	6/6	0.85	0.15	58,66,72,80	0
5	GOL	B	612	6/6	0.85	0.14	57,60,65,69	0
5	GOL	B	614	6/6	0.85	0.11	59,61,61,62	0
5	GOL	B	609	6/6	0.86	0.17	67,69,73,77	0
5	GOL	B	607	6/6	0.87	0.13	53,62,65,65	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	B	604	6/6	0.88	0.14	60,64,65,69	0
5	GOL	B	613	6/6	0.88	0.11	60,62,63,64	0
5	GOL	B	611	6/6	0.88	0.13	58,63,66,66	0
5	GOL	B	605	6/6	0.89	0.11	56,63,66,67	0
5	GOL	B	603	6/6	0.90	0.12	47,51,52,56	0
5	GOL	A	604	6/6	0.92	0.10	54,60,64,67	0
5	GOL	B	606	6/6	0.92	0.10	54,56,58,60	0
4	GDP	A	601	28/28	0.98	0.05	30,32,35,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.