



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:22 PM UTC

PDB ID : 2VB7 / pdb_00002vb7
Title : beta-ketoacyl-ACP synthase I (KAS) from E. coli, apo structure after soak in PEG solution
Authors : Pappenberger, G.; Schulz-Gasch, T.; Bailly, J.; Hennig, M.
Deposited on : 2007-09-06
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	 90% 9%
1	B	406	 91% 8%
1	D	406	 92% 8%
2	C	406	 93% 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	2979	1855	517	581	26	0	17	0
1	B	404	2993	1857	519	592	25	0	19	0
1	D	404	2967	1847	512	583	25	0	14	0

- Molecule 2 is a protein called 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	406	3006	1871	513	594	28	0	19	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	658	Total 658	O 658	0	0
3	B	622	Total 622	O 622	0	0
3	C	648	Total 648	O 648	0	0
3	D	600	Total 600	O 600	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 1



- Molecule 1: 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 1



- Molecule 1: 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 1



- Molecule 2: 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.13Å 139.46Å 212.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	116.25 – 1.60 116.25 – 1.60	Depositor EDS
% Data completeness (in resolution range)	95.8 (116.25-1.60) 95.8 (116.25-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.20 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.156 , 0.186 (Not available) , 0.205	Depositor DCC
R_{free} test set	11193 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.5	Xtrriage
Anisotropy	0.152	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 70.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14473	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/3126	0.80	0/4225
1	B	0.72	0/3155	0.81	1/4265 (0.0%)
1	D	0.72	0/3096	0.82	0/4188
2	C	0.79	1/3147 (0.0%)	0.82	1/4252 (0.0%)
All	All	0.73	1/12524 (0.0%)	0.81	2/16930 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	163	CSD	C-N	18.02	1.58	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	163	CSD	O-C-N	-5.88	113.59	123.00
1	B	133	VAL	N-CA-C	5.50	116.28	110.72

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	163	CSD	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2979	0	2921	35	0
1	B	2993	0	2921	27	0
1	D	2967	0	2896	21	0
2	C	3006	0	2941	18	0
3	A	658	0	0	10	0
3	B	622	0	0	8	0
3	C	648	0	0	3	0
3	D	600	0	0	6	0
All	All	14473	0	11679	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27[B]:ARG:NH1	3:B:2094:HOH:O	1.73	1.21
1:B:53:LYS:HE2	3:B:2182:HOH:O	1.66	0.95
1:A:257[B]:ILE:HD13	1:A:401[B]:MET:HG2	1.50	0.94
1:A:295[A]:LEU:HD12	1:A:386[A]:MET:HE2	1.53	0.90
1:B:257[B]:ILE:HD13	1:B:401[B]:MET:HG2	1.55	0.89
1:A:27[B]:ARG:NH1	3:A:2112:HOH:O	1.99	0.81
1:A:295[B]:LEU:CD1	1:A:386[B]:MET:HG2	2.13	0.78
1:D:163:CYS:SG	3:D:2592:HOH:O	2.40	0.78
1:A:113:GLN:HG2	3:B:2384:HOH:O	1.86	0.76
1:B:28:GLU:OE1	1:B:30:ARG:NH1	2.16	0.73
1:B:113:GLN:NE2	3:B:2304:HOH:O	2.20	0.73
1:D:200:GLU:OE2	3:D:2372:HOH:O	2.07	0.73
1:A:295[B]:LEU:HD13	1:A:386[B]:MET:CG	2.19	0.73
1:D:6[A]:ILE:HD11	1:D:257[A]:ILE:HD11	1.69	0.72
1:A:295[B]:LEU:HD13	1:A:386[B]:MET:HG2	1.71	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:257[B]:ILE:CD1	2:C:401[B]:MET:HG2	2.20	0.72
1:B:204:MET:HE1	1:B:270:VAL:HG11	1.73	0.70
1:D:6[B]:ILE:HD11	1:D:257[B]:ILE:HD11	1.73	0.70
1:A:177[A]:ILE:HD12	1:A:240:VAL:HG12	1.74	0.69
1:A:257[B]:ILE:CD1	1:A:401[B]:MET:HG2	2.22	0.69
2:C:172[B]:ASN:ND2	3:C:2387:HOH:O	2.26	0.68
1:A:110:PRO:HG2	1:A:197:MET:HB2	1.76	0.67
1:B:285[A]:MET:SD	1:B:292:ILE:HD11	2.37	0.64
1:B:257[B]:ILE:CD1	1:B:401[B]:MET:HG2	2.26	0.63
1:A:268:ASP:OD1	1:A:270:VAL:HG22	1.99	0.62
1:A:295[A]:LEU:CD1	1:A:386[A]:MET:HE2	2.26	0.62
1:D:161:SER:OG	1:D:168:HIS:HD2	1.82	0.61
1:A:6:ILE:HD11	1:A:257[A]:ILE:HD11	1.82	0.61
1:A:172[B]:ASN:ND2	3:A:2406:HOH:O	2.34	0.61
2:C:161:SER:OG	2:C:168:HIS:HD2	1.83	0.61
1:A:151[A]:LYS:NZ	3:A:2403:HOH:O	2.01	0.60
1:A:108:GLY:HA3	3:A:2348:HOH:O	2.02	0.60
2:C:200:GLU:OE2	1:D:113:GLN:NE2	2.36	0.59
1:D:53:LYS:HE2	3:D:2177:HOH:O	2.03	0.58
1:B:161:SER:OG	1:B:168:HIS:HD2	1.85	0.58
1:B:30:ARG:HG2	3:B:2088:HOH:O	2.04	0.58
2:C:257[B]:ILE:HD13	2:C:401[B]:MET:HG2	1.85	0.57
1:A:161:SER:OG	1:A:168:HIS:HD2	1.88	0.57
1:B:177[B]:ILE:CD1	1:B:240:VAL:HG12	2.34	0.57
1:A:113:GLN:HE21	1:B:110:PRO:HA	1.70	0.56
1:B:28:GLU:HG3	3:B:2099:HOH:O	2.06	0.56
2:C:177[A]:ILE:HD12	2:C:240:VAL:HG12	1.88	0.56
2:C:110:PRO:HG2	2:C:197:MET:HB2	1.88	0.55
1:D:257[B]:ILE:HD13	1:D:401[B]:MET:HG2	1.87	0.55
1:B:177[B]:ILE:HD12	1:B:240:VAL:HG12	1.88	0.55
2:C:113:GLN:HE21	1:D:110:PRO:HA	1.72	0.55
1:A:172[B]:ASN:HD21	1:B:172[B]:ASN:HD21	1.55	0.55
1:D:30:ARG:HG2	3:D:2076:HOH:O	2.06	0.54
1:D:300:THR:OG1	1:D:302[B]:THR:HG23	2.07	0.54
1:A:302[A]:THR:HG22	3:A:2550:HOH:O	2.08	0.53
1:A:177[A]:ILE:CD1	1:A:240:VAL:HG12	2.38	0.53
1:D:155[B]:VAL:CG2	1:D:157:TYR:CE2	2.93	0.52
1:A:295[A]:LEU:HD12	1:A:386[A]:MET:CE	2.35	0.51
1:D:314[B]:ARG:NH1	1:D:369:ALA:O	2.37	0.51
1:A:380:ARG:NH1	3:A:2637:HOH:O	2.43	0.51
1:B:300:THR:OG1	1:B:302[B]:THR:HG23	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:HB2	3:B:2367:HOH:O	2.10	0.50
1:B:324:ILE:HB	1:B:373[A]:ILE:HD13	1.92	0.50
1:A:200:GLU:OE2	1:B:113:GLN:OE1	2.29	0.50
2:C:30:ARG:NE	3:C:2108:HOH:O	2.45	0.50
2:C:200:GLU:OE2	3:C:2412:HOH:O	2.20	0.49
1:A:273:SER:O	3:A:2514:HOH:O	2.19	0.49
1:A:324:ILE:HB	1:A:373[A]:ILE:HD13	1.94	0.49
1:B:110:PRO:HG2	1:B:197:MET:HB2	1.95	0.48
1:A:101[A]:LEU:HD23	1:A:186:PHE:HB2	1.97	0.47
2:C:163:CSD:HB3	2:C:391:GLY:HA2	1.96	0.47
1:A:283:MET:HE1	3:B:2281:HOH:O	2.15	0.47
2:C:6:ILE:HD11	2:C:257[A]:ILE:HD11	1.97	0.47
1:B:204:MET:HE1	1:B:270:VAL:CG1	2.44	0.46
2:C:113:GLN:HG2	3:D:2372:HOH:O	2.15	0.46
2:C:285[A]:MET:SD	2:C:292:ILE:HD11	2.55	0.46
1:D:1:MET:HA	3:D:2350:HOH:O	2.16	0.46
1:D:257[B]:ILE:CD1	1:D:401[B]:MET:HG2	2.46	0.46
1:A:30:ARG:HG2	3:A:2105:HOH:O	2.17	0.45
2:C:300:THR:OG1	2:C:302[B]:THR:HG23	2.16	0.45
1:D:159:ILE:O	1:D:165:THR:HG23	2.17	0.44
1:A:159:ILE:O	1:A:165:THR:HG23	2.16	0.44
1:A:295[A]:LEU:CD1	1:A:386[A]:MET:CE	2.94	0.44
2:C:124:ARG:HB2	2:C:128:ALA:HB2	1.99	0.43
1:A:110:PRO:HB3	1:B:113:GLN:CD	2.43	0.43
2:C:49:TRP:CE3	2:C:193:LEU:HG	2.54	0.43
1:B:20:GLN:H	1:B:20:GLN:NE2	2.16	0.43
1:A:368:ALA:N	3:A:2616:HOH:O	2.50	0.43
1:B:252:HIS:HE1	1:B:351:GLU:OE2	2.02	0.43
1:D:204:MET:HE1	1:D:270:VAL:HG11	2.01	0.43
1:B:252:HIS:CE1	1:B:351:GLU:OE2	2.71	0.42
1:D:349:MET:HB3	1:D:354:PHE:O	2.18	0.42
2:C:159:ILE:O	2:C:165:THR:HG23	2.19	0.42
1:B:165:THR:HB	1:B:336:GLY:HA2	2.02	0.42
1:D:165:THR:HB	1:D:336:GLY:HA2	2.01	0.41
1:B:6[B]:ILE:CD1	1:B:240:VAL:HG22	2.51	0.41
1:A:151[A]:LYS:HE2	3:A:2401:HOH:O	2.21	0.41
1:D:49:TRP:CE3	1:D:193:LEU:HG	2.55	0.41
1:D:300:THR:O	1:D:301:SER:CB	2.69	0.41
1:A:165:THR:HB	1:A:336:GLY:HA2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/406 (103%)	405 (96%)	13 (3%)	2 (0%)	24	10
1	B	422/406 (104%)	408 (97%)	12 (3%)	2 (0%)	24	10
1	D	417/406 (103%)	403 (97%)	12 (3%)	2 (0%)	24	10
2	C	423/406 (104%)	410 (97%)	11 (3%)	2 (0%)	24	10
All	All	1682/1624 (104%)	1626 (97%)	48 (3%)	8 (0%)	24	10

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	299	GLY
1	B	299	GLY
2	C	299	GLY
1	D	299	GLY
1	B	301	SER
2	C	301	SER
1	D	301	SER
1	A	301	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/309 (102%)	313 (99%)	3 (1%)	70	55
1	B	322/309 (104%)	317 (98%)	5 (2%)	55	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	312/309 (101%)	309 (99%)	3 (1%)	68	50
2	C	318/308 (103%)	315 (99%)	3 (1%)	70	55
All	All	1268/1235 (103%)	1254 (99%)	14 (1%)	68	46

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	CYS
1	A	358	SER
1	A	364	LEU
1	B	13[A]	SER
1	B	13[B]	SER
1	B	20	GLN
1	B	270	VAL
1	B	358	SER
2	C	13[A]	SER
2	C	13[B]	SER
2	C	358	SER
1	D	177	ILE
1	D	358	SER
1	D	362	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	37	GLN
1	A	95	ASN
1	A	168	HIS
1	A	176	GLN
1	A	178	GLN
1	A	396	ASN
1	B	20	GLN
1	B	113	GLN
1	B	168	HIS
1	B	176	GLN
1	B	178	GLN
1	B	252	HIS
1	B	396	ASN
2	C	37	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	168	HIS
2	C	176	GLN
2	C	372	ASN
2	C	396	ASN
1	D	95	ASN
1	D	168	HIS
1	D	176	GLN
1	D	178	GLN
1	D	252	HIS
1	D	296	ASN
1	D	372	ASN
1	D	396	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSD	C	163	2	4,7,8	18.08	1 (25%)	1,8,10	6.19	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	C	163	2	-	0/2/6/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	163	CSD	OD1-SG	36.10	1.80	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	163	CSD	OD1-SG-CB	6.19	117.01	105.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	163	CSD	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/406 (99%)	-0.36	4 (0%) 79 83	7, 14, 25, 39	17 (4%)
1	B	404/406 (99%)	-0.48	4 (0%) 79 83	7, 13, 22, 36	19 (4%)
1	D	404/406 (99%)	-0.38	4 (0%) 79 83	7, 14, 27, 41	14 (3%)
2	C	405/406 (99%)	-0.54	2 (0%) 87 90	7, 12, 22, 36	19 (4%)
All	All	1617/1624 (99%)	-0.44	14 (0%) 81 84	7, 13, 24, 41	69 (4%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	270	VAL	5.8
1	D	319	ASP	4.6
1	D	270	VAL	4.5
2	C	270	VAL	4.5
1	B	270	VAL	4.4
1	A	392	PHE	3.1
1	D	271	ALA	3.0
1	A	273	SER	2.8
1	B	268[A]	ASP	2.7
2	C	392	PHE	2.7
1	B	271	ALA	2.7
1	A	268	ASP	2.5
1	B	392	PHE	2.3
1	D	392	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CSD	C	163	8/9	0.88	0.13	14,16,22,25	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.