



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:25 AM UTC

PDB ID : 5VB8 / pdb\_00005vb8  
Title : Crystal structure of the NavAb voltage-gated sodium channel in an open state  
Authors : Lenaeus, M.J.; Catterall, W.A.  
Deposited on : 2017-03-28  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3554 atoms, of which 1764 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	215	3299	1135	1636	244	272	12	0	0	0


There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	983	MET	-	initiating methionine	UNP A8EVM5
A	984	ASP	-	expression tag	UNP A8EVM5
A	985	TYR	-	expression tag	UNP A8EVM5
A	986	LYS	-	expression tag	UNP A8EVM5
A	987	ASP	-	expression tag	UNP A8EVM5
A	988	ASP	-	expression tag	UNP A8EVM5
A	989	ASP	-	expression tag	UNP A8EVM5
A	990	ASP	-	expression tag	UNP A8EVM5
A	991	LYS	-	expression tag	UNP A8EVM5
A	992	GLY	-	expression tag	UNP A8EVM5
A	993	SER	-	expression tag	UNP A8EVM5
A	994	LEU	-	expression tag	UNP A8EVM5
A	995	VAL	-	expression tag	UNP A8EVM5
A	996	PRO	-	expression tag	UNP A8EVM5
A	997	ARG	-	expression tag	UNP A8EVM5
A	998	GLY	-	expression tag	UNP A8EVM5
A	999	SER	-	expression tag	UNP A8EVM5
A	1000	HIS	-	expression tag	UNP A8EVM5
A	1217	CYS	ILE	conflict	UNP A8EVM5

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is 1,2-DIMYRISTOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PX4) (formula: C<sub>36</sub>H<sub>73</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
3	A	1	52	16	27	8	1	0	0
3	A	1	62	18	34	1	8	1	0
3	A	1	83	26	48	8	1	0	0
3	A	1	40	12	19	8	1	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	12	Total	O	0	0
			12	12		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ion transport protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.30Å 125.30Å 192.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.53 – 2.85 29.53 – 2.85	Depositor EDS
% Data completeness (in resolution range)	91.3 (29.53-2.85) 91.1 (29.53-2.85)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.06 (at 2.85Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.228 , 0.254 0.234 , 0.261	Depositor DCC
$R_{free}$ test set	854 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	88.7	Xtrriage
Anisotropy	0.321	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 89.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3554	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	105.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PX4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	1/1708 (0.1%)	0.69	0/2335

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1127	ILE	CA-CB	5.17	1.56	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1663	1636	1636	42	0
2	A	1	0	0	0	0
3	A	109	128	114	3	0
4	A	5	0	0	0	0
5	A	12	0	0	0	0
All	All	1790	1764	1750	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1177:GLU:OE2	1:A:1178:SER:OG	2.07	0.72
1:A:1083:VAL:HG21	1:A:1105:ARG:HA	1.72	0.71
1:A:1143:ILE:O	1:A:1147:MET:HG3	2.06	0.56
1:A:1025:ASN:OD1	1:A:1105:ARG:HD2	2.07	0.55
1:A:1051:ILE:O	1:A:1054:THR:HB	2.09	0.53
1:A:1162:THR:HG22	1:A:1164:GLY:H	1.76	0.50
1:A:1029:MET:CE	1:A:1103:VAL:HA	2.42	0.49
1:A:1080:ASP:O	1:A:1084:VAL:HG23	2.13	0.49
1:A:1206:THR:O	1:A:1206:THR:HG22	2.13	0.48
1:A:1199:ILE:HB	1:A:1200:PRO:HD3	1.94	0.48
1:A:1082:PHE:CE2	1:A:1086:ILE:HD11	2.47	0.48
1:A:1076:TRP:HE3	1:A:1111:THR:HG22	1.78	0.48
1:A:1052:VAL:HG12	1:A:1056:PHE:CE2	2.49	0.47
1:A:1054:THR:HG22	1:A:1055:ILE:N	2.29	0.47
1:A:1191:TYR:CD1	3:A:1302:PX4:H14	2.50	0.47
1:A:1006:THR:O	1:A:1010:GLU:HG2	2.15	0.46
1:A:1162:THR:HG22	1:A:1163:LEU:N	2.30	0.46
1:A:1201:PHE:O	1:A:1205:VAL:HG22	2.14	0.46
1:A:1208:VAL:O	1:A:1212:LEU:N	2.39	0.46
1:A:1097:ILE:O	1:A:1100:VAL:N	2.46	0.45
1:A:1049:ASN:O	1:A:1053:ILE:HG22	2.17	0.44
1:A:1005:ILE:O	1:A:1008:ILE:N	2.47	0.43
1:A:1207:PHE:HA	1:A:1210:ILE:HD11	1.99	0.42
1:A:1025:ASN:O	1:A:1029:MET:HG2	2.18	0.42
1:A:1030:GLY:O	1:A:1033:THR:HB	2.19	0.42
3:A:1303:PX4:H10	3:A:1303:PX4:O3	2.19	0.42
3:A:1303:PX4:H17	3:A:1303:PX4:H46	2.01	0.42
1:A:1083:VAL:HG21	1:A:1105:ARG:CA	2.47	0.42
1:A:1212:LEU:O	1:A:1212:LEU:HD12	2.19	0.42
1:A:1059:GLU:O	1:A:1063:ARG:HG3	2.20	0.42
1:A:1060:ILE:HG22	1:A:1064:ILE:HD11	2.00	0.42
1:A:1184:VAL:O	1:A:1188:MET:HG3	2.20	0.42
1:A:1119:ILE:O	1:A:1123:LEU:HG	2.20	0.42
1:A:1037:PHE:CD1	1:A:1037:PHE:C	2.98	0.42
1:A:1001:MET:HE3	1:A:1001:MET:HA	2.02	0.42
1:A:1020:TYR:CZ	1:A:1024:LEU:HD21	2.55	0.42
1:A:1207:PHE:CD1	1:A:1207:PHE:C	2.98	0.41
1:A:1020:TYR:CE2	1:A:1024:LEU:HD21	2.55	0.41
1:A:1162:THR:CG2	1:A:1163:LEU:N	2.83	0.41
1:A:1082:PHE:CZ	1:A:1086:ILE:HD11	2.56	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1210:ILE:O	1:A:1214:VAL:HG23	2.21	0.40
1:A:1010:GLU:OE2	1:A:1010:GLU:N	2.54	0.40
1:A:1078:LEU:O	1:A:1079:PHE:C	2.63	0.40
1:A:1207:PHE:HA	1:A:1210:ILE:CD1	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	211/244 (86%)	191 (90%)	20 (10%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	171/223 (77%)	157 (92%)	14 (8%)	10 23

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1010	GLU
1	A	1029	MET

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1033	THR
1	A	1053	ILE
1	A	1054	THR
1	A	1058	ILE
1	A	1070	SER
1	A	1083	VAL
1	A	1123	LEU
1	A	1124	ILE
1	A	1126	VAL
1	A	1139	LEU
1	A	1178	SER
1	A	1187	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PX4	A	1303	-	27,27,45	1.48	3 (11%)	33,35,53	1.86	5 (15%)
3	PX4	A	1302	-	24,24,45	1.47	5 (20%)	27,29,53	1.66	5 (18%)
3	PX4	A	1305	-	20,20,45	1.38	3 (15%)	23,25,53	1.59	3 (13%)
3	PX4	A	1304	-	34,34,45	1.33	3 (8%)	37,39,53	1.59	5 (13%)
4	SO4	A	1306	-	4,4,4	0.25	0	6,6,6	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PX4	A	1303	-	-	14/30/30/49	-
3	PX4	A	1302	-	-	8/26/26/49	-
3	PX4	A	1305	-	-	12/21/21/49	-
3	PX4	A	1304	-	-	8/36/36/49	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1304	PX4	O7-C23	3.90	1.45	1.34
3	A	1304	PX4	O5-C9	3.47	1.43	1.33
3	A	1303	PX4	O7-C23	3.36	1.43	1.34
3	A	1302	PX4	P1-O3	3.23	1.66	1.54
3	A	1302	PX4	O7-C23	2.97	1.42	1.34
3	A	1305	PX4	O7-C23	2.87	1.42	1.34
3	A	1302	PX4	O5-C8	-2.75	1.39	1.45
3	A	1304	PX4	P1-O3	2.72	1.64	1.54
3	A	1303	PX4	P1-O4	2.66	1.69	1.59
3	A	1305	PX4	P1-O3	2.61	1.64	1.54
3	A	1302	PX4	O5-C9	2.58	1.40	1.33
3	A	1303	PX4	O7-C7	-2.34	1.41	1.46
3	A	1302	PX4	O7-C7	-2.13	1.41	1.46
3	A	1305	PX4	P1-O4	2.02	1.66	1.60

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1303	PX4	O7-C23-C24	8.11	129.02	111.48
3	A	1305	PX4	O7-C23-C24	5.82	124.06	111.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1304	PX4	O7-C23-C24	5.80	124.02	111.48
3	A	1302	PX4	O7-C23-C24	4.48	121.18	111.48
3	A	1302	PX4	O5-C9-C10	3.60	122.82	111.83
3	A	1304	PX4	O5-C9-C10	3.27	121.80	111.83
3	A	1303	PX4	O7-C23-O8	-3.02	116.64	123.70
3	A	1302	PX4	O1-P1-O4	2.67	113.63	106.67
3	A	1305	PX4	O7-C23-O8	-2.65	117.50	123.70
3	A	1304	PX4	C8-C7-C6	-2.51	105.93	111.78
3	A	1303	PX4	O8-C23-C24	-2.42	114.31	123.78
3	A	1303	PX4	O1-P1-O4	2.39	118.40	107.57
3	A	1302	PX4	C8-C7-C6	-2.35	106.30	111.78
3	A	1305	PX4	O5-C9-C10	2.24	121.90	112.38
3	A	1304	PX4	O8-C23-C24	-2.22	115.09	123.78
3	A	1304	PX4	O5-C8-C7	2.09	114.41	108.40
3	A	1303	PX4	C8-O5-C9	-2.07	111.98	117.08
3	A	1302	PX4	O7-C7-C8	2.06	115.73	108.34

There are no chirality outliers.

All (42) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1302	PX4	C7-C6-O4-P1
3	A	1303	PX4	C6-O4-P1-O1
3	A	1303	PX4	C6-O4-P1-O3
3	A	1303	PX4	O4-C6-C7-O7
3	A	1303	PX4	O8-C23-O7-C7
3	A	1303	PX4	C24-C23-O7-C7
3	A	1304	PX4	C6-O4-P1-O1
3	A	1304	PX4	C6-O4-P1-O3
3	A	1304	PX4	O8-C23-O7-C7
3	A	1304	PX4	C24-C23-O7-C7
3	A	1303	PX4	O6-C9-O5-C8
3	A	1303	PX4	C10-C9-O5-C8
3	A	1305	PX4	C10-C9-O5-C8
3	A	1304	PX4	O7-C7-C8-O5
3	A	1305	PX4	O6-C9-O5-C8
3	A	1305	PX4	C23-C24-C25-C26
3	A	1302	PX4	O7-C7-C8-O5
3	A	1302	PX4	C24-C25-C26-C27
3	A	1302	PX4	C24-C23-O7-C7
3	A	1305	PX4	C25-C26-C27-C28
3	A	1303	PX4	C24-C25-C26-C27

*Continued on next page...*

*Continued from previous page...*



Mol	Chain	Res	Type	Atoms
3	A	1305	PX4	C24-C23-O7-C7
3	A	1303	PX4	O4-C6-C7-C8
3	A	1302	PX4	O8-C23-O7-C7
3	A	1305	PX4	O8-C23-O7-C7
3	A	1304	PX4	C18-C19-C20-C21
3	A	1304	PX4	C6-O4-P1-O2
3	A	1305	PX4	O4-C6-C7-O7
3	A	1302	PX4	C6-C7-C8-O5
3	A	1304	PX4	C6-C7-C8-O5
3	A	1305	PX4	O7-C23-C24-C25
3	A	1303	PX4	C7-C8-O5-C9
3	A	1305	PX4	O4-C6-C7-C8
3	A	1303	PX4	C1-O3-P1-O1
3	A	1303	PX4	C1-O3-P1-O4
3	A	1303	PX4	C6-O4-P1-O2
3	A	1303	PX4	C26-C27-C28-C29
3	A	1305	PX4	C8-C7-O7-C23
3	A	1305	PX4	C7-C6-O4-P1
3	A	1302	PX4	O7-C23-C24-C25
3	A	1305	PX4	O8-C23-C24-C25
3	A	1302	PX4	O8-C23-C24-C25



There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1303	PX4	2	0
3	A	1302	PX4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/244 (88%)	0.49	27 (12%) <b>8</b> <b>6</b>	57, 110, 150, 170	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1000	HIS	6.0
1	A	1219	ASP	4.7
1	A	1096	GLU	4.0
1	A	1002	TYR	4.0
1	A	1126	VAL	3.2
1	A	1040	SER	3.1
1	A	1089	VAL	3.1
1	A	999	SER	3.0
1	A	1066	VAL	2.9
1	A	1218	VAL	2.9
1	A	1013	PHE	2.9
1	A	1007	ASN	2.9
1	A	1216	ILE	2.8
1	A	1213	VAL	2.7
1	A	1130	MET	2.7
1	A	1132	SER	2.6
1	A	1128	PRO	2.5
1	A	1044	TYR	2.5
1	A	1217	CYS	2.4
1	A	1131	LEU	2.3
1	A	1150	GLN	2.3
1	A	1010	GLU	2.2
1	A	1123	LEU	2.2
1	A	1069	ILE	2.2
1	A	1215	ALA	2.2
1	A	1119	ILE	2.2
1	A	1006	THR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NA	A	1301	1/1	0.51	0.38	76,76,76,76	1
3	PX4	A	1305	21/46	0.78	0.23	98,128,160,169	0
4	SO4	A	1306	5/5	0.85	0.23	135,136,136,136	0
3	PX4	A	1302	25/46	0.89	0.15	78,97,123,132	0
3	PX4	A	1303	28/46	0.91	0.18	76,102,135,142	0
3	PX4	A	1304	35/46	0.91	0.17	84,109,124,131	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around PX4 A 1305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around PX4 A 1302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around PX4 A 1303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.