wwPDB X-ray Structure Validation Summary Report (i)

Aug 19, 2026 – 08:07 AM EDT

PDB ID	:	8VK1 / pdb_00008vk1
Title	:	X-ray crystal structure of human IgE 4C8 Fab complex with Der p 2.0103
Authors	:	Khatri, K.; Ball, A.; Smith, S.A.; Chapman, M.D.; Pomes, A.; Chruszcz, M.
Deposited on	:	2024-01-08
Resolution	:	3.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	212	85% 14%
1	E	212	81% 17%
2	A	129	84% 16%
2	D	129	92% 8%
3	C	232	85% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	232	

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IgE 4C8 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	210	Total	C	N	O	S	0	0	0
			1558	974	257	323	4			
1	B	210	Total	C	N	O	S	0	0	0
			1562	977	258	323	4			

- Molecule 2 is a protein called Der p 2 variant 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	129	Total	C	N	O	S	0	0	0
			980	617	170	185	8			
2	A	129	Total	C	N	O	S	0	0	0
			976	614	169	185	8			

- Molecule 3 is a protein called IgE 4C8 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	226	Total	C	N	O	S	0	0	0
			1641	1029	274	331	7			
3	C	225	Total	C	N	O	S	0	0	0
			1639	1029	274	330	6			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	12	Total	O	0	0
			12	12		
4	D	17	Total	O	0	0
			17	17		
4	F	7	Total	O	0	0
			7	7		
4	B	6	Total	O	0	0
			6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	7	Total O 7 7	0	0
4	A	10	Total O 10 10	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: IgE 4C8 light chain

Chain E:






• Molecule 1: IgE 4C8 light chain

Chain B:






• Molecule 2: Der p 2 variant 3

Chain D:






• Molecule 2: Der p 2 variant 3

Chain A:






• Molecule 3: IgE 4C8 heavy chain


Chain F:





• Molecule 3: IgE 4C8 heavy chain

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	225.25Å 95.30Å 69.21Å 90.00° 105.72° 90.00°	Depositor
Resolution (Å)	39.58 – 3.05 39.58 – 3.05	Depositor EDS
% Data completeness (in resolution range)	98.5 (39.58-3.05) 98.5 (39.58-3.05)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ [1]	2.48 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.197 , 0.248 0.195 , 0.240	Depositor DCC
R_{free} test set	1353 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	60.9	Xtriage
Anisotropy	0.822	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.26 , 49.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.045 for -h-2*l,-k,l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	8415	wwPDB-VP
Average B, all atoms (Å^2)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	B	0.59	0/1601	0.93	0/2193
1	E	0.59	0/1597	0.95	0/2189
2	A	0.55	0/995	0.91	0/1351
2	D	0.59	0/999	0.94	0/1355
3	C	0.56	0/1677	0.94	0/2284
3	F	0.59	0/1679	0.97	0/2288
All	All	0.58	0/8548	0.94	0/11660

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	60	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1562	0	1494	23	0
1	E	1558	0	1483	29	0
2	A	976	0	965	17	0
2	D	980	0	976	5	0
3	C	1639	0	1600	21	0
3	F	1641	0	1594	18	0
4	A	10	0	0	0	0
4	B	6	0	0	0	0
4	C	7	0	0	0	0
4	D	17	0	0	0	0
4	E	12	0	0	0	0
4	F	7	0	0	0	0
All	All	8415	0	8112	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:54:ILE:HD13	2:A:68:ILE:HG21	1.57	0.84
2:A:55:LYS:HG3	2:A:107:THR:HB	1.67	0.76
1:B:132:LEU:HD12	1:B:178:LEU:HD23	1.73	0.70
3:F:173:VAL:HG22	3:F:192:VAL:HG12	1.81	0.62
1:B:60:ARG:HH21	1:B:81:ASP:CG	2.08	0.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	208/212 (98%)	198 (95%)	10 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	208/212 (98%)	199 (96%)	9 (4%)	0	100	100
2	A	127/129 (98%)	120 (94%)	7 (6%)	0	100	100
2	D	127/129 (98%)	123 (97%)	4 (3%)	0	100	100
3	C	223/232 (96%)	216 (97%)	7 (3%)	0	100	100
3	F	224/232 (97%)	220 (98%)	4 (2%)	0	100	100
All	All	1117/1146 (98%)	1076 (96%)	41 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	176/181 (97%)	176 (100%)	0	100	100
1	E	175/181 (97%)	175 (100%)	0	100	100
2	A	107/110 (97%)	105 (98%)	2 (2%)	50	69
2	D	108/110 (98%)	107 (99%)	1 (1%)	70	78
3	C	183/192 (95%)	183 (100%)	0	100	100
3	F	183/192 (95%)	183 (100%)	0	100	100
All	All	932/966 (96%)	929 (100%)	3 (0%)	86	85

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	8	CYS
2	A	8	CYS
2	A	53	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	209	ASN
1	B	41	GLN
2	A	85	GLN
1	B	108	GLN
1	E	65	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	B	210/212 (99%)	-0.35	1 (0%)	87	72	51, 73, 94, 122	0
1	E	210/212 (99%)	-0.29	1 (0%)	87	72	49, 65, 91, 118	0
2	A	129/129 (100%)	-0.36	0	100	100	56, 80, 113, 125	0
2	D	129/129 (100%)	-0.42	0	100	100	48, 68, 88, 100	0
3	C	225/232 (96%)	-0.34	0	100	100	52, 69, 97, 195	0
3	F	226/232 (97%)	-0.43	0	100	100	49, 63, 95, 192	0
All	All	1129/1146 (98%)	-0.36	2 (0%)	91	83	48, 69, 98, 195	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	128	ASN	2.8
1	B	128	ASN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.