



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2026 – 12:07 AM JST

PDB ID : 7VN0 / pdb_00007vn0
Title : CATPO mutant - T188A
Authors : Yuzugullu Karakus, Y.; Balci Unver, S.; Zengin Karatas, M.; Goc, G.; Pearson, A.R.; Yorke, B.
Deposited on : 2021-10-10
Resolution : 1.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	720	 6% 87% 7% 6%
1	B	720	 4% 87% 7% 6%
1	C	720	 % 86% 7% 6%
1	D	720	 % 88% 6% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HDD	A	701	X	-	-	-
2	HDD	B	701	X	-	-	-
2	HDD	C	701	X	-	-	-
2	HDD	D	702	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 24123 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	674	Total	C	N	O	S	0	35	0
			5562	3496	987	1067	12			
1	B	678	Total	C	N	O	S	0	31	0
			5548	3489	982	1065	12			
1	C	676	Total	C	N	O	S	0	26	0
			5486	3458	964	1052	12			
1	D	678	Total	C	N	O	S	0	18	0
			5434	3425	953	1045	11			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP M4GGR7
A	-20	GLY	-	expression tag	UNP M4GGR7
A	-19	SER	-	expression tag	UNP M4GGR7
A	-18	SER	-	expression tag	UNP M4GGR7
A	-17	HIS	-	expression tag	UNP M4GGR7
A	-16	HIS	-	expression tag	UNP M4GGR7
A	-15	HIS	-	expression tag	UNP M4GGR7
A	-14	HIS	-	expression tag	UNP M4GGR7
A	-13	HIS	-	expression tag	UNP M4GGR7
A	-12	HIS	-	expression tag	UNP M4GGR7
A	-11	SER	-	expression tag	UNP M4GGR7
A	-10	SER	-	expression tag	UNP M4GGR7
A	-9	GLY	-	expression tag	UNP M4GGR7
A	-8	GLU	-	expression tag	UNP M4GGR7
A	-7	ASN	-	expression tag	UNP M4GGR7
A	-6	LEU	-	expression tag	UNP M4GGR7
A	-5	TYR	-	expression tag	UNP M4GGR7
A	-4	PHE	-	expression tag	UNP M4GGR7
A	-3	GLN	-	expression tag	UNP M4GGR7
A	-2	GLY	-	expression tag	UNP M4GGR7
A	-1	HIS	-	expression tag	UNP M4GGR7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	188	ALA	THR	engineered mutation	UNP M4GGR7
B	-21	MET	-	initiating methionine	UNP M4GGR7
B	-20	GLY	-	expression tag	UNP M4GGR7
B	-19	SER	-	expression tag	UNP M4GGR7
B	-18	SER	-	expression tag	UNP M4GGR7
B	-17	HIS	-	expression tag	UNP M4GGR7
B	-16	HIS	-	expression tag	UNP M4GGR7
B	-15	HIS	-	expression tag	UNP M4GGR7
B	-14	HIS	-	expression tag	UNP M4GGR7
B	-13	HIS	-	expression tag	UNP M4GGR7
B	-12	HIS	-	expression tag	UNP M4GGR7
B	-11	SER	-	expression tag	UNP M4GGR7
B	-10	SER	-	expression tag	UNP M4GGR7
B	-9	GLY	-	expression tag	UNP M4GGR7
B	-8	GLU	-	expression tag	UNP M4GGR7
B	-7	ASN	-	expression tag	UNP M4GGR7
B	-6	LEU	-	expression tag	UNP M4GGR7
B	-5	TYR	-	expression tag	UNP M4GGR7
B	-4	PHE	-	expression tag	UNP M4GGR7
B	-3	GLN	-	expression tag	UNP M4GGR7
B	-2	GLY	-	expression tag	UNP M4GGR7
B	-1	HIS	-	expression tag	UNP M4GGR7
B	188	ALA	THR	engineered mutation	UNP M4GGR7
C	-21	MET	-	initiating methionine	UNP M4GGR7
C	-20	GLY	-	expression tag	UNP M4GGR7
C	-19	SER	-	expression tag	UNP M4GGR7
C	-18	SER	-	expression tag	UNP M4GGR7
C	-17	HIS	-	expression tag	UNP M4GGR7
C	-16	HIS	-	expression tag	UNP M4GGR7
C	-15	HIS	-	expression tag	UNP M4GGR7
C	-14	HIS	-	expression tag	UNP M4GGR7
C	-13	HIS	-	expression tag	UNP M4GGR7
C	-12	HIS	-	expression tag	UNP M4GGR7
C	-11	SER	-	expression tag	UNP M4GGR7
C	-10	SER	-	expression tag	UNP M4GGR7
C	-9	GLY	-	expression tag	UNP M4GGR7
C	-8	GLU	-	expression tag	UNP M4GGR7
C	-7	ASN	-	expression tag	UNP M4GGR7
C	-6	LEU	-	expression tag	UNP M4GGR7
C	-5	TYR	-	expression tag	UNP M4GGR7
C	-4	PHE	-	expression tag	UNP M4GGR7
C	-3	GLN	-	expression tag	UNP M4GGR7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP M4GGR7
C	-1	HIS	-	expression tag	UNP M4GGR7
C	188	ALA	THR	engineered mutation	UNP M4GGR7
D	-21	MET	-	initiating methionine	UNP M4GGR7
D	-20	GLY	-	expression tag	UNP M4GGR7
D	-19	SER	-	expression tag	UNP M4GGR7
D	-18	SER	-	expression tag	UNP M4GGR7
D	-17	HIS	-	expression tag	UNP M4GGR7
D	-16	HIS	-	expression tag	UNP M4GGR7
D	-15	HIS	-	expression tag	UNP M4GGR7
D	-14	HIS	-	expression tag	UNP M4GGR7
D	-13	HIS	-	expression tag	UNP M4GGR7
D	-12	HIS	-	expression tag	UNP M4GGR7
D	-11	SER	-	expression tag	UNP M4GGR7
D	-10	SER	-	expression tag	UNP M4GGR7
D	-9	GLY	-	expression tag	UNP M4GGR7
D	-8	GLU	-	expression tag	UNP M4GGR7
D	-7	ASN	-	expression tag	UNP M4GGR7
D	-6	LEU	-	expression tag	UNP M4GGR7
D	-5	TYR	-	expression tag	UNP M4GGR7
D	-4	PHE	-	expression tag	UNP M4GGR7
D	-3	GLN	-	expression tag	UNP M4GGR7
D	-2	GLY	-	expression tag	UNP M4GGR7
D	-1	HIS	-	expression tag	UNP M4GGR7
D	188	ALA	THR	engineered mutation	UNP M4GGR7

- Molecule 2 is CIS-HEME D HYDROXYCHLORIN GAMMA-SPIROLACTONE (CCD ID: HDD) (formula: $C_{34}H_{32}FeN_4O_5$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 44	C 34	Fe 1	N 4	O 5	0	0
2	B	1	Total 44	C 34	Fe 1	N 4	O 5	0	0
2	C	1	Total 44	C 34	Fe 1	N 4	O 5	0	0
2	D	1	Total 44	C 34	Fe 1	N 4	O 5	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Ca 3 3	0	0
3	B	2	Total Ca 2 2	0	0
3	C	2	Total Ca 2 2	0	0
3	D	2	Total Ca 2 2	0	0

- Molecule 4 is ALANINE (CCD ID: ALA) (formula: $\text{C}_3\text{H}_7\text{NO}_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			5	3	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	470	Total	O	0	0
			470	470		
5	B	464	Total	O	0	0
			464	464		
5	C	493	Total	O	0	0
			493	493		
5	D	475	Total	O	0	1
			476	476		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Catalase




• Molecule 1: Catalase





• Molecule 1: Catalase





● Molecule 1: Catalase



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	124.94Å 120.79Å 184.75Å 90.00° 101.90° 90.00°	Depositor
Resolution (Å)	56.37 – 1.40 56.37 – 1.40	Depositor EDS
% Data completeness (in resolution range)	98.9 (56.37-1.40) 99.1 (56.37-1.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.161 , 0.182 0.161 , 0.182	Depositor DCC
R_{free} test set	25989 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	10.3	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	24123	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9150e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, HDD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.03	2/5699 (0.0%)	1.16	6/7738 (0.1%)
1	B	1.02	1/5686 (0.0%)	1.15	6/7721 (0.1%)
1	C	1.02	0/5626	1.12	7/7643 (0.1%)
1	D	1.05	3/5572 (0.1%)	1.17	6/7575 (0.1%)
All	All	1.03	6/22583 (0.0%)	1.15	25/30677 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	572	THR	C-O	-6.91	1.16	1.24
1	D	292	VAL	C-O	6.25	1.30	1.24
1	B	173	HIS	CE1-NE2	5.95	1.38	1.32
1	D	241	LYS	C-O	5.85	1.31	1.23
1	A	408	HIS	CE1-NE2	5.12	1.37	1.32

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	697	ASP	CA-C-O	-7.17	108.62	120.80
1	A	619	SER	CA-C-N	6.67	130.45	120.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	619	SER	C-N-CA	6.67	130.45	120.31
1	D	29	ASP	CA-CB-CG	-6.36	106.24	112.60
1	B	157	ILE	CB-CA-C	6.17	116.12	110.13

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	75	ARG	Sidechain
1	D	75	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5562	0	5296	49	0
1	B	5548	0	5292	55	0
1	C	5486	0	5238	53	0
1	D	5434	0	5177	43	0
2	A	44	0	31	0	0
2	B	44	0	31	0	0
2	C	44	0	31	0	0
2	D	44	0	31	0	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	D	5	0	4	0	0
5	A	470	0	0	9	0
5	B	464	0	0	14	0
5	C	493	0	0	8	0
5	D	476	0	0	10	0
All	All	24123	0	21131	188	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 188 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:405[B]:MET:CE	5:B:1029:HOH:O	1.71	1.29
1:A:246[B]:HIS:CE1	1:A:248[B]:LYS:NZ	2.00	1.29
1:B:246[B]:HIS:ND1	1:B:248[B]:LYS:HD3	1.52	1.22
1:B:246[B]:HIS:CE1	1:B:248[B]:LYS:NZ	2.06	1.22
1:C:127[A]:ARG:O	5:C:802:HOH:O	1.58	1.22

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	705/720 (98%)	685 (97%)	20 (3%)	0	100	100
1	B	707/720 (98%)	685 (97%)	22 (3%)	0	100	100
1	C	698/720 (97%)	682 (98%)	16 (2%)	0	100	100
1	D	694/720 (96%)	678 (98%)	16 (2%)	0	100	100
All	All	2804/2880 (97%)	2730 (97%)	74 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	592/596 (99%)	580 (98%)	12 (2%)	48	17
1	B	591/596 (99%)	587 (99%)	4 (1%)	76	52
1	C	585/596 (98%)	578 (99%)	7 (1%)	63	34
1	D	577/596 (97%)	569 (99%)	8 (1%)	59	28
All	All	2345/2384 (98%)	2314 (99%)	31 (1%)	61	30

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	145	THR
1	D	275	ASP
1	C	108	ASN
1	D	640	ARG
1	D	108	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	262	GLN
1	C	461	GLN
1	C	375	ASN
1	D	108	ASN
1	B	46	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 9 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HDD	A	701	1,5	50,52,52	1.83	14 (28%)	56,89,89	1.94	15 (26%)
4	ALA	D	701	1	3,4,5	1.31	0	2,4,6	2.39	1 (50%)
2	HDD	B	701	1	50,52,52	1.97	18 (36%)	56,89,89	2.33	18 (32%)
2	HDD	C	701	1,5	50,52,52	2.04	19 (38%)	56,89,89	1.83	15 (26%)
2	HDD	D	702	1,5	50,52,52	1.87	18 (36%)	56,89,89	1.76	15 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HDD	A	701	1,5	1/1/8/14	2/9/89/89	0/1/9/9
4	ALA	D	701	1	-	0/0/2/4	-
2	HDD	B	701	1	1/1/8/14	2/9/89/89	0/1/9/9
2	HDD	C	701	1,5	1/1/8/14	2/9/89/89	0/1/9/9
2	HDD	D	702	1,5	1/1/8/14	2/9/89/89	0/1/9/9

The worst 5 of 69 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	701	HDD	C1D-ND	4.34	1.38	1.35
2	C	701	HDD	C1B-NB	-4.23	1.33	1.40
2	C	701	HDD	FE-NB	4.18	2.07	1.94
2	A	701	HDD	CHA-C1A	4.12	1.48	1.39
2	D	702	HDD	FE-NB	4.05	2.07	1.94

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	HDD	C3B-C2B-C1B	-9.78	99.23	106.49
2	A	701	HDD	C3B-C2B-C1B	-5.04	102.75	106.49
2	C	701	HDD	C3C-C2C-C1C	-4.99	102.31	107.08
2	D	702	HDD	C4B-C3B-C2B	-4.53	103.52	107.11
2	A	701	HDD	CAA-C2A-C1A	4.44	133.64	124.89

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	701	HDD	NC
2	B	701	HDD	NC
2	C	701	HDD	NC
2	D	702	HDD	NC


5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	701	HDD	CAA-CBA-CGA-O1A
2	C	701	HDD	CAA-CBA-CGA-O1A
2	C	701	HDD	CAA-CBA-CGA-O2A
2	D	702	HDD	CAA-CBA-CGA-O1A
2	A	701	HDD	CAA-CBA-CGA-O1A


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	674/720 (93%)	-0.05	46 (6%)	23 24	4, 12, 32, 59	35 (5%)
1	B	678/720 (94%)	-0.12	28 (4%)	41 43	4, 13, 34, 72	31 (4%)
1	C	676/720 (93%)	-0.38	6 (0%)	81 84	4, 11, 22, 62	26 (3%)
1	D	678/720 (94%)	-0.18	9 (1%)	75 79	4, 13, 26, 46	18 (2%)
All	All	2706/2880 (93%)	-0.18	89 (3%)	49 51	4, 12, 28, 72	110 (4%)

The worst 5 of 89 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	621	ALA	6.1
1	C	618	ALA	5.5
1	B	655	VAL	5.1
1	D	621	ALA	4.8
1	C	619	SER	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ALA	D	701	5/6	0.74	0.16	22,24,25,26	0
3	CA	B	703	1/1	0.97	0.05	18,18,18,18	1
3	CA	D	704	1/1	0.98	0.04	15,15,15,15	1
2	HDD	D	702	44/44	0.99	0.04	7,8,10,12	0
3	CA	A	703	1/1	0.99	0.04	14,14,14,14	1
3	CA	A	704	1/1	0.99	0.08	11,11,11,11	1
3	CA	B	702	1/1	0.99	0.05	14,14,14,14	0
2	HDD	A	701	44/44	0.99	0.04	7,8,10,13	0
3	CA	C	703	1/1	0.99	0.03	14,14,14,14	1
2	HDD	B	701	44/44	0.99	0.04	7,8,10,12	0
2	HDD	C	701	44/44	0.99	0.04	6,7,10,11	0
3	CA	D	703	1/1	1.00	0.05	15,15,15,15	0
3	CA	C	702	1/1	1.00	0.02	9,9,9,9	1
3	CA	A	702	1/1	1.00	0.05	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HDD D 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HDD A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HDD B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HDD C 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.