



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:52 AM UTC

PDB ID : 2VXT / pdb_00002vxt
Title : Crystal structure of human IL-18 complexed to murine reference antibody 125-2H Fab
Authors : Argiriadi, M.A.; Xiang, T.; Wu, C.; Ghayur, T.; Borhani, D.W.
Deposited on : 2008-07-10
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	216	 12% 88% 8%
2	I	157	 15% 85% 13%
3	L	214	 7% 92% 8%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5375 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MURINE IGG 125-2H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	207	1692	1068	278	337	9	0	16	0

- Molecule 2 is a protein called INTERLEUKIN-18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	I	156	1328	837	223	260	8	0	9	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	74	ALA	CYS	engineered mutation	UNP Q14116
I	104	ALA	CYS	engineered mutation	UNP Q14116
I	112	ALA	CYS	engineered mutation	UNP Q14116
I	163	ALA	CYS	engineered mutation	UNP Q14116

- Molecule 3 is a protein called MURINE IGG 125-2H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1759	1087	292	371	9	0	14	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total Cl 1 1	0	0
4	I	1	Total Cl 1 1	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	I	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	212	Total O 212 212	0	0
6	I	128	Total O 128 128	0	0
6	L	253	Total O 253 253	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MURINE IGG 125-2H



- Molecule 2: INTERLEUKIN-18



- Molecule 3: MURINE IGG 125-2H



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.34Å 62.44Å 104.73Å 90.00° 100.39° 90.00°	Depositor
Resolution (Å)	31.16 – 1.49 31.16 – 1.50	Depositor EDS
% Data completeness (in resolution range)	95.5 (31.16-1.49) 95.4 (31.16-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.162 , 0.196 0.172 , 0.204	Depositor DCC
R_{free} test set	4364 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5375	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	H	0.72	1/1736 (0.1%)	0.87	0/2369
2	I	0.61	0/1348	0.84	0/1807
3	L	0.69	0/1796	0.90	0/2435
All	All	0.68	1/4880 (0.0%)	0.88	0/6611

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1
2	I	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	96	LEU	C-N	7.88	1.43	1.33

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	40	SER	Mainchain
2	I	92	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1692	0	1629	12	0
2	I	1328	0	1318	33	0
3	L	1759	0	1655	14	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
5	I	1	0	0	0	0
6	H	212	0	0	0	0
6	I	128	0	0	5	0
6	L	253	0	0	5	0
All	All	5375	0	4602	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:69:MET:HE1	2:I:74:ALA:HA	1.37	1.05
2:I:69:MET:CE	2:I:74:ALA:HA	1.90	1.00
2:I:92:GLN:HE22	2:I:96[B]:MET:CE	1.82	0.92
2:I:69:MET:CE	2:I:74:ALA:CA	2.56	0.83
2:I:92:GLN:HE22	2:I:96[B]:MET:HE2	1.42	0.82
2:I:136[B]:ILE:CD1	6:I:2064:HOH:O	2.36	0.72
2:I:69:MET:HE3	2:I:74:ALA:N	2.05	0.71
2:I:94:ARG:O	2:I:140[B]:ARG:NE	2.26	0.68
2:I:42:GLU:OE1	2:I:87[A]:MET:CE	2.43	0.67
2:I:42:GLU:OE1	2:I:87[A]:MET:HE1	1.95	0.65
2:I:92:GLN:NE2	2:I:96[B]:MET:HE2	2.12	0.65
3:L:146:VAL:CG1	3:L:175[A]:MET:HE1	2.27	0.65
2:I:42:GLU:HB2	2:I:87[A]:MET:HE3	1.81	0.63
2:I:69:MET:HE3	2:I:74:ALA:CA	2.29	0.61
3:L:146:VAL:HG11	3:L:175[A]:MET:HE1	1.83	0.60
2:I:136[B]:ILE:HD12	6:I:2064:HOH:O	2.01	0.58
2:I:87[A]:MET:HG2	2:I:98:VAL:HG12	1.84	0.58
2:I:45:LEU:HD23	2:I:84:ILE:HG13	1.87	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:69:MET:HE3	2:I:74:ALA:HA	1.81	0.56
3:L:78:LEU:HD21	3:L:104[A]:LEU:HD21	1.89	0.55
1:H:148:GLU:HG3	1:H:175:TYR:CZ	2.42	0.54
2:I:62:ASN:O	2:I:119:PHE:HD2	1.91	0.53
2:I:87[A]:MET:HE2	6:I:2030:HOH:O	2.07	0.53
3:L:117:ILE:HG13	3:L:134[A]:CYS:SG	2.47	0.53
2:I:93:PRO:HB2	6:L:2054:HOH:O	2.09	0.53
2:I:92:GLN:HE22	2:I:96[B]:MET:HE1	1.72	0.52
1:H:143:LYS:HE3	1:H:171:GLN:HE22	1.76	0.51
3:L:146:VAL:HG11	3:L:175[A]:MET:CE	2.40	0.51
1:H:184:PRO:HB2	1:H:187[A]:THR:HG23	1.94	0.50
3:L:187:GLU:OE1	6:L:2224:HOH:O	2.19	0.50
2:I:69:MET:HE1	2:I:74:ALA:CA	2.20	0.50
2:I:69:MET:HE3	2:I:73:ASP:C	2.37	0.49
1:H:41[B]:HIS:HB2	1:H:43[B]:LYS:HG3	1.95	0.48
1:H:148:GLU:HG3	1:H:175:TYR:CE2	2.49	0.47
2:I:136[B]:ILE:HD11	6:I:2064:HOH:O	2.05	0.47
3:L:143[A]:ASP:OD2	6:L:2171:HOH:O	2.21	0.47
1:H:43[B]:LYS:HB2	1:H:44[B]:SER:H	1.64	0.46
3:L:10[A]:SER:HB2	6:L:2023:HOH:O	2.17	0.45
2:I:111:SER:O	2:I:117[B]:ILE:HD12	2.18	0.44
1:H:20:VAL:HG21	1:H:80:MET:HE2	2.00	0.44
1:H:47:TRP:CZ2	1:H:49:GLY:HA2	2.53	0.44
1:H:20:VAL:CG2	1:H:80:MET:HE2	2.48	0.43
2:I:58:ILE:CG1	2:I:62:ASN:HA	2.48	0.43
1:H:184:PRO:HD2	1:H:187[A]:THR:HG21	2.00	0.43
3:L:119:PRO:HB3	3:L:209:PHE:CE1	2.53	0.43
1:H:50:ASP:OD1	1:H:58[A]:SER:HB3	2.18	0.43
2:I:42:GLU:OE1	2:I:87[A]:MET:HE3	2.19	0.43
3:L:83:PHE:HA	3:L:104[A]:LEU:HB3	2.00	0.42
3:L:3:GLN:NE2	6:L:2006:HOH:O	2.52	0.42
3:L:11:LEU:HB3	3:L:104[B]:LEU:CD2	2.49	0.42
3:L:11:LEU:HB3	3:L:104[B]:LEU:HD22	2.03	0.41
2:I:69:MET:CE	2:I:74:ALA:CB	2.98	0.41
1:H:80:MET:HE1	1:H:90:TYR:CG	2.56	0.41
2:I:96[B]:MET:HE2	6:I:2037:HOH:O	2.21	0.41
2:I:50:ASN:OD1	2:I:50:ASN:C	2.64	0.40
2:I:131:THR:HA	2:I:136[B]:ILE:HD13	2.04	0.40
2:I:69:MET:HE2	2:I:74:ALA:HB2	2.03	0.40
2:I:180:LEU:HD21	3:L:91:TYR:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	220/216 (102%)	212 (96%)	8 (4%)	0	100	100
2	I	163/157 (104%)	159 (98%)	4 (2%)	0	100	100
3	L	226/214 (106%)	223 (99%)	3 (1%)	0	100	100
All	All	609/587 (104%)	594 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	197/187 (105%)	197 (100%)	0	100	100
2	I	151/144 (105%)	150 (99%)	1 (1%)	76	57
3	L	205/191 (107%)	203 (99%)	2 (1%)	68	45
All	All	553/522 (106%)	550 (100%)	3 (0%)	78	66

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	92	GLN
3	L	89	LEU
3	L	213	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such

sidechains are listed below:

Mol	Chain	Res	Type
1	H	171	GLN
2	I	52	ASN
2	I	92	GLN
2	I	123	ASN
3	L	156	GLN
3	L	189	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	207/216 (95%)	0.81	26 (12%) 8 8	7, 18, 32, 41	16 (7%)
2	I	156/157 (99%)	1.18	24 (15%) 5 5	9, 21, 35, 45	9 (5%)
3	L	213/214 (99%)	0.84	16 (7%) 20 22	7, 20, 29, 39	14 (6%)
All	All	576/587 (98%)	0.92	66 (11%) 9 10	7, 20, 32, 45	39 (6%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	170	LEU	4.8
1	H	171	GLN	4.1
3	L	134[A]	CYS	3.9
1	H	172	SER	3.9
2	I	74	ALA	3.6
3	L	68	GLY	3.6
2	I	170	PHE	3.5
1	H	189	PRO	3.4
1	H	169	VAL	3.3
1	H	211	VAL	3.3
2	I	93	PRO	3.3
1	H	163	VAL	3.2
2	I	169	LEU	3.1
2	I	105	GLU	3.1
2	I	63	ARG	3.0
1	H	127	GLY	3.0
2	I	76	ASP	3.0
3	L	163[A]	TRP	3.0
3	L	67	SER	3.0
2	I	136[A]	ILE	2.9
1	H	173	ASP	2.9
1	H	174	LEU	2.9
1	H	188	TRP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	187[A]	THR	2.9
2	I	71	ASP	2.8
3	L	213	GLU	2.7
1	H	175	TYR	2.7
2	I	92	GLN	2.7
2	I	168	ASP	2.7
3	L	184	ASP	2.7
3	L	70	ASP	2.6
1	H	161	SER	2.6
1	H	190	SER	2.6
2	I	72	SER	2.6
2	I	191	ASN	2.6
3	L	69	SER	2.6
2	I	61	GLY	2.6
2	I	158	GLY	2.6
2	I	104	ALA	2.5
3	L	44	PHE	2.5
2	I	90	ASP	2.5
1	H	162	GLY	2.5
1	H	159	LEU	2.5
3	L	98	PHE	2.4
1	H	41[A]	HIS	2.4
1	H	191	GLU	2.4
1	H	182	THR	2.3
2	I	91	SER	2.3
2	I	116	ILE	2.3
1	H	168	ALA	2.3
3	L	127[A]	SER	2.3
2	I	78	ALA	2.2
1	H	136[A]	VAL	2.2
2	I	156	TYR	2.2
3	L	210	ASN	2.2
2	I	38	PHE	2.2
2	I	192	GLU	2.2
3	L	212	ASN	2.1
2	I	159	TYR	2.1
1	H	154	TRP	2.1
3	L	165	ASP	2.1
3	L	211	ARG	2.1
1	H	157	GLY	2.1
1	H	167	PRO	2.1
3	L	95	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	166	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	H	1214	1/1	0.97	0.12	27,27,27,27	0
4	CL	I	1193	1/1	0.97	0.17	28,28,28,28	0
5	MG	I	1194	1/1	0.98	0.11	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.