Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 11:32 AM EDT

PDB ID	:	2WBG / pdb_00002wbg
Title	:	Structure of family 1 beta-glucosidase from Thermotoga maritima in complex with 3-imino-2-oxa-(+)-castanospermine
Authors	:	Aguilar, M.; Gloster, T.M.; Turkenburg, J.P.; Garcia-Moreno, M.I.; Ortiz Mellet, C.; Davies, G.J.; Garcia Fernandez, J.M.
Deposited on	:	2009-02-27
Resolution	:	1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	87% 7% • 5%
1	B	468	86% 7% • 5%
1	C	468	90% 5% 5%
1	D	468	86% 8% • 5%

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 16641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-GLUCOSIDASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	18	0
			3709	2421	619	663	6			
1	B	443	Total	C	N	O	S	0	9	0
			3647	2382	603	655	7			
1	C	443	Total	C	N	O	S	0	22	0
			3755	2449	625	675	6			
1	D	443	Total	C	N	O	S	0	15	0
			3671	2393	606	666	6			

- Molecule 2 is (3Z,5S,6R,7S,8R,8aR)-3-(octylimino)hexahydro[1,3]oxazolo[3,4-a]pyridine-5,6,7,8-tetrol (CCD ID: LGS) (formula: $C_{15}H_{28}N_2O_5$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			22	15	2	5		
2	B	1	Total	C	N	O	0	0
			22	15	2	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C N O 22 15 2 5	0	0
2	D	1	Total C N O 22 15 2 5	0	0

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total C O 4 2 2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	411	Total O 411 411	0	0
4	B	379	Total O 379 379	0	0
4	C	522	Total O 522 522	0	0
4	D	455	Total O 455 455	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: BETA-GLUCOSIDASE A

Chain A:




• Molecule 1: BETA-GLUCOSIDASE A

Chain B:




• Molecule 1: BETA-GLUCOSIDASE A

Chain C:



• Molecule 1: BETA-GLUCOSIDASE A

Chain D:




N246
Y276
V300
K301
F302
D303
P304
D305
A306
P307
A308
K309
V310
M322
K336
E340
A355
S361
N370
W383
Q387
N403
E408
R413
N442
E445
ASP

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	89.99Å 73.19Å 138.09Å 90.00° 94.03° 90.00°	Depositor
Resolution (Å)	137.36 – 1.85 137.36 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (137.36-1.85) 99.8 (137.36-1.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.18 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.4.0077	Depositor
R, R_{free}	0.153 , 0.199 0.153 , 0.199	Depositor DCC
R_{free} test set	7667 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	16.3	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 55.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	16641	wwPDB-VP
Average B, all atoms (Å^2)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4638e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, LGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.87	0/3877	0.88	6/5269 (0.1%)
1	B	0.79	1/3788 (0.0%)	0.85	2/5150 (0.0%)
1	C	0.94	0/3931	0.89	0/5339
1	D	0.86	0/3832	0.89	0/5212
All	All	0.87	1/15428 (0.0%)	0.88	8/20970 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	344	PRO	CA-C	7.56	1.56	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	327	VAL	CA-C-N	6.45	126.21	119.05
1	B	327	VAL	C-N-CA	6.45	126.21	119.05
1	A	82	TRP	CA-C-N	-5.67	113.92	120.04
1	A	82	TRP	C-N-CA	-5.67	113.92	120.04
1	A	343[A]	ASN	CA-C-N	-5.15	115.95	119.66
1	A	343[A]	ASN	C-N-CA	-5.15	115.95	119.66
1	A	343[B]	ASN	CA-C-N	-5.15	115.95	119.66
1	A	343[B]	ASN	C-N-CA	-5.15	115.95	119.66

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3709	0	3609	24	0
1	B	3647	0	3505	28	0
1	C	3755	0	3666	17	0
1	D	3671	0	3512	28	0
2	A	22	0	28	0	0
2	B	22	0	28	0	0
2	C	22	0	28	0	0
2	D	22	0	28	0	0
3	C	4	0	3	1	0
4	A	411	0	0	4	0
4	B	379	0	0	5	0
4	C	522	0	0	1	0
4	D	455	0	0	4	0
All	All	16641	0	14407	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:192[B]:ARG:HH11	1:C:192[B]:ARG:CG	1.54	1.20
1:C:192[B]:ARG:HG2	1:C:192[B]:ARG:NH1	1.50	1.08
1:B:185:ARG:HG3	1:B:185:ARG:HH11	1.20	1.06
1:B:216:LYS:HD3	4:B:2311:HOH:O	1.69	0.91
1:D:370:ASN:HB3	4:D:2401:HOH:O	1.72	0.89
1:C:192[B]:ARG:HH11	1:C:192[B]:ARG:HG2	0.74	0.89
1:B:185:ARG:HH11	1:B:185:ARG:CG	1.90	0.85
1:A:333:TRP:CZ2	1:A:337:LYS:HD2	2.17	0.79
1:A:144:ALA:HB2	1:A:203[A]:ARG:HG2	1.65	0.79
1:A:144:ALA:HB2	1:A:203[A]:ARG:CG	2.17	0.75
1:B:333:TRP:CZ2	1:B:337:LYS:HD3	2.23	0.74
1:B:246:ASN:HB3	4:B:2239:HOH:O	1.91	0.70
1:D:304:PRO:O	1:D:305:ASP:CG	2.36	0.69
1:D:304:PRO:O	1:D:305:ASP:CB	2.41	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:ALA:HB3	1:B:20:GLN:HE21	1.60	0.67
1:A:141:ASP:OD1	1:A:203[A]:ARG:NH1	2.29	0.65
1:A:333:TRP:CZ2	1:A:337:LYS:CD	2.80	0.64
1:D:304:PRO:O	1:D:305:ASP:OD1	2.15	0.63
1:B:144:ALA:HB2	1:B:203[A]:ARG:HG2	1.81	0.63
1:C:148[B]:ARG:HG3	1:C:149:VAL:N	2.14	0.63
1:D:370:ASN:CB	4:D:2401:HOH:O	2.39	0.61
1:B:185:ARG:CG	1:B:185:ARG:NH1	2.58	0.58
1:D:7:PRO:HG2	4:D:2417:HOH:O	2.02	0.58
1:C:246:ASN:OD1	3:C:1447:ACT:H1	2.04	0.58
1:D:302:PHE:CE2	1:D:304:PRO:HD3	2.40	0.56
1:D:213:LYS:O	1:D:214:ASP:HB2	2.06	0.55
1:D:5:LYS:HE2	1:D:442:ASN:HB3	1.89	0.55
1:D:383:TRP:O	1:D:387:GLN:HG2	2.07	0.55
1:A:88[B]:GLU:OE2	1:A:92:ARG:NH2	2.38	0.55
1:C:109[B]:LEU:HD11	1:C:157:ARG:HB3	1.89	0.54
1:A:144:ALA:HB2	1:A:203[A]:ARG:HG3	1.89	0.52
1:A:184:MET:HE2	1:C:184:MET:HE2	1.91	0.52
1:C:274[A]:GLU:O	1:C:275[A]:ASN:HB2	2.10	0.51
1:A:262[A]:GLU:HG3	1:A:263:LEU:N	2.25	0.51
1:B:185:ARG:HG3	1:B:185:ARG:NH1	2.02	0.51
1:B:152[B]:GLU:OE1	1:B:152[B]:GLU:CA	2.57	0.51
1:A:262[A]:GLU:HG3	1:A:263:LEU:HG	1.93	0.51
1:A:185:ARG:NH1	4:A:2205:HOH:O	2.43	0.51
1:D:236:ILE:O	1:D:240[A]:ARG:HG3	2.12	0.50
1:A:307:PRO:C	1:A:309:LYS:H	2.20	0.49
1:A:333:TRP:CE2	1:A:337:LYS:HD2	2.46	0.49
1:D:304:PRO:O	1:D:305:ASP:HB2	2.11	0.49
1:B:300:VAL:HG13	1:B:310[B]:VAL:HG23	1.93	0.49
1:C:300[A]:VAL:HG13	1:C:310:VAL:HG13	1.94	0.49
1:B:322:MET:HE1	1:B:408:GLU:HG3	1.95	0.48
1:B:141:ASP:OD1	1:B:203[B]:ARG:NH2	2.46	0.48
1:A:74:LYS:HD3	4:A:2102:HOH:O	2.14	0.48
1:B:265:LEU:O	1:B:269:ARG:HB2	2.13	0.48
1:A:90:THR:HG22	4:A:2118:HOH:O	2.14	0.47
1:A:20:GLN:O	1:A:403:ASN:HB2	2.13	0.47
1:C:192[B]:ARG:CG	1:C:192[B]:ARG:NH1	2.28	0.46
1:B:144:ALA:HB2	1:B:203[A]:ARG:CG	2.45	0.46
1:B:141:ASP:OD1	1:B:203[A]:ARG:NH1	2.37	0.46
1:B:238:ALA:HB1	1:B:310[A]:VAL:HG12	1.96	0.46
1:B:275:ASN:HB2	4:B:2264:HOH:O	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:223:ASN:HA	1:B:247:ASN:OD1	2.16	0.46
1:B:4:LYS:HE3	4:B:2004:HOH:O	2.16	0.45
1:B:26:LEU:HD21	1:B:32:MET:HG2	1.99	0.45
1:B:121:HIS:O	1:B:122:TRP:HB2	2.16	0.45
1:D:216:LYS:HD2	4:D:2372:HOH:O	2.16	0.45
1:A:5:LYS:HE2	1:A:441:ASN:O	2.16	0.45
1:C:109[A]:LEU:HA	1:C:109[A]:LEU:HD13	1.68	0.44
1:A:262[A]:GLU:CG	1:A:263:LEU:N	2.80	0.44
1:A:247:ASN:C	1:A:249:PRO:CD	2.91	0.43
1:B:303:ASP:O	1:B:309:LYS:HA	2.18	0.43
1:C:322:MET:HE1	1:C:408:GLU:HG3	2.00	0.43
1:C:336:LYS:HB3	1:C:336:LYS:HE2	1.87	0.43
1:D:355:ALA:HA	1:D:413:ARG:O	2.18	0.43
1:B:336:LYS:HB3	1:B:336:LYS:HE2	1.91	0.43
1:A:316:ASP:OD1	4:A:2312:HOH:O	2.22	0.43
1:D:336:LYS:O	1:D:340[A]:GLU:HG3	2.20	0.42
1:C:46:LYS:HD3	4:C:2479:HOH:O	2.19	0.41
1:D:82:TRP:HB3	1:D:83:PRO:HD3	2.02	0.41
1:B:398:TRP:HA	1:B:399:SER:HA	1.85	0.41
1:D:322:MET:HE1	1:D:408:GLU:HG3	2.01	0.41
1:A:231:GLU:O	1:A:232:LYS:C	2.63	0.41
1:C:274[A]:GLU:O	1:C:275[A]:ASN:CB	2.68	0.41
1:D:276:TYR:CD1	1:D:276:TYR:C	2.99	0.41
1:A:68:ILE:HG23	1:A:73:VAL:HB	2.01	0.41
1:A:300:VAL:HG13	1:A:310[A]:VAL:CG1	2.51	0.41
1:B:403:ASN:ND2	4:B:2343:HOH:O	2.50	0.41
1:C:24:SER:N	1:C:25:PRO:CD	2.84	0.41
1:C:405:GLU:O	1:C:405:GLU:HG3	2.21	0.41
1:D:207:VAL:O	1:D:211:THR:HG23	2.20	0.41
1:D:403:ASN:HD22	1:D:403:ASN:C	2.29	0.41
1:B:248:TYR:N	1:B:249:PRO:CD	2.84	0.40
1:D:28:ASP:HB3	1:D:96:LYS:HD3	2.03	0.40
1:D:26:LEU:HD21	1:D:32:MET:HG2	2.03	0.40
1:D:300[A]:VAL:HG13	1:D:310:VAL:HG13	2.03	0.40
1:D:307:PRO:O	1:D:308:ALA:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	459/468 (98%)	451 (98%)	8 (2%)	0	100	100
1	B	450/468 (96%)	443 (98%)	7 (2%)	0	100	100
1	C	464/468 (99%)	461 (99%)	2 (0%)	1 (0%)	43	31
1	D	457/468 (98%)	453 (99%)	2 (0%)	2 (0%)	30	17
All	All	1830/1872 (98%)	1808 (99%)	19 (1%)	3 (0%)	43	31

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	305	ASP
1	D	122	TRP
1	C	122	TRP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/399 (98%)	376 (97%)	13 (3%)	33	18
1	B	376/399 (94%)	369 (98%)	7 (2%)	50	38
1	C	397/399 (100%)	387 (98%)	10 (2%)	42	27
1	D	379/399 (95%)	372 (98%)	7 (2%)	51	40
All	All	1541/1596 (97%)	1504 (98%)	37 (2%)	48	28

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	VAL
1	A	5	LYS
1	A	74	LYS
1	A	246[A]	ASN
1	A	246[B]	ASN
1	A	317[A]	LEU
1	A	317[B]	LEU
1	A	337	LYS
1	A	350	THR
1	A	361	SER
1	A	365[A]	ARG
1	A	365[B]	ARG
1	A	403	ASN
1	B	80	ILE
1	B	109	LEU
1	B	185	ARG
1	B	216	LYS
1	B	269	ARG
1	B	317	LEU
1	B	403	ASN
1	C	3	VAL
1	C	109[A]	LEU
1	C	109[B]	LEU
1	C	185	ARG
1	C	300[A]	VAL
1	C	300[B]	VAL
1	C	316	ASP
1	C	317[A]	LEU
1	C	317[B]	LEU
1	C	403	ASN
1	D	3	VAL
1	D	203	ARG
1	D	300[A]	VAL
1	D	300[B]	VAL
1	D	361	SER
1	D	370	ASN
1	D	403	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	GLN
1	A	275	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	381	GLN
1	A	387	GLN
1	A	403	ASN
1	A	437	ASN
1	B	20	GLN
1	B	95	GLN
1	B	129	GLN
1	B	343	ASN
1	B	381	GLN
1	B	387	GLN
1	B	403	ASN
1	C	129	GLN
1	C	381	GLN
1	C	387	GLN
1	C	403	ASN
1	D	47	ASN
1	D	129	GLN
1	D	387	GLN
1	D	403	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	LGS	A	1446	-	22,23,23	1.85	3 (13%)	22,31,31	1.95	5 (22%)
2	LGS	D	1446	-	22,23,23	1.93	2 (9%)	22,31,31	2.14	6 (27%)
2	LGS	B	1446	-	22,23,23	1.64	2 (9%)	22,31,31	2.40	7 (31%)
3	ACT	C	1447	-	3,3,3	0.70	0	3,3,3	1.44	0
2	LGS	C	1446	-	22,23,23	1.48	3 (13%)	22,31,31	2.94	11 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LGS	B	1446	-	-	1/7/42/42	0/2/2/2
2	LGS	A	1446	-	-	0/7/42/42	0/2/2/2
2	LGS	D	1446	-	-	1/7/42/42	0/2/2/2
2	LGS	C	1446	-	-	5/7/42/42	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1446	LGS	C7-N2	7.91	1.34	1.26
2	A	1446	LGS	C7-N2	7.28	1.34	1.26
2	B	1446	LGS	C7-N2	6.61	1.33	1.26
2	C	1446	LGS	C7-N2	5.02	1.31	1.26
2	C	1446	LGS	O1-C1	3.50	1.45	1.40
2	D	1446	LGS	O1-C1	2.94	1.44	1.40
2	A	1446	LGS	O1-C1	2.94	1.44	1.40
2	B	1446	LGS	C2-C1	2.84	1.56	1.53
2	C	1446	LGS	C2-C1	2.23	1.55	1.53
2	A	1446	LGS	C2-C1	2.19	1.55	1.53

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1446	LGS	C6-C5-N1	6.67	105.19	99.99
2	C	1446	LGS	C6-C5-N1	6.28	104.89	99.99
2	C	1446	LGS	C3-C4-C5	-5.99	101.24	111.37
2	C	1446	LGS	C9-C8-N2	-5.81	100.38	110.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	1446	LGS	C6-C5-N1	5.71	104.44	99.99
2	D	1446	LGS	C3-C4-C5	-5.20	102.59	111.37
2	B	1446	LGS	C3-C4-C5	-4.76	103.33	111.37
2	B	1446	LGS	O1-C1-N1	-4.01	102.80	111.40
2	D	1446	LGS	C3-C2-C1	3.84	115.42	109.01
2	B	1446	LGS	C8-N2-C7	3.74	128.53	121.05
2	D	1446	LGS	O3-C3-C2	-3.67	101.72	110.38
2	C	1446	LGS	C3-C2-C1	3.66	115.11	109.01
2	A	1446	LGS	C8-N2-C7	3.58	128.20	121.05
2	D	1446	LGS	C8-N2-C7	3.55	128.15	121.05
2	C	1446	LGS	C10-C9-C8	-3.52	101.10	113.38
2	A	1446	LGS	O1-C1-N1	-3.42	104.06	111.40
2	D	1446	LGS	C6-C5-N1	3.31	102.57	99.99
2	B	1446	LGS	C3-C2-C1	3.17	114.30	109.01
2	A	1446	LGS	C3-C4-C5	-3.17	106.01	111.37
2	C	1446	LGS	O3-C3-C2	-3.11	103.04	110.38
2	C	1446	LGS	O1-C1-N1	-3.00	104.95	111.40
2	D	1446	LGS	O1-C1-N1	-2.51	106.02	111.40
2	C	1446	LGS	C11-C10-C9	-2.47	101.87	114.37
2	C	1446	LGS	C12-C11-C10	-2.47	101.87	114.37
2	A	1446	LGS	C3-C2-C1	2.33	112.90	109.01
2	C	1446	LGS	C13-C12-C11	-2.32	102.65	114.37
2	B	1446	LGS	O3-C3-C2	-2.26	105.05	110.38
2	B	1446	LGS	C9-C8-N2	-2.17	106.82	110.67
2	C	1446	LGS	O2-C2-C1	-2.01	102.81	109.62

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1446	LGS	N2-C8-C9-C10
2	C	1446	LGS	C9-C10-C11-C12
2	C	1446	LGS	C11-C10-C9-C8
2	C	1446	LGS	C10-C11-C12-C13
2	D	1446	LGS	C11-C12-C13-C14
2	C	1446	LGS	C12-C13-C14-C15
2	B	1446	LGS	C12-C13-C14-C15


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1447	ACT	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand LGS A 1446



Ligand LGS D 1446







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	443/468 (94%)	-0.58	1 (0%) 91 93	8, 16, 28, 39	18 (4%)
1	B	443/468 (94%)	-0.49	0 100 100	9, 19, 32, 41	9 (2%)
1	C	443/468 (94%)	-0.84	0 100 100	4, 10, 23, 38	22 (4%)
1	D	443/468 (94%)	-0.70	1 (0%) 91 93	6, 14, 28, 44	15 (3%)
All	All	1772/1872 (94%)	-0.65	2 (0%) 92 93	4, 15, 29, 44	64 (3%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	304	PRO	2.9
1	A	3	VAL	2.6

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	ACT	C	1447	4/4	0.88	0.08	23,24,25,26	0
2	LGS	C	1446	22/22	0.97	0.05	5,8,20,24	0
2	LGS	A	1446	22/22	0.98	0.04	8,11,25,26	0
2	LGS	D	1446	22/22	0.98	0.05	6,10,24,29	0
2	LGS	B	1446	22/22	0.98	0.05	11,14,27,33	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LGS C 1446:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LGS A 1446:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LGS D 1446:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LGS B 1446:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers

There are no such residues in this entry.