



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 23, 2026 – 05:33 pm BST

PDB ID : 2WJM / pdb_00002wjm
Title : Lipidic sponge phase crystal structure of the photosynthetic reaction centre from *Blastochloris viridis* (low dose)
Authors : Woehri, A.B.; Wahlgren, W.Y.; Malmerberg, E.; Johansson, L.C.; Neutze, R.; Katona, G.
Deposited on : 2009-05-27
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	336	4% 94% 5%
2	H	258	24% 90% 6%
3	L	274	19% 93% 7%
4	M	324	3% 94% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	HEC	C	401	X	-	-	-
5	HEC	C	402	X	-	-	-
5	HEC	C	403	X	-	-	-
5	HEC	C	404	X	-	-	-
7	BCB	L	301	X	-	-	-
7	BCB	L	302	X	-	-	-
7	BCB	M	401	X	-	-	-
7	BCB	M	402	X	-	-	-

2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 10671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOTOSYNTHETIC REACTION CENTER CYTOCHROME C SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	332	Total	C	N	O	S	0	0	0
			2598	1637	465	478	18			

- Molecule 2 is a protein called REACTION CENTER PROTEIN H CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	243	Total	C	N	O	S	0	0	0
			1880	1205	326	347	2			


- Molecule 3 is a protein called REACTION CENTER PROTEIN L CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	273	Total	C	N	O	S	0	2	0
			2170	1458	350	355	7			


- Molecule 4 is a protein called REACTION CENTER PROTEIN M CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	M	323	Total	C	N	O	S	0	0	0
			2548	1697	417	423	11			

- Molecule 5 is HEME C (CCD ID: HEC) (formula: C₃₄H₃₆FeN₄O₄).




- Molecule 6 is DIACYL GLYCEROL (CCD ID: DGA) (formula: $C_{39}H_{76}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			37	33	4		

- Molecule 7 is BACTERIOCHLOROPHYLL B (CCD ID: BCB) (formula: C₅₅H₇₂MgN₄O₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
7	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
7	M	1	Total	C	Mg	N	O	0	0
			65	54	1	4	6		
7	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 8 is BACTERIOPHEOPHYTIN B (CCD ID: BPB) (formula: C₅₅H₇₄N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	L	1	Total	C	N	O	0	0
			65	55	4	6		
8	M	1	Total	C	N	O	0	0
			61	51	4	6		

- Molecule 9 is [(Z)-octadec-9-enyl] (2R)-2,3-bis(oxidanyl)propanoate (CCD ID: MPG) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	L	1	Total	C	O	0	0
			25	21	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	L	1	Total C O 25 21 4	0	0
9	M	1	Total C 17 17	0	0

- Molecule 10 is FE (II) ION (CCD ID: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	L	1	Total Fe 1 1	0	0

- Molecule 11 is MENAQUINONE-7 (CCD ID: MQ7) (formula: C₄₆H₆₄O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	M	1	Total C O 48 46 2	0	0

- Molecule 12 is 15-cis-1,2-dihydroneurosporene (CCD ID: NS5) (formula: C₄₀H₆₀).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	M	1	Total	C		0	0
			40	40			

- Molecule 13 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	M	1	Total	O	P	0	0
			5	4	1		
13	M	1	Total	O	P	0	0
			5	4	1		

- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	C	311	Total 311	O 311	0	0
14	H	146	Total 146	O 146	0	0
14	L	92	Total 92	O 92	0	0
14	M	162	Total 162	O 162	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PHOTOSYNTHETIC REACTION CENTER CYTOCHROME C SUBUNIT




• Molecule 2: REACTION CENTER PROTEIN H CHAIN



• Molecule 3: REACTION CENTER PROTEIN L CHAIN



• Molecule 4: REACTION CENTER PROTEIN M CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.45Å 138.50Å 177.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	176.78 – 1.95 176.78 – 1.95	Depositor EDS
% Data completeness (in resolution range)	95.8 (176.78-1.95) 95.9 (176.78-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.173 , 0.201 0.186 , 0.213	Depositor DCC
R_{free} test set	7313 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10671	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, BCB, PO4, FME, FE2, BPB, MPG, NS5, MQ7, DGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.87	1/2665 (0.0%)	0.92	3/3633 (0.1%)
2	H	0.75	0/1913	0.81	1/2613 (0.0%)
3	L	0.79	0/2263	0.87	1/3089 (0.0%)
4	M	0.82	0/2652	0.87	2/3630 (0.1%)
All	All	0.81	1/9493 (0.0%)	0.87	7/12965 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	231	ALA	CA-CB	5.38	1.61	1.53

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	48	GLU	N-CA-C	-8.50	94.15	108.26
1	C	3	GLU	CA-C-N	-6.18	115.21	119.66
1	C	3	GLU	C-N-CA	-6.18	115.21	119.66
2	H	12	ILE	N-CA-C	5.93	116.11	110.42
4	M	100	TYR	N-CA-C	-5.82	105.38	112.88

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	248	HIS	Peptide
1	C	47	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2598	0	2567	9	0
2	H	1880	0	1869	6	0
3	L	2170	0	2100	13	0
4	M	2548	0	2432	14	0
5	C	172	0	120	1	0
6	C	37	0	58	0	0
7	L	132	0	144	6	0
7	M	131	0	140	4	0
8	L	65	0	74	7	0
8	M	61	0	63	6	0
9	L	50	0	80	3	0
9	M	17	0	31	0	0
10	L	1	0	0	0	0
11	M	48	0	64	0	0
12	M	40	0	60	7	0
13	M	10	0	0	0	0
14	C	311	0	0	1	0
14	H	146	0	0	2	0
14	L	92	0	0	0	0
14	M	162	0	0	0	0
All	All	10671	0	9802	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:M:403:BPB:HHC	8:M:403:BPB:HBBB	1.44	0.98
7:M:401:BCB:HBB2	7:M:401:BCB:HHC	1.53	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:L:303:BPB:HHC	8:L:303:BPB:HBBB	1.55	0.88
2:H:142:ASP:OD1	14:H:301:HOH:O	2.04	0.73
7:M:401:BCB:HHC	7:M:401:BCB:CBB	2.19	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	330/336 (98%)	321 (97%)	9 (3%)	0	100	100
2	H	239/258 (93%)	235 (98%)	4 (2%)	0	100	100
3	L	273/274 (100%)	266 (97%)	7 (3%)	0	100	100
4	M	321/324 (99%)	312 (97%)	9 (3%)	0	100	100
All	All	1163/1192 (98%)	1134 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	280/282 (99%)	280 (100%)	0	100	100
2	H	192/212 (91%)	188 (98%)	4 (2%)	47	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	L	218/219 (100%)	214 (98%)	4 (2%)	51	47
4	M	247/250 (99%)	244 (99%)	3 (1%)	63	61
All	All	937/963 (97%)	926 (99%)	11 (1%)	63	61

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	L	249	ILE
4	M	29	VAL
4	M	214	PHE
4	M	181	ILE
3	L	16	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	L	183	ASN
3	L	213	ASN
4	M	16	HIS
1	C	302	GLN
1	C	120	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FME	H	1	2	8,9,10	0.83	1 (12%)	7,9,11	3.07	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FME	H	1	2	-	4/7/9/11	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	1	FME	CN-N	2.05	1.40	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	FME	CA-N-CN	-7.35	111.52	122.82
2	H	1	FME	CE-SD-CG	2.25	108.13	100.40

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	1	FME	O1-CN-N-CA
2	H	1	FME	C-CA-CB-CG
2	H	1	FME	N-CA-CB-CG
2	H	1	FME	CB-CG-SD-CE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	1	FME	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 19 ligands modelled in this entry, 1 is monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	MQ7	M	404	-	49,49,49	1.44	2 (4%)	60,63,63	1.18	6 (10%)
7	BCB	L	301	3	68,74,74	1.94	17 (25%)	79,115,115	3.13	28 (35%)
5	HEC	C	401	1	50,50,50	1.94	10 (20%)	68,82,82	1.62	9 (13%)
7	BCB	L	302	3	68,74,74	1.86	15 (22%)	79,115,115	2.98	24 (30%)
8	BPB	L	303	-	57,70,70	2.48	15 (26%)	56,101,101	2.30	15 (26%)
5	HEC	C	402	1	50,50,50	1.90	7 (14%)	68,82,82	1.58	12 (17%)
8	BPB	M	403	-	53,66,70	2.54	14 (26%)	51,96,101	2.57	17 (33%)
7	BCB	M	402	4	68,74,74	1.91	15 (22%)	79,115,115	2.99	25 (31%)
9	MPG	L	305	-	24,24,24	1.50	2 (8%)	24,25,25	1.64	3 (12%)
12	NS5	M	405	-	39,39,39	2.17	5 (12%)	44,46,46	2.16	15 (34%)
9	MPG	L	304	-	24,24,24	1.40	2 (8%)	24,25,25	1.51	2 (8%)
9	MPG	M	406	-	16,16,24	0.27	0	15,15,25	0.62	0
5	HEC	C	404	1	50,50,50	1.83	7 (14%)	68,82,82	1.32	7 (10%)
13	PO4	M	408	-	4,4,4	0.61	0	6,6,6	1.85	2 (33%)
6	DGA	C	405	1	36,36,43	1.28	2 (5%)	38,38,45	1.20	2 (5%)
7	BCB	M	401	4	67,73,74	1.98	16 (23%)	77,113,115	3.12	17 (22%)
5	HEC	C	403	1	50,50,50	1.95	9 (18%)	68,82,82	1.62	11 (16%)
13	PO4	M	407	-	4,4,4	0.64	0	6,6,6	1.50	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	MQ7	M	404	-	-	3/41/61/61	0/2/2/2
7	BCB	L	301	3	3/3/21/26	5/41/137/137	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEC	C	401	1	1/1/3/7	6/14/54/54	-
7	BCB	L	302	3	3/3/21/26	8/41/137/137	-
8	BPB	L	303	-	-	7/37/105/105	0/5/6/6
5	HEC	C	402	1	1/1/3/7	7/14/54/54	-
8	BPB	M	403	-	-	17/33/101/105	0/5/6/6
7	BCB	M	402	4	3/3/21/26	7/41/137/137	-
9	MPG	L	305	-	-	13/25/25/25	-
12	NS5	M	405	-	-	13/43/43/43	-
9	MPG	L	304	-	-	16/25/25/25	-
9	MPG	M	406	-	-	3/14/14/25	-
5	HEC	C	404	1	1/1/3/7	3/14/54/54	-
6	DGA	C	405	1	-	15/37/37/45	-
7	BCB	M	401	4	3/3/20/26	6/40/136/137	-
5	HEC	C	403	1	1/1/3/7	4/14/54/54	-

The worst 5 of 138 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	M	403	BPB	C1B-C2B	8.74	1.49	1.39
12	M	405	NS5	C35-C36	7.73	1.54	1.32
8	L	303	BPB	CAC-C3C	7.62	1.52	1.33
5	C	404	HEC	C3D-C2D	7.50	1.52	1.36
11	M	404	MQ7	C3-C2	7.45	1.48	1.35

The worst 5 of 197 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	M	401	BCB	C1C-NC-C4C	-20.68	97.41	106.71
7	L	302	BCB	C1C-NC-C4C	-17.91	98.66	106.71
7	L	301	BCB	C1C-NC-C4C	-17.81	98.70	106.71
7	M	402	BCB	C1C-NC-C4C	-17.02	99.05	106.71
7	L	301	BCB	C2B-C1B-NB	7.59	115.32	110.23

5 of 16 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	C	401	HEC	NA
5	C	402	HEC	NA
5	C	403	HEC	NA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
5	C	404	HEC	NA
7	L	301	BCB	ND

5 of 133 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	C	405	DGA	CA2-CA1-OG1-CG1
6	C	405	DGA	OA1-CA1-OG1-CG1
6	C	405	DGA	OG1-CG1-CG2-OG2
6	C	405	DGA	OG1-CG1-CG2-CG3
7	L	302	BCB	C2C-C3C-CAC-CBC


There are no ring outliers.


9 monomers are involved in 31 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	L	301	BCB	3	0
7	L	302	BCB	3	0
8	L	303	BPB	7	0
5	C	402	HEC	1	0
8	M	403	BPB	6	0
7	M	402	BCB	1	0
12	M	405	NS5	7	0
9	L	304	MPG	3	0
7	M	401	BCB	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	332/336 (98%)	0.65	12 (3%) 46 53	27, 32, 38, 53	0
2	H	242/258 (93%)	1.47	63 (26%) 1 1	26, 31, 43, 58	0
3	L	273/274 (99%)	1.34	52 (19%) 3 3	21, 32, 41, 54	2 (0%)
4	M	323/324 (99%)	0.49	10 (3%) 51 58	27, 31, 40, 43	0
All	All	1170/1192 (98%)	0.94	137 (11%) 9 11	21, 31, 41, 58	2 (0%)

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	63	ALA	6.7
2	H	83	PRO	6.6
2	H	84	GLU	6.2
2	H	85	THR	5.9
3	L	1	ALA	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FME	H	1	10/11	0.66	0.21	31,36,49,51	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	DGA	C	405	37/44	0.72	0.26	54,66,76,76	0
13	PO4	M	407	5/5	0.77	0.18	50,52,60,61	0
9	MPG	L	304	25/25	0.78	0.23	47,56,76,76	0
9	MPG	L	305	25/25	0.79	0.26	34,43,48,49	25
12	NS5	M	405	40/40	0.82	0.21	38,50,65,66	0
9	MPG	M	406	17/25	0.83	0.21	53,56,57,57	0
13	PO4	M	408	5/5	0.90	0.30	42,45,49,54	0
11	MQ7	M	404	48/48	0.92	0.12	26,30,44,46	0
8	BPB	L	303	65/65	0.94	0.10	22,29,34,36	0
7	BCB	L	302	66/66	0.94	0.11	22,25,49,52	0
8	BPB	M	403	61/65	0.95	0.11	19,25,56,58	0
7	BCB	L	301	66/66	0.96	0.09	19,25,35,38	0
7	BCB	M	401	65/66	0.96	0.11	20,23,65,67	0
7	BCB	M	402	66/66	0.96	0.09	19,22,35,36	0
5	HEC	C	401	43/43	0.97	0.09	26,30,32,33	0
5	HEC	C	404	43/43	0.97	0.09	20,24,35,43	0
5	HEC	C	403	43/43	0.98	0.08	19,23,25,27	0
5	HEC	C	402	43/43	0.98	0.09	23,26,32,35	0
10	FE2	L	306	1/1	0.99	0.07	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around DGA C 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MPG L 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MPG L 305:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NS5 M 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MPG M 406:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around MQ7 M 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BPB L 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around BCB L 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BPB M 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around BCB M 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB M 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC C 404:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.