



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 01:40 AM UTC

PDB ID : 1WPB / pdb\_00001wpb  
Title : Structure of Escherichia coli yfbU gene product  
Authors : Borek, D.; Chen, Y.; Zheng, M.; Skarina, T.; Savchenko, A.; Edwards, A.;  
Otwinowski, Z.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2004-09-01  
Resolution : 2.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*







The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	
1	B	172	
1	C	172	
1	D	172	
1	E	172	
1	F	172	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	172	
1	H	172	
1	I	172	
1	J	172	
1	K	172	
1	L	172	
1	M	172	
1	N	172	
1	O	172	
1	P	172	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	C	2011	-	-	-	X
2	CL	E	2013	-	-	-	X
2	CL	O	2036	-	-	-	X
3	GOL	B	3003	-	X	-	-
3	GOL	C	3036	-	-	X	-
3	GOL	D	3005	-	-	X	-
3	GOL	E	3004	-	-	X	-
3	GOL	G	3001	-	-	X	-
3	GOL	H	3006	-	-	X	-
3	GOL	K	3011	-	-	X	-
3	GOL	M	3043	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 24406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein yfbU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	168	1410	879	257	261	13	0	2	0
1	B	168	1408	878	254	263	13	0	2	0
1	C	168	1403	875	254	261	13	0	1	0
1	D	168	1405	876	256	260	13	0	1	0
1	E	168	1425	887	260	265	13	0	5	0
1	F	168	1405	876	254	262	13	0	2	0
1	G	168	1407	877	254	263	13	0	2	0
1	H	168	1403	875	254	261	13	0	1	0
1	I	168	1403	875	254	261	13	0	1	0
1	J	168	1403	875	254	261	13	0	1	0
1	K	168	1406	877	256	260	13	0	1	0
1	L	168	1407	877	255	262	13	0	2	0
1	M	167	1394	870	252	259	13	0	1	0
1	N	168	1403	875	254	261	13	0	1	0
1	O	168	1403	875	254	261	13	0	1	0
1	P	168	1403	875	254	261	13	0	1	0

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Cl 1	0	0
2	B	1	Total 1	Cl 1	0	0
2	C	5	Total 5	Cl 5	0	0
2	D	3	Total 3	Cl 3	0	0
2	E	2	Total 2	Cl 2	0	0
2	F	3	Total 3	Cl 3	0	0
2	G	1	Total 1	Cl 1	0	0
2	H	3	Total 3	Cl 3	0	0
2	I	4	Total 4	Cl 4	0	0
2	J	3	Total 3	Cl 3	0	0
2	K	2	Total 2	Cl 2	0	0
2	L	2	Total 2	Cl 2	0	0
2	M	2	Total 2	Cl 2	0	0
2	N	1	Total 1	Cl 1	0	0
2	O	2	Total 2	Cl 2	0	0
2	P	1	Total 1	Cl 1	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	E	1	Total C O 6 3 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total 6	C 3	O 3	0	0
3	E	1	Total 6	C 3	O 3	0	0
3	E	1	Total 6	C 3	O 3	0	0
3	F	1	Total 6	C 3	O 3	0	0
3	G	1	Total 6	C 3	O 3	0	0
3	G	1	Total 6	C 3	O 3	0	0
3	H	1	Total 6	C 3	O 3	0	0
3	H	1	Total 6	C 3	O 3	0	0
3	H	1	Total 6	C 3	O 3	0	0
3	I	1	Total 6	C 3	O 3	0	0
3	I	1	Total 6	C 3	O 3	0	0
3	J	1	Total 6	C 3	O 3	0	0
3	J	1	Total 6	C 3	O 3	0	0
3	K	1	Total 6	C 3	O 3	0	0
3	K	1	Total 6	C 3	O 3	0	0
3	K	1	Total 6	C 3	O 3	0	0
3	K	1	Total 6	C 3	O 3	0	0
3	L	1	Total 6	C 3	O 3	0	0
3	L	1	Total 6	C 3	O 3	0	0
3	M	1	Total 6	C 3	O 3	0	0
3	M	1	Total 6	C 3	O 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	M	1	Total	C	O	0	0
			6	3	3		
3	N	1	Total	C	O	0	0
			6	3	3		
3	N	1	Total	C	O	0	0
			6	3	3		
3	N	1	Total	C	O	0	0
			6	3	3		
3	N	1	Total	C	O	0	0
			6	3	3		
3	O	1	Total	C	O	0	0
			6	3	3		
3	O	1	Total	C	O	0	0
			6	3	3		
3	P	1	Total	C	O	0	0
			6	3	3		
3	P	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	96	Total	O	0	0
			96	96		
4	B	96	Total	O	0	0
			96	96		
4	C	103	Total	O	0	0
			103	103		
4	D	110	Total	O	0	0
			110	110		
4	E	88	Total	O	0	0
			88	88		
4	F	85	Total	O	0	0
			85	85		
4	G	89	Total	O	0	0
			89	89		
4	H	104	Total	O	0	0
			104	104		
4	I	130	Total	O	0	0
			130	130		
4	J	121	Total	O	0	0
			121	121		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	K	117	Total 117	O 117	0	0
4	L	102	Total 102	O 102	0	0
4	M	92	Total 92	O 92	0	0
4	N	92	Total 92	O 92	0	0
4	O	85	Total 85	O 85	0	0
4	P	108	Total 108	O 108	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: hypothetical protein yfbU




- Molecule 1: hypothetical protein yfbU




- Molecule 1: hypothetical protein yfbU



- Molecule 1: hypothetical protein yfbU





- Molecule 1: hypothetical protein yfbU








- Molecule 1: hypothetical protein yfbU





- Molecule 1: hypothetical protein yfbU




- Molecule 1: hypothetical protein yfbU





- Molecule 1: hypothetical protein yfbU




- Molecule 1: hypothetical protein yfbU




- Molecule 1: hypothetical protein yfbU




- Molecule 1: hypothetical protein yfbU




• Molecule 1: hypothetical protein yfbU




• Molecule 1: hypothetical protein yfbU



• Molecule 1: hypothetical protein yfbU



• Molecule 1: hypothetical protein yfbU



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	230.52Å 230.52Å 230.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.00 15.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-2.00) 99.7 (15.00-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.45 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.190 , 0.227 (Not available) , 0.227	Depositor DCC
$R_{free}$ test set	1481 reflections (0.50%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.4	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 55.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.015 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	24406	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.07	2/1447 (0.1%)	0.97	4/1950 (0.2%)
1	B	1.05	1/1445 (0.1%)	0.97	1/1948 (0.1%)
1	C	1.06	1/1436 (0.1%)	1.02	2/1936 (0.1%)
1	D	1.10	1/1439 (0.1%)	1.06	4/1939 (0.2%)
1	E	1.04	2/1476 (0.1%)	1.05	1/1988 (0.1%)
1	F	0.97	0/1443	0.93	0/1946
1	G	0.94	1/1445 (0.1%)	0.96	2/1948 (0.1%)
1	H	1.06	0/1436	1.03	1/1936 (0.1%)
1	I	1.13	1/1436 (0.1%)	0.98	0/1936
1	J	1.02	0/1436	0.97	1/1936 (0.1%)
1	K	1.07	1/1439 (0.1%)	1.01	2/1939 (0.1%)
1	L	1.14	1/1444 (0.1%)	1.03	2/1947 (0.1%)
1	M	1.06	1/1427 (0.1%)	1.05	5/1924 (0.3%)
1	N	1.00	0/1436	0.98	2/1936 (0.1%)
1	O	1.02	1/1436 (0.1%)	0.96	2/1936 (0.1%)
1	P	1.08	1/1436 (0.1%)	1.01	2/1936 (0.1%)
All	All	1.05	14/23057 (0.1%)	1.00	31/31081 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	D	1	3
1	E	0	2
1	G	0	2
1	H	0	1
1	K	0	1

*Continued on next page...*

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1
1	O	0	1
1	P	0	2
All	All	1	15

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	M	23	MET	SD-CE	-8.26	1.58	1.79
1	G	23	MET	SD-CE	-6.92	1.62	1.79
1	I	23	MET	SD-CE	-6.11	1.64	1.79
1	A	103	ALA	CA-CB	-5.85	1.44	1.53
1	E	103	ALA	CA-CB	-5.65	1.44	1.53

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	88	GLN	N-CA-C	8.79	129.51	110.80
1	B	47	GLY	N-CA-C	7.59	121.84	112.73
1	P	129	GLY	N-CA-C	7.53	125.47	114.10
1	D	89	GLN	N-CA-C	7.36	120.97	107.73
1	M	14	ARG	NE-CZ-NH2	7.35	125.81	119.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	88	GLN	CA

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	89	GLN	Peptide
1	C	129	GLY	Peptide
1	D	86	GLN	Peptide
1	D	87	ASP	Peptide
1	D	88	GLN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1410	0	1353	11	3
1	B	1408	0	1346	7	0
1	C	1403	0	1344	12	0
1	D	1405	0	1351	11	0
1	E	1425	0	1366	16	0
1	F	1405	0	1347	11	0
1	G	1407	0	1346	8	0
1	H	1403	0	1344	7	0
1	I	1403	0	1344	6	0
1	J	1403	0	1344	8	0
1	K	1406	0	1351	13	0
1	L	1407	0	1346	7	0
1	M	1394	0	1336	10	0
1	N	1403	0	1344	16	0
1	O	1403	0	1344	6	0
1	P	1403	0	1344	14	0
2	A	1	0	0	1	0
2	B	1	0	0	0	0
2	C	5	0	0	0	0
2	D	3	0	0	0	0
2	E	2	0	0	0	0
2	F	3	0	0	1	0
2	G	1	0	0	0	0
2	H	3	0	0	1	0
2	I	4	0	0	0	0
2	J	3	0	0	0	0
2	K	2	0	0	0	0
2	L	2	0	0	0	0
2	M	2	0	0	0	0
2	N	1	0	0	0	0
2	O	2	0	0	0	0
2	P	1	0	0	0	0
3	A	12	0	16	3	0
3	B	30	0	39	4	0
3	C	12	0	16	4	1
3	D	24	0	32	4	0
3	E	24	0	32	7	2
3	F	6	0	8	0	0
3	G	12	0	16	5	0
3	H	18	0	24	6	0
3	I	12	0	16	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	12	0	16	0	0
3	K	24	0	32	7	0
3	L	12	0	16	1	0
3	M	18	0	23	5	1
3	N	24	0	31	4	0
3	O	12	0	16	0	0
3	P	12	0	15	3	0
4	A	96	0	0	1	0
4	B	96	0	0	1	0
4	C	103	0	0	1	1
4	D	110	0	0	2	0
4	E	88	0	0	2	0
4	F	85	0	0	0	0
4	G	89	0	0	2	0
4	H	104	0	0	1	0
4	I	130	0	0	3	0
4	J	121	0	0	0	0
4	K	117	0	0	0	0
4	L	102	0	0	1	0
4	M	92	0	0	3	0
4	N	92	0	0	0	1
4	O	85	0	0	0	0
4	P	108	0	0	0	1
All	All	24406	0	21898	167	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:GLN:NE2	3:B:3015:GOL:O3	1.95	0.99
1:N:88:GLN:O	1:N:90:SER:N	2.01	0.92
1:E:156:GLN:NE2	3:E:3002:GOL:O1	2.09	0.84
1:C:156:GLN:HE22	3:C:3036:GOL:H32	1.44	0.82
1:K:49:GLN:HE21	3:K:3011:GOL:C1	1.92	0.81

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:3043:GOL:O1	4:N:3085:HOH:O[5_555]	1.84	0.36
1:A:156:GLN:OE1	3:E:3004:GOL:O2[9_555]	1.85	0.35
1:A:52:GLU:OE1	3:C:3036:GOL:O2[5_555]	2.16	0.04
1:A:156:GLN:NE2	3:E:3004:GOL:O1[9_555]	2.16	0.04
4:C:3070:HOH:O	4:P:3141:HOH:O[10_646]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/172 (98%)	162 (96%)	4 (2%)	2 (1%)	10	6
1	B	168/172 (98%)	165 (98%)	2 (1%)	1 (1%)	21	17
1	C	167/172 (97%)	163 (98%)	4 (2%)	0	100	100
1	D	24/172 (14%)	23 (96%)	1 (4%)	0	100	100
1	E	143/172 (83%)	136 (95%)	6 (4%)	1 (1%)	18	14
1	F	168/172 (98%)	162 (96%)	5 (3%)	1 (1%)	21	17
1	G	168/172 (98%)	167 (99%)	1 (1%)	0	100	100
1	H	167/172 (97%)	161 (96%)	4 (2%)	2 (1%)	10	6
1	I	167/172 (97%)	166 (99%)	1 (1%)	0	100	100
1	J	167/172 (97%)	164 (98%)	2 (1%)	1 (1%)	21	17
1	K	167/172 (97%)	165 (99%)	1 (1%)	1 (1%)	21	17
1	L	168/172 (98%)	163 (97%)	4 (2%)	1 (1%)	21	17
1	M	166/172 (96%)	162 (98%)	4 (2%)	0	100	100
1	N	167/172 (97%)	163 (98%)	3 (2%)	1 (1%)	21	17
1	O	167/172 (97%)	161 (96%)	6 (4%)	0	100	100
1	P	167/172 (97%)	161 (96%)	4 (2%)	2 (1%)	10	6
All	All	2509/2752 (91%)	2444 (97%)	52 (2%)	13 (0%)	24	21

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	SER
1	E	90	SER
1	H	87	ASP
1	H	88	GLN
1	K	90	SER

### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 80 ligands modelled in this entry, 36 are monoatomic - leaving 44 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	B	3017	-	5,5,5	0.51	0	5,5,5	0.89	0
3	GOL	N	3029	-	5,5,5	0.73	0	5,5,5	0.61	0
3	GOL	N	3012	-	5,5,5	1.24	1 (20%)	5,5,5	1.64	2 (40%)
3	GOL	E	3022	-	5,5,5	0.78	0	5,5,5	1.45	0
3	GOL	P	3034	-	5,5,5	1.27	1 (20%)	5,5,5	0.85	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	3003	-	5,5,5	1.42	1 (20%)	5,5,5	1.73	1 (20%)
3	GOL	I	3032	-	5,5,5	0.59	0	5,5,5	0.67	0
3	GOL	H	3006	-	5,5,5	0.95	0	5,5,5	1.19	1 (20%)
3	GOL	P	3014	-	5,5,5	0.28	0	5,5,5	0.86	0
3	GOL	I	3031	-	5,5,5	0.71	0	5,5,5	1.02	0
3	GOL	J	3033	-	5,5,5	0.70	0	5,5,5	1.33	1 (20%)
3	GOL	C	3036	-	5,5,5	1.04	0	5,5,5	2.24	2 (40%)
3	GOL	D	3005	-	5,5,5	0.50	0	5,5,5	1.23	1 (20%)
3	GOL	H	3016	-	5,5,5	0.84	0	5,5,5	1.24	1 (20%)
3	GOL	K	3041	-	5,5,5	0.81	0	5,5,5	1.13	0
3	GOL	L	3028	-	5,5,5	0.55	0	5,5,5	0.86	0
3	GOL	O	3026	-	5,5,5	0.87	0	5,5,5	1.36	1 (20%)
3	GOL	M	3043	-	5,5,5	0.50	0	5,5,5	0.79	0
3	GOL	O	3013	-	5,5,5	0.43	0	5,5,5	0.97	0
3	GOL	K	3010	-	5,5,5	0.63	0	5,5,5	1.11	0
3	GOL	L	3027	-	5,5,5	0.37	0	5,5,5	1.15	0
3	GOL	D	3020	-	5,5,5	0.53	0	5,5,5	0.80	0
3	GOL	A	3037	-	5,5,5	1.00	0	5,5,5	1.65	2 (40%)
3	GOL	D	3018	-	5,5,5	0.83	0	5,5,5	1.38	0
3	GOL	G	3038	-	5,5,5	0.65	0	5,5,5	0.75	0
3	GOL	E	3035	-	5,5,5	0.70	0	5,5,5	1.04	0
3	GOL	N	3008	-	5,5,5	0.86	0	5,5,5	0.41	0
3	GOL	B	3024	-	5,5,5	0.53	0	5,5,5	0.49	0
3	GOL	D	3019	-	5,5,5	0.53	0	5,5,5	0.59	0
3	GOL	G	3001	-	5,5,5	0.63	0	5,5,5	0.95	0
3	GOL	A	3040	-	5,5,5	0.95	0	5,5,5	1.41	1 (20%)
3	GOL	K	3011	-	5,5,5	0.80	0	5,5,5	0.40	0
3	GOL	H	3025	-	5,5,5	0.65	0	5,5,5	0.66	0
3	GOL	C	3021	-	5,5,5	0.56	0	5,5,5	0.51	0
3	GOL	N	3044	-	5,5,5	0.54	0	5,5,5	0.75	0
3	GOL	B	3015	-	5,5,5	0.86	0	5,5,5	1.37	1 (20%)
3	GOL	E	3002	-	5,5,5	0.91	0	5,5,5	0.63	0
3	GOL	J	3030	-	5,5,5	0.47	0	5,5,5	1.21	0
3	GOL	E	3004	-	5,5,5	0.53	0	5,5,5	0.58	0
3	GOL	M	3042	-	5,5,5	0.83	0	5,5,5	0.95	0
3	GOL	F	3039	-	5,5,5	1.05	1 (20%)	5,5,5	1.09	0
3	GOL	B	3023	-	5,5,5	0.88	0	5,5,5	1.22	0
3	GOL	M	3007	-	5,5,5	1.34	1 (20%)	5,5,5	1.81	2 (40%)
3	GOL	K	3009	-	5,5,5	0.65	0	5,5,5	1.84	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	3017	-	-	1/4/4/4	-
3	GOL	N	3029	-	-	0/4/4/4	-
3	GOL	N	3012	-	-	2/4/4/4	-
3	GOL	E	3022	-	-	0/4/4/4	-
3	GOL	P	3034	-	-	2/4/4/4	-
3	GOL	B	3003	-	-	4/4/4/4	-
3	GOL	I	3032	-	-	3/4/4/4	-
3	GOL	H	3006	-	-	2/4/4/4	-
3	GOL	P	3014	-	-	0/4/4/4	-
3	GOL	I	3031	-	-	2/4/4/4	-
3	GOL	J	3033	-	-	3/4/4/4	-
3	GOL	C	3036	-	-	2/4/4/4	-
3	GOL	D	3005	-	-	2/4/4/4	-
3	GOL	H	3016	-	-	1/4/4/4	-
3	GOL	K	3041	-	-	2/4/4/4	-
3	GOL	L	3028	-	-	2/4/4/4	-
3	GOL	O	3026	-	-	2/4/4/4	-
3	GOL	M	3043	-	-	0/4/4/4	-
3	GOL	O	3013	-	-	2/4/4/4	-
3	GOL	K	3010	-	-	4/4/4/4	-
3	GOL	L	3027	-	-	4/4/4/4	-
3	GOL	D	3020	-	-	3/4/4/4	-
3	GOL	A	3037	-	-	1/4/4/4	-
3	GOL	D	3018	-	-	3/4/4/4	-
3	GOL	G	3038	-	-	0/4/4/4	-
3	GOL	E	3035	-	-	4/4/4/4	-
3	GOL	N	3008	-	-	0/4/4/4	-
3	GOL	B	3024	-	-	0/4/4/4	-
3	GOL	D	3019	-	-	4/4/4/4	-
3	GOL	G	3001	-	-	2/4/4/4	-
3	GOL	A	3040	-	-	2/4/4/4	-
3	GOL	K	3011	-	-	2/4/4/4	-
3	GOL	H	3025	-	-	1/4/4/4	-
3	GOL	C	3021	-	-	4/4/4/4	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	N	3044	-	-	4/4/4/4	-
3	GOL	B	3015	-	-	2/4/4/4	-
3	GOL	E	3002	-	-	2/4/4/4	-
3	GOL	J	3030	-	-	4/4/4/4	-
3	GOL	E	3004	-	-	2/4/4/4	-
3	GOL	M	3042	-	-	2/4/4/4	-
3	GOL	F	3039	-	-	4/4/4/4	-
3	GOL	B	3023	-	-	2/4/4/4	-
3	GOL	M	3007	-	-	2/4/4/4	-
3	GOL	K	3009	-	-	0/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	M	3007	GOL	O2-C2	-2.95	1.34	1.43
3	N	3012	GOL	O2-C2	-2.65	1.35	1.43
3	P	3034	GOL	O2-C2	-2.48	1.36	1.43
3	B	3003	GOL	O2-C2	-2.28	1.36	1.43
3	F	3039	GOL	O2-C2	-2.13	1.37	1.43

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3036	GOL	C3-C2-C1	3.76	125.59	111.80
3	B	3003	GOL	O3-C3-C2	-3.56	94.37	110.38
3	K	3009	GOL	O1-C1-C2	-3.35	95.29	110.38
3	M	3007	GOL	C3-C2-C1	2.82	122.14	111.80
3	M	3007	GOL	O2-C2-C3	-2.80	97.58	109.18

There are no chirality outliers.

5 of 90 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	3003	GOL	O1-C1-C2-C3
3	B	3003	GOL	C1-C2-C3-O3
3	B	3023	GOL	C1-C2-C3-O3
3	C	3021	GOL	O1-C1-C2-C3
3	C	3021	GOL	C1-C2-C3-O3

There are no ring outliers.

21 monomers are involved in 57 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	N	3029	GOL	1	0
3	N	3012	GOL	1	0
3	P	3034	GOL	1	0
3	B	3003	GOL	1	0
3	H	3006	GOL	5	0
3	P	3014	GOL	2	0
3	C	3036	GOL	4	1
3	D	3005	GOL	4	0
3	H	3016	GOL	1	0
3	M	3043	GOL	3	1
3	L	3027	GOL	1	0
3	N	3008	GOL	2	0
3	G	3001	GOL	5	0
3	A	3040	GOL	3	0
3	K	3011	GOL	5	0
3	B	3015	GOL	2	0
3	E	3002	GOL	3	0
3	E	3004	GOL	4	2
3	B	3023	GOL	1	0
3	M	3007	GOL	2	0
3	K	3009	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	168/172 (97%)	0.51	18 (10%) 11 10	14, 35, 59, 73	11 (6%)
1	B	168/172 (97%)	0.49	23 (13%) 6 6	14, 35, 62, 73	10 (5%)
1	C	168/172 (97%)	0.59	23 (13%) 6 6	14, 35, 61, 71	9 (5%)
1	D	168/172 (97%)	0.46	12 (7%) 22 20	15, 34, 56, 66	10 (5%)
1	E	168/172 (97%)	0.45	19 (11%) 10 9	13, 34, 61, 73	13 (7%)
1	F	168/172 (97%)	0.36	12 (7%) 22 20	15, 35, 63, 71	14 (8%)
1	G	168/172 (97%)	0.50	17 (10%) 12 11	15, 37, 62, 70	13 (7%)
1	H	168/172 (97%)	0.42	10 (5%) 27 26	14, 33, 59, 69	14 (8%)
1	I	168/172 (97%)	0.21	10 (5%) 27 26	13, 33, 45, 59	12 (7%)
1	J	168/172 (97%)	0.31	9 (5%) 31 30	20, 35, 59, 67	9 (5%)
1	K	168/172 (97%)	0.29	10 (5%) 27 26	14, 33, 57, 68	11 (6%)
1	L	168/172 (97%)	0.64	23 (13%) 6 6	13, 33, 54, 64	13 (7%)
1	M	167/172 (97%)	0.46	16 (9%) 13 12	15, 35, 64, 76	10 (5%)
1	N	168/172 (97%)	0.36	8 (4%) 35 34	15, 35, 61, 72	10 (5%)
1	O	168/172 (97%)	0.54	13 (7%) 19 18	15, 36, 61, 73	9 (5%)
1	P	168/172 (97%)	0.37	14 (8%) 17 16	14, 34, 59, 72	12 (7%)
All	All	2687/2752 (97%)	0.43	237 (8%) 15 14	13, 35, 60, 76	180 (6%)

The worst 5 of 237 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	124[A]	THR	6.1
1	O	3	GLN	5.7
1	P	148[A]	ASN	5.7
1	M	148[A]	ASN	5.5
1	N	148[A]	ASN	5.4

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	O	3026	6/6	0.52	0.25	60,62,63,64	0
3	GOL	D	3018	6/6	0.57	0.17	50,57,58,59	0
3	GOL	B	3024	6/6	0.59	0.23	68,68,69,69	0
3	GOL	C	3036	6/6	0.61	0.21	47,49,52,55	0
3	GOL	O	3013	6/6	0.68	0.20	58,60,62,66	0
3	GOL	H	3016	6/6	0.68	0.18	49,53,56,58	0
3	GOL	B	3023	6/6	0.72	0.19	46,52,54,58	0
3	GOL	E	3035	6/6	0.73	0.17	53,57,59,60	0
3	GOL	L	3028	6/6	0.73	0.20	56,57,59,60	0
2	CL	O	2036	1/1	0.74	0.41	66,66,66,66	1
3	GOL	M	3042	6/6	0.74	0.17	47,52,54,56	0
3	GOL	A	3040	6/6	0.75	0.17	41,54,57,58	0
3	GOL	I	3032	6/6	0.76	0.16	50,55,56,57	0
3	GOL	N	3029	6/6	0.76	0.17	55,56,58,58	0
3	GOL	J	3030	6/6	0.76	0.15	46,54,56,60	0
3	GOL	B	3015	6/6	0.76	0.20	48,52,55,56	0
3	GOL	P	3014	6/6	0.76	0.14	42,47,52,55	0
3	GOL	F	3039	6/6	0.77	0.18	43,47,48,50	0
3	GOL	K	3010	6/6	0.77	0.16	55,59,60,63	0
3	GOL	M	3007	6/6	0.78	0.16	46,47,52,56	0
3	GOL	D	3019	6/6	0.78	0.17	52,54,55,56	0
3	GOL	D	3020	6/6	0.78	0.15	44,56,58,59	0
2	CL	G	2033	1/1	0.78	0.30	52,52,52,52	1
2	CL	E	2013	1/1	0.78	0.52	62,62,62,62	1
3	GOL	G	3001	6/6	0.78	0.17	49,51,54,54	0
3	GOL	P	3034	6/6	0.78	0.16	38,48,52,57	0
3	GOL	N	3008	6/6	0.79	0.20	54,55,56,60	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	J	3033	6/6	0.79	0.14	47,58,59,60	0
3	GOL	N	3044	6/6	0.79	0.24	66,68,69,70	0
2	CL	C	2011	1/1	0.79	0.51	70,70,70,70	1
3	GOL	I	3031	6/6	0.79	0.14	48,53,56,59	0
3	GOL	D	3005	6/6	0.79	0.15	49,54,59,59	0
3	GOL	G	3038	6/6	0.79	0.14	51,57,58,60	0
3	GOL	K	3009	6/6	0.80	0.12	42,44,48,53	0
3	GOL	E	3002	6/6	0.80	0.15	52,53,55,57	0
3	GOL	B	3003	6/6	0.80	0.17	35,38,44,47	0
3	GOL	C	3021	6/6	0.81	0.13	54,55,56,56	0
3	GOL	N	3012	6/6	0.81	0.15	49,50,53,54	0
2	CL	F	2034	1/1	0.81	0.39	64,64,64,64	1
2	CL	J	2016	1/1	0.81	0.47	55,55,55,55	1
3	GOL	H	3025	6/6	0.82	0.16	62,63,64,65	0
2	CL	M	2029	1/1	0.82	0.24	50,50,50,50	1
2	CL	N	2030	1/1	0.82	0.42	62,62,62,62	1
3	GOL	B	3017	6/6	0.82	0.14	55,57,60,61	0
3	GOL	E	3004	6/6	0.82	0.15	51,54,58,60	0
3	GOL	E	3022	6/6	0.82	0.16	54,55,56,57	0
3	GOL	K	3041	6/6	0.83	0.12	38,50,52,57	0
2	CL	M	2017	1/1	0.83	0.40	61,61,61,61	1
3	GOL	H	3006	6/6	0.83	0.15	45,49,53,53	0
3	GOL	A	3037	6/6	0.83	0.15	41,48,52,53	0
2	CL	A	2020	1/1	0.83	0.40	57,57,57,57	1
2	CL	J	2026	1/1	0.83	0.40	56,56,56,56	1
2	CL	H	2014	1/1	0.84	0.47	61,61,61,61	1
3	GOL	M	3043	6/6	0.84	0.15	46,51,51,55	0
3	GOL	K	3011	6/6	0.85	0.14	50,52,55,57	0
2	CL	H	2024	1/1	0.87	0.29	53,53,53,53	1
2	CL	I	2015	1/1	0.87	0.41	59,59,59,59	1
2	CL	C	2021	1/1	0.87	0.30	53,53,53,53	1
3	GOL	L	3027	6/6	0.87	0.10	44,54,57,57	0
2	CL	P	2019	1/1	0.88	0.43	53,53,53,53	1
2	CL	K	2027	1/1	0.88	0.40	52,52,52,52	1
2	CL	L	2028	1/1	0.88	0.43	58,58,58,58	1
2	CL	D	2022	1/1	0.89	0.39	52,52,52,52	1
2	CL	D	2012	1/1	0.89	0.42	55,55,55,55	1
2	CL	E	2032	1/1	0.90	0.43	58,58,58,58	1
2	CL	F	2023	1/1	0.90	0.37	53,53,53,53	1
2	CL	C	2031	1/1	0.90	0.46	61,61,61,61	1
2	CL	O	2018	1/1	0.92	0.41	55,55,55,55	1
2	CL	B	2010	1/1	0.93	0.35	54,54,54,54	1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	I	2025	1/1	0.93	0.25	47,47,47,47	1
2	CL	L	2009	1/1	0.94	0.51	57,57,57,57	1
2	CL	F	2035	1/1	0.95	0.23	42,42,42,42	1
2	CL	K	2003	1/1	0.99	0.02	25,25,25,25	0
2	CL	C	2006	1/1	0.99	0.03	27,27,27,27	0
2	CL	D	2008	1/1	0.99	0.03	24,24,24,24	0
2	CL	J	2002	1/1	0.99	0.03	21,21,21,21	0
2	CL	I	2001	1/1	0.99	0.02	19,19,19,19	0
2	CL	I	2007	1/1	0.99	0.02	24,24,24,24	0
2	CL	H	2005	1/1	1.00	0.02	20,20,20,20	0
2	CL	C	2004	1/1	1.00	0.03	23,23,23,23	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.