



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 01:18 PM UTC

PDB ID : 2WT2 / pdb_00002wt2
Title : Galectin domain of porcine adenovirus type 4 NADC-1 isolate fibre complexed with tri(N-acetyl-lactosamine)
Authors : Guardado-Calvo, P.; Munoz, E.M.; Llamas-Saiz, A.L.; Fox, G.C.; Glasgow, J.N.; van Raaij, M.J.
Deposited on : 2009-09-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	5% 76% 11% 13%
1	B	343	3% 77% 10% 13%
2	C	6	50% 50%
2	D	6	50% 33% 17%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PUTATIVE FIBER PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	0	0	0
			2354	1521	397	431	5			
1	B	300	Total	C	N	O	S	0	0	0
			2354	1521	397	431	5			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	361	MET	-	expression tag	UNP Q83467
A	362	GLY	-	expression tag	UNP Q83467
A	363	SER	-	expression tag	UNP Q83467
A	364	SER	-	expression tag	UNP Q83467
A	365	HIS	-	expression tag	UNP Q83467
A	366	HIS	-	expression tag	UNP Q83467
A	367	HIS	-	expression tag	UNP Q83467
A	368	HIS	-	expression tag	UNP Q83467
A	369	HIS	-	expression tag	UNP Q83467
A	370	HIS	-	expression tag	UNP Q83467
A	371	SER	-	expression tag	UNP Q83467
A	372	SER	-	expression tag	UNP Q83467
A	373	GLY	-	expression tag	UNP Q83467
A	374	LEU	-	expression tag	UNP Q83467
A	375	VAL	-	expression tag	UNP Q83467
A	376	PRO	-	expression tag	UNP Q83467
A	377	ARG	-	expression tag	UNP Q83467
A	378	GLY	-	expression tag	UNP Q83467
A	379	SER	-	expression tag	UNP Q83467
A	380	HIS	-	expression tag	UNP Q83467
A	381	MET	-	expression tag	UNP Q83467
A	382	ALA	-	expression tag	UNP Q83467
A	383	SER	-	expression tag	UNP Q83467
A	384	MET	-	expression tag	UNP Q83467
A	385	THR	-	expression tag	UNP Q83467

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	386	GLY	-	expression tag	UNP Q83467
A	387	GLY	-	expression tag	UNP Q83467
A	388	GLN	-	expression tag	UNP Q83467
A	389	GLN	-	expression tag	UNP Q83467
A	390	GLY	-	expression tag	UNP Q83467
A	391	ARG	-	expression tag	UNP Q83467
A	392	ILE	-	expression tag	UNP Q83467
B	361	MET	-	expression tag	UNP Q83467
B	362	GLY	-	expression tag	UNP Q83467
B	363	SER	-	expression tag	UNP Q83467
B	364	SER	-	expression tag	UNP Q83467
B	365	HIS	-	expression tag	UNP Q83467
B	366	HIS	-	expression tag	UNP Q83467
B	367	HIS	-	expression tag	UNP Q83467
B	368	HIS	-	expression tag	UNP Q83467
B	369	HIS	-	expression tag	UNP Q83467
B	370	HIS	-	expression tag	UNP Q83467
B	371	SER	-	expression tag	UNP Q83467
B	372	SER	-	expression tag	UNP Q83467
B	373	GLY	-	expression tag	UNP Q83467
B	374	LEU	-	expression tag	UNP Q83467
B	375	VAL	-	expression tag	UNP Q83467
B	376	PRO	-	expression tag	UNP Q83467
B	377	ARG	-	expression tag	UNP Q83467
B	378	GLY	-	expression tag	UNP Q83467
B	379	SER	-	expression tag	UNP Q83467
B	380	HIS	-	expression tag	UNP Q83467
B	381	MET	-	expression tag	UNP Q83467
B	382	ALA	-	expression tag	UNP Q83467
B	383	SER	-	expression tag	UNP Q83467
B	384	MET	-	expression tag	UNP Q83467
B	385	THR	-	expression tag	UNP Q83467
B	386	GLY	-	expression tag	UNP Q83467
B	387	GLY	-	expression tag	UNP Q83467
B	388	GLN	-	expression tag	UNP Q83467
B	389	GLN	-	expression tag	UNP Q83467
B	390	GLY	-	expression tag	UNP Q83467
B	391	ARG	-	expression tag	UNP Q83467
B	392	ILE	-	expression tag	UNP Q83467

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	0	0	0
			76	42	3	31			
2	D	6	Total	C	N	O	0	0	0
			76	42	3	31			

- Molecule 3 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			4	1	3		
3	A	1	Total	N	O	0	0
			4	1	3		
3	A	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	61	Total 61	O 61	0	0
4	B	68	Total 68	O 68	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PUTATIVE FIBER PROTEIN



• Molecule 1: PUTATIVE FIBER PROTEIN



• Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MCH
GAL2
NAG3
GAL4
NAG5
GAL6

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	176.13Å 38.26Å 86.52Å 90.00° 92.12° 90.00°	Depositor
Resolution (Å)	35.00 – 2.50 35.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (35.00-2.50) 99.8 (35.00-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, R_{free}	0.200 , 0.268 0.198 , 0.264	Depositor DCC
R_{free} test set	1126 reflections (5.51%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.646	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.020 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5013	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0277e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GAL, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2421	0.73	0/3313
1	B	0.60	0/2421	0.75	1/3313 (0.0%)
All	All	0.59	0/4842	0.74	1/6626 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	533	LEU	N-CA-C	5.13	117.79	110.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2354	0	2350	24	0
1	B	2354	0	2350	23	0
2	C	76	0	66	0	0
2	D	76	0	66	1	0
3	A	12	0	0	1	0
3	B	12	0	0	0	0
4	A	61	0	0	1	0
4	B	68	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5013	0	4832	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1687:NO3:O1	4:A:2059:HOH:O	2.08	0.72
1:B:508:ARG:HD2	4:B:2038:HOH:O	1.96	0.65
1:B:398:LEU:HD13	1:B:402:VAL:HG22	1.85	0.59
1:B:398:LEU:HD13	1:B:402:VAL:CG2	2.34	0.57
1:A:449:VAL:HG13	1:A:462:GLU:HG2	1.85	0.56
1:B:539:ALA:HB3	1:B:542:ASP:OD1	2.06	0.56
1:A:640:ARG:HD2	1:A:643:ASP:OD1	2.07	0.54
1:A:552:LEU:HD13	1:A:556:LEU:HD13	1.90	0.54
1:B:487:TYR:CZ	1:B:501:GLN:HG3	2.43	0.53
1:A:475:PRO:CD	1:A:538:ALA:HB2	2.39	0.53
1:B:414:LEU:HD22	1:B:418:ALA:HB1	1.91	0.52
1:A:475:PRO:HD2	1:A:538:ALA:HB2	1.90	0.52
1:B:414:LEU:HD22	1:B:418:ALA:CB	2.40	0.52
1:B:473:GLU:HA	1:B:544:ILE:HD13	1.92	0.51
1:B:558:PRO:HA	1:B:639:THR:HG22	1.94	0.49
1:A:552:LEU:HD13	1:A:556:LEU:CD1	2.43	0.49
1:B:439:ARG:CZ	1:B:449:VAL:HG21	2.43	0.49
1:A:562:LEU:HD23	1:A:682:HIS:HB2	1.95	0.48
1:B:402:VAL:CG1	1:B:482:ILE:HG13	2.44	0.48
1:A:511:LEU:HD11	1:A:515:LEU:HD12	1.95	0.48
1:A:578:ASN:HB3	1:A:580:VAL:HG13	1.96	0.48
1:A:473:GLU:HA	1:A:544:ILE:HD13	1.95	0.48
1:A:511:LEU:HD11	1:A:515:LEU:CD1	2.44	0.47
1:A:427:THR:O	1:A:427:THR:HG23	2.15	0.46
1:B:481:VAL:HG23	4:B:2019:HOH:O	2.17	0.45
1:B:578:ASN:HB3	1:B:580:VAL:HG13	1.98	0.45
1:B:639:THR:HG23	1:B:658:TYR:OH	2.17	0.45
1:B:402:VAL:HG11	1:B:482:ILE:HG13	1.99	0.44
1:B:658:TYR:CE2	1:B:663:LEU:HD11	2.52	0.44
1:B:424:ASP:OD1	1:B:432:LEU:HD13	2.18	0.44
1:A:580:VAL:HG12	1:A:590:VAL:HG13	2.00	0.44
1:A:647:LEU:N	1:A:647:LEU:HD12	2.32	0.43
1:B:573:GLU:HB3	2:D:5:NAG:H5	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:639:THR:HG23	1:A:658:TYR:OH	2.18	0.43
1:A:562:LEU:HD23	1:A:682:HIS:CB	2.49	0.42
1:B:640:ARG:CZ	1:B:645:GLN:HB2	2.50	0.42
1:A:424:ASP:OD1	1:A:432:LEU:HD13	2.19	0.42
1:B:529:ARG:N	1:B:530:PRO:CD	2.82	0.42
1:B:597:GLY:O	1:B:604:GLY:HA3	2.19	0.42
1:A:392:ILE:HD13	1:A:396:LEU:CD1	2.50	0.41
1:A:521:THR:HG21	1:A:523:TYR:CE2	2.55	0.41
1:B:502:ARG:O	1:B:503:LEU:HD23	2.21	0.41
1:A:606:ILE:HB	1:A:622:VAL:HB	2.03	0.41
1:A:474:LYS:HD2	1:A:538:ALA:HB3	2.03	0.41
1:B:427:THR:HG23	1:B:431:SER:HB2	2.03	0.41
1:A:539:ALA:HB1	1:A:540:PRO:HD2	2.02	0.40
1:A:402:VAL:HG11	1:A:482:ILE:HG13	2.02	0.40
1:A:421:PHE:CE2	1:A:438:VAL:HG21	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/343 (87%)	286 (96%)	11 (4%)	1 (0%)	36	55
1	B	298/343 (87%)	288 (97%)	10 (3%)	0	100	100
All	All	596/686 (87%)	574 (96%)	21 (4%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	429	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/302 (88%)	263 (99%)	2 (1%)	73	88
1	B	265/302 (88%)	262 (99%)	3 (1%)	65	84
All	All	530/604 (88%)	525 (99%)	5 (1%)	70	87

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	556	LEU
1	A	621	THR
1	B	404	THR
1	B	417	SER
1	B	529	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	657	GLN
1	A	679	HIS
1	B	504	GLN
1	B	655	GLN
1	B	657	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2	15,15,15	0.43	0	21,21,21	0.99	1 (4%)
2	GAL	C	2	2	11,11,12	0.70	0	15,15,17	0.69	0
2	NAG	C	3	2	14,14,15	0.58	0	17,19,21	0.94	0
2	GAL	C	4	2	11,11,12	0.72	0	15,15,17	1.08	2 (13%)
2	NAG	C	5	2	14,14,15	0.59	0	17,19,21	1.28	2 (11%)
2	GAL	C	6	2	11,11,12	0.81	0	15,15,17	0.79	0
2	NAG	D	1	2	15,15,15	0.45	0	21,21,21	0.80	0
2	GAL	D	2	2	11,11,12	0.75	0	15,15,17	0.69	0
2	NAG	D	3	2	14,14,15	0.45	0	17,19,21	0.93	0
2	GAL	D	4	2	11,11,12	0.69	0	15,15,17	1.02	2 (13%)
2	NAG	D	5	2	14,14,15	0.52	0	17,19,21	1.34	2 (11%)
2	GAL	D	6	2	11,11,12	0.95	1 (9%)	15,15,17	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2	-	0/6/26/26	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	NAG	C	3	2	-	1/6/23/26	0/1/1/1
2	GAL	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	2/6/23/26	0/1/1/1
2	GAL	C	6	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	NAG	D	3	2	-	0/6/23/26	0/1/1/1
2	GAL	D	4	2	-	0/2/19/22	0/1/1/1
2	NAG	D	5	2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	D	6	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	6	GAL	O5-C1	-2.25	1.39	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	5	NAG	O5-C1-C2	-3.84	105.35	111.29
2	C	5	NAG	O5-C1-C2	-3.74	105.50	111.29
2	D	5	NAG	C4-C3-C2	2.96	115.35	111.02
2	C	5	NAG	C4-C3-C2	2.79	115.11	111.02
2	D	4	GAL	C1-C2-C3	2.22	112.88	109.64
2	C	4	GAL	C1-C2-C3	2.15	112.78	109.64
2	C	1	NAG	O5-C1-C2	-2.06	107.45	109.52
2	C	4	GAL	C1-O5-C5	2.01	114.88	112.19
2	D	4	GAL	C1-O5-C5	2.00	114.87	112.19

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	5	NAG	C4-C5-C6-O6
2	C	5	NAG	O5-C5-C6-O6
2	C	3	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	5	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NO3	A	1688	-	1,3,3	3.41	1 (100%)	0,3,3	-	-
3	NO3	B	1686	-	1,3,3	3.13	1 (100%)	0,3,3	-	-
3	NO3	B	1687	-	1,3,3	3.06	1 (100%)	0,3,3	-	-
3	NO3	B	1688	-	1,3,3	3.69	1 (100%)	0,3,3	-	-
3	NO3	A	1686	-	1,3,3	3.09	1 (100%)	0,3,3	-	-
3	NO3	A	1687	-	1,3,3	2.98	1 (100%)	0,3,3	-	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1688	NO3	O1-N	3.69	1.42	1.24
3	A	1688	NO3	O1-N	3.41	1.41	1.24
3	B	1686	NO3	O1-N	3.13	1.39	1.24
3	A	1686	NO3	O1-N	3.09	1.39	1.24
3	B	1687	NO3	O1-N	3.06	1.39	1.24
3	A	1687	NO3	O1-N	2.98	1.39	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1687	NO3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	300/343 (87%)	-0.22	16 (5%) 32 28	11, 26, 72, 160	0
1	B	300/343 (87%)	-0.27	9 (3%) 52 48	10, 26, 64, 115	0
All	All	600/686 (87%)	-0.24	25 (4%) 40 36	10, 26, 70, 160	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	526	GLY	5.0
1	B	685	PRO	3.7
1	A	540	PRO	3.4
1	A	527	ASP	3.2
1	A	528	PRO	3.1
1	A	530	PRO	3.1
1	B	528	PRO	2.9
1	A	539	ALA	2.9
1	A	543	VAL	2.8
1	A	541	LEU	2.8
1	A	685	PRO	2.8
1	B	531	THR	2.8
1	B	526	GLY	2.7
1	A	529	ARG	2.7
1	A	531	THR	2.7
1	A	532	THR	2.6
1	B	532	THR	2.6
1	B	527	ASP	2.5
1	A	387	GLY	2.4
1	A	684	VAL	2.4
1	B	530	PRO	2.3
1	A	534	LEU	2.3
1	B	540	PRO	2.2
1	A	388	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	539	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	D	5	14/15	0.86	0.11	31,36,40,43	0
2	NAG	D	1	15/15	0.87	0.10	27,34,41,41	0
2	GAL	C	6	11/12	0.87	0.10	34,43,49,50	0
2	NAG	C	1	15/15	0.88	0.09	31,37,42,46	0
2	NAG	C	5	14/15	0.90	0.09	33,39,44,47	0
2	GAL	D	6	11/12	0.90	0.10	28,37,40,41	0
2	GAL	D	4	11/12	0.93	0.06	20,23,27,27	0
2	GAL	D	2	11/12	0.93	0.07	27,29,31,36	0
2	NAG	D	3	14/15	0.93	0.08	18,26,43,43	0
2	GAL	C	4	11/12	0.94	0.07	24,28,31,35	0
2	NAG	C	3	14/15	0.95	0.07	23,24,35,35	0
2	GAL	C	2	11/12	0.95	0.06	21,26,29,32	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NO3	B	1688	4/4	0.88	0.12	31,35,38,40	0
3	NO3	A	1688	4/4	0.94	0.07	32,36,37,39	0
3	NO3	A	1686	4/4	0.94	0.13	18,20,23,27	0
3	NO3	A	1687	4/4	0.96	0.10	20,20,21,25	0
3	NO3	B	1687	4/4	0.97	0.06	24,25,26,31	0
3	NO3	B	1686	4/4	0.97	0.07	18,19,23,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.