Full wwPDB X-ray Structure Validation Report (i)

Aug 25, 2026 – 05:42 am BST

PDB ID	:	2X80 / pdb_00002x80
Title	:	P450 BM3 F87A in complex with DMSO
Authors	:	Kuper, J.; Wong, T.S.; Roccatano, D.; Wilmanns, M.; Schwaneberg, U.
Deposited on	:	2010-03-05
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	8% 89% 10%
1	B	455	5% 89% 10% •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BIFUNCTIONAL P-450/NADPH-P450 REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	455	3665	2339	623	686	17	0	0	0
1	B	455	3665	2339	623	686	17	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	ALA	PHE	engineered mutation	UNP P14779
B	87	ALA	PHE	engineered mutation	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	1	1	1	0	0
			Total	Zn		
3	B	2	2	2	0	0

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
4	A	1	4	2	1	1	0	0
			Total	C	O	S		
4	B	1	4	2	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	178	178	178	0	0
			Total	O		
5	B	228	228	228	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BIFUNCTIONAL P-450/NADPH-P450 REDUCTASE

Chain A:





• Molecule 1: BIFUNCTIONAL P-450/NADPH-P450 REDUCTASE

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	82.21Å 96.64Å 164.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.10 – 2.30 41.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (41.10-2.30) 99.6 (41.10-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.12 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.250 0.203 , 0.248	Depositor DCC
R_{free} test set	2979 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	31.7	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 34.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	7833	wwPDB-VP
Average B, all atoms (Å^2)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, DMS, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.63	1/3749 (0.0%)	0.90	0/5068
1	B	0.66	0/3749	0.90	1/5068 (0.0%)
All	All	0.65	1/7498 (0.0%)	0.90	1/10136 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	357	ILE	CA-CB	6.33	1.58	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	357	ILE	N-CA-CB	5.99	114.27	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3665	0	3641	25	0
1	B	3665	0	3641	42	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
5	A	178	0	0	1	0
5	B	228	0	0	2	0
All	All	7833	0	7354	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:MET:HE1	1:B:409:GLU:CA	1.91	0.99
1:B:119:MET:CE	1:B:409:GLU:HB2	1.94	0.97
1:B:119:MET:HE1	1:B:409:GLU:CB	1.98	0.93
1:B:119:MET:HE1	1:B:409:GLU:HB2	1.46	0.93
1:B:361:HIS:HE1	1:B:391:LYS:H	1.24	0.84
1:B:119:MET:HE1	1:B:409:GLU:HA	1.60	0.82
1:A:100:HIS:HD2	5:A:2177:HOH:O	1.64	0.80
1:B:171:HIS:HD2	1:B:173:PHE:H	1.31	0.79
1:A:171:HIS:HD2	1:A:173:PHE:H	1.33	0.77
1:A:361:HIS:HE1	1:A:391:LYS:H	1.33	0.74
1:B:119:MET:CE	1:B:409:GLU:HA	2.18	0.74
1:B:323:ARG:HA	1:B:361:HIS:CD2	2.23	0.73
1:B:150:LEU:HD21	1:B:177:MET:HE1	1.70	0.72
1:B:119:MET:HE3	1:B:409:GLU:HB2	1.73	0.70
1:B:272:LEU:HD13	1:B:322:LEU:HG	1.76	0.68
1:B:361:HIS:CE1	1:B:391:LYS:H	2.11	0.66
1:B:122:ILE:HG22	1:B:148:LEU:HD22	1.77	0.65
1:B:100:HIS:HD2	5:B:2225:HOH:O	1.79	0.64
1:A:192:ASN:HB2	1:A:195:ASP:HB2	1.78	0.64
1:A:122:ILE:HG22	1:A:148:LEU:HD22	1.82	0.59
1:B:119:MET:CE	1:B:409:GLU:CB	2.68	0.58
1:B:203:ARG:NH2	1:B:207:GLU:OE1	2.42	0.53
1:A:361:HIS:CE1	1:A:391:LYS:H	2.22	0.52
1:B:388:HIS:HD2	1:B:391:LYS:HZ3	1.58	0.52
1:A:3:LYS:HB2	1:A:344:GLU:HB3	1.94	0.50
1:B:280:LEU:HB3	1:B:287:LEU:HG	1.94	0.50
1:A:5:MET:HE1	1:A:50:ARG:HE	1.76	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:LEU:HD13	1:A:322:LEU:HG	1.92	0.49
1:B:229:GLN:NE2	1:B:239:ASN:HD21	2.11	0.49
1:B:280:LEU:HD22	1:B:287:LEU:HA	1.95	0.48
1:B:171:HIS:CD2	1:B:173:PHE:H	2.22	0.47
1:A:148:LEU:HD11	1:A:413:VAL:HG21	1.97	0.47
1:B:388:HIS:HD2	1:B:391:LYS:NZ	2.13	0.46
1:A:7:GLN:NE2	1:A:43:GLU:HB2	2.30	0.46
1:A:127:VAL:O	1:A:131:GLU:HG2	2.15	0.46
1:A:171:HIS:CD2	1:A:173:PHE:H	2.22	0.46
1:B:403:GLN:NE2	5:B:2195:HOH:O	2.49	0.46
1:A:391:LYS:HZ3	1:A:395:ASN:HD22	1.62	0.45
1:A:109:GLN:O	1:A:112:MET:HB2	2.16	0.45
1:B:38:GLU:HB2	1:B:54:SER:HB3	1.98	0.45
1:B:323:ARG:HA	1:B:361:HIS:HD2	1.77	0.45
1:A:103:LEU:HD21	1:A:237:MET:HG3	1.99	0.45
1:B:103:LEU:HD21	1:B:237:MET:CG	2.47	0.45
1:A:20:LEU:HG	1:A:42:PHE:CZ	2.53	0.44
1:B:173:PHE:CD2	1:B:173:PHE:C	2.96	0.44
1:B:361:HIS:HE1	1:B:391:LYS:N	2.04	0.44
1:B:229:GLN:HE21	1:B:239:ASN:HD21	1.64	0.44
1:A:348:GLU:N	1:A:351:ASP:OD2	2.50	0.44
1:B:64:GLU:OE2	1:B:397:GLN:HG2	2.18	0.43
1:B:329:PRO:O	1:B:358:PRO:HD3	2.18	0.43
1:B:361:HIS:CE1	1:B:390:PHE:HA	2.53	0.43
1:A:25:PRO:HD2	1:A:435:GLU:OE1	2.19	0.42
1:A:141:VAL:O	1:A:145:MET:HG2	2.18	0.42
1:A:319:ASN:HD22	1:A:319:ASN:HA	1.74	0.42
1:B:192:ASN:HA	1:B:193:PRO:HD3	1.73	0.42
1:A:302:VAL:HA	1:A:303:PRO:HD3	1.92	0.42
1:B:286:VAL:HG11	1:B:374:PHE:HE2	1.84	0.42
1:B:302:VAL:HA	1:B:303:PRO:HD3	1.97	0.41
1:A:195:ASP:HA	1:A:196:PRO:HD3	1.93	0.41
1:B:110:GLN:CD	1:B:110:GLN:H	2.28	0.41
1:A:183:GLU:O	1:A:187:LYS:HG3	2.21	0.41
1:B:62:CYS:HB3	1:B:395:ASN:ND2	2.36	0.41
1:B:119:MET:CE	1:B:409:GLU:CA	2.73	0.41
1:B:253:ASN:O	1:B:257:GLN:HG2	2.21	0.41
1:B:112:MET:SD	1:B:405:PHE:HA	2.61	0.40
1:B:119:MET:HE1	1:B:409:GLU:N	2.31	0.40
1:A:345:TYR:HA	1:A:346:PRO:HD3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/455 (100%)	432 (95%)	19 (4%)	2 (0%)	30	38
1	B	453/455 (100%)	438 (97%)	15 (3%)	0	100	100
All	All	906/910 (100%)	870 (96%)	34 (4%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	ASP
1	A	45	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	398/398 (100%)	391 (98%)	7 (2%)	51	70
1	B	398/398 (100%)	383 (96%)	15 (4%)	29	44
All	All	796/796 (100%)	774 (97%)	22 (3%)	38	56

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	126	LEU
1	A	148	LEU
1	A	181	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	182	ASP
1	A	302	VAL
1	A	356	LEU
1	B	1	THR
1	B	110	GLN
1	B	126	LEU
1	B	148	LEU
1	B	181	LEU
1	B	192	ASN
1	B	203	ARG
1	B	204	GLN
1	B	229	GLN
1	B	263	ILE
1	B	270	SER
1	B	287	LEU
1	B	302	VAL
1	B	380	GLU
1	B	409	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	95	ASN
1	A	100	HIS
1	A	110	GLN
1	A	171	HIS
1	A	201	ASN
1	A	229	GLN
1	A	319	ASN
1	A	359	GLN
1	A	361	HIS
1	A	388	HIS
1	A	395	ASN
1	A	404	GLN
1	A	420	HIS
1	A	426	HIS
1	B	27	GLN
1	B	95	ASN
1	B	100	HIS
1	B	109	GLN
1	B	171	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	204	GLN
1	B	239	ASN
1	B	319	ASN
1	B	361	HIS
1	B	381	ASN
1	B	388	HIS
1	B	395	ASN
1	B	403	GLN
1	B	426	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	HEM	A	1456	1	50,50,50	1.97	8 (16%)	66,82,82	1.54	10 (15%)
4	DMS	B	1459	-	3,3,3	2.74	1 (33%)	3,3,3	0.50	0
2	HEM	B	1456	1	50,50,50	1.92	9 (18%)	66,82,82	1.24	5 (7%)
4	DMS	A	1458	-	3,3,3	2.84	1 (33%)	3,3,3	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1456	1	-	4/14/54/54	-
2	HEM	B	1456	1	-	2/14/54/54	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1456	HEM	C3D-C2D	7.44	1.52	1.36
2	B	1456	HEM	C3D-C2D	7.27	1.52	1.36
2	A	1456	HEM	FE-NC	5.40	2.13	1.95
2	B	1456	HEM	FE-ND	5.11	2.10	1.94
2	A	1456	HEM	FE-ND	5.08	2.10	1.94
4	A	1458	DMS	O-S	4.79	1.82	1.50
4	B	1459	DMS	O-S	4.62	1.81	1.50
2	A	1456	HEM	FE-NA	4.57	2.10	1.95
2	B	1456	HEM	FE-NB	3.74	2.06	1.94
2	B	1456	HEM	FE-NA	3.36	2.06	1.95
2	B	1456	HEM	CAC-C3C	3.02	1.55	1.47
2	B	1456	HEM	CAB-C3B	2.98	1.55	1.47
2	A	1456	HEM	CAB-C3B	2.82	1.55	1.47
2	B	1456	HEM	CMA-C3A	2.76	1.56	1.50
2	A	1456	HEM	CMC-C2C	2.59	1.56	1.50
2	B	1456	HEM	CMC-C2C	2.56	1.56	1.50
2	B	1456	HEM	CMB-C2B	2.21	1.55	1.50
2	A	1456	HEM	CAC-C3C	2.10	1.53	1.47
2	A	1456	HEM	CMD-C2D	2.03	1.55	1.50

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1456	HEM	C4D-ND-C1D	5.33	110.58	105.07
2	A	1456	HEM	C4D-ND-C1D	5.26	110.51	105.07
2	A	1456	HEM	C4C-C3C-C2C	3.54	109.66	106.75
2	A	1456	HEM	C4A-NA-C1A	3.44	108.71	105.35
2	A	1456	HEM	C4C-NC-C1C	3.38	108.66	105.35
2	A	1456	HEM	CMD-C2D-C1D	3.03	129.66	125.04
2	A	1456	HEM	C1B-NB-C4B	2.81	107.98	105.07
2	A	1456	HEM	CBC-CAC-C3C	-2.65	114.43	127.62
2	A	1456	HEM	CHC-C4B-NB	2.50	127.13	124.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1456	HEM	CAD-C3D-C4D	2.46	128.96	124.66
2	B	1456	HEM	C4A-NA-C1A	2.45	107.74	105.35
2	A	1456	HEM	CBD-CAD-C3D	-2.36	106.07	112.63
2	B	1456	HEM	CAD-CBD-CGD	-2.28	108.69	113.60
2	B	1456	HEM	CHD-C4C-NC	2.26	126.89	124.44
2	A	1456	HEM	CHA-C4D-ND	2.16	127.04	124.37

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1456	HEM	C2C-C3C-CAC-CBC
2	A	1456	HEM	C4C-C3C-CAC-CBC
2	B	1456	HEM	CAD-CBD-CGD-O1D
2	A	1456	HEM	CAD-CBD-CGD-O1D
2	A	1456	HEM	CAD-CBD-CGD-O2D
2	B	1456	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEM A 1456



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM B 1456



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	455/455 (100%)	0.47	37 (8%)	18	19	28, 43, 68, 85	0
1	B	455/455 (100%)	0.26	21 (4%)	37	39	25, 41, 66, 80	0
All	All	910/910 (100%)	0.37	58 (6%)	25	27	25, 42, 67, 85	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ILE	6.4
1	A	21	ASN	5.8
1	A	231	ASP	4.6
1	B	225	ALA	4.5
1	A	1	THR	4.4
1	B	1	THR	4.4
1	A	193	PRO	4.2
1	B	227	GLY	4.0
1	A	45	PRO	3.9
1	A	230	SER	3.8
1	B	229	GLN	3.8
1	A	191	ALA	3.7
1	A	196	PRO	3.4
1	A	192	ASN	3.1
1	A	243	PRO	3.1
1	A	455	LEU	3.1
1	A	198	TYR	3.0
1	A	245	THR	3.0
1	B	92	HIS	3.0
1	A	14	LEU	2.9
1	B	194	ASP	2.9
1	A	4	GLU	2.8
1	B	193	PRO	2.8
1	B	192	ASN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	229	GLN	2.7
1	B	196	PRO	2.7
1	A	370	ASP	2.7
1	B	228	GLU	2.6
1	A	47	ARG	2.6
1	A	44	ALA	2.5
1	A	136	ASP	2.5
1	A	197	ALA	2.5
1	A	369	ASP	2.5
1	A	10	THR	2.5
1	A	2	ILE	2.4
1	A	263	ILE	2.4
1	B	195	ASP	2.3
1	A	244	GLU	2.2
1	A	200	GLU	2.2
1	A	23	ASP	2.2
1	B	47	ARG	2.2
1	B	190	ARG	2.2
1	B	4	GLU	2.2
1	B	197	ALA	2.2
1	A	232	ASP	2.2
1	A	248	PRO	2.1
1	B	226	SER	2.1
1	B	244	GLU	2.1
1	A	11	PHE	2.1
1	A	188	LEU	2.1
1	A	384	ALA	2.1
1	A	20	LEU	2.1
1	A	169	GLN	2.1
1	A	46	GLY	2.1
1	B	3	LYS	2.0
1	B	380	GLU	2.0
1	A	227	GLY	2.0
1	B	288	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	DMS	A	1458	4/4	0.82	0.28	67,67,67,68	0
4	DMS	B	1459	4/4	0.89	0.21	68,68,68,69	0
3	ZN	B	1458	1/1	0.95	0.07	65,65,65,65	0
3	ZN	A	1457	1/1	0.97	0.07	65,65,65,65	0
2	HEM	B	1456	43/43	0.98	0.07	25,28,32,33	0
2	HEM	A	1456	43/43	0.98	0.07	28,31,34,36	0
3	ZN	B	1457	1/1	0.98	0.05	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM B 1456:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HEM A 1456:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.