wwPDB EM Validation Summary Report (i)

Aug 4, 2026 – 03:28 PM EDT

PDB ID	:	6XI0 / pdb_00006xi0
EMDB ID	:	EMD-22189
Title	:	R. capsulatus cyt bc1 (CIII2) at 3.3A
Authors	:	Steimle, S.; Van Eeuwen, T.; Ozturk, Y.; Kim, H.J.; Braitbard, M.; Selamoglu, N.; Garcia, B.A.; Schneidman-Duhovny, D.; Murakami, K.; Daldal, F.
Deposited on	:	2020-06-19
Resolution	:	3.30 Å(reported)
Based on initial model	:	1ZRT

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15087 (2.80 - 3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	191	73%; 47%; 47%; • 5%
1	R	191	73%; 46%; 48%; • 5%
2	C	437	5%; 76%; 20%; • •
2	P	437	6%; 75%; 21%; • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	279	12% 65% 18% • 16%
3	Q	279	12% 65% 18% • 16%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	HEC	C	502	X	-	-	-
5	HEC	D	501	X	-	-	-
5	HEC	P	502	X	-	-	-
5	HEC	Q	501	X	-	-	-

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 13079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquinol-cytochrome c reductase iron-sulfur subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	E	181	1365	855	246	256	8	0	0
1	R	181	1365	855	246	256	8	0	0

- Molecule 2 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	423	3244	2186	519	525	14	0	0
2	P	423	3244	2186	519	525	14	0	0

- Molecule 3 is a protein called Cytochrome c1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	D	235	1801	1148	301	337	15	0	0
3	Q	235	1794	1144	300	335	15	0	0

- Molecule 4 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			AltConf
			Total	Fe	S	
4	E	1	4	2	2	0
			Total	Fe	S	
4	R	1	4	2	2	0

- Molecule 5 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{36}FeN_4O_4$).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	Fe	N	O	
5	C	1	43	34	1	4	4	0
			Total	C	Fe	N	O	
5	C	1	43	34	1	4	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
			Total	C	Fe	N	O	
5	D	1	43	34	1	4	4	0
5	P	1	43	34	1	4	4	0
5	P	1	43	34	1	4	4	0
5	Q	1	43	34	1	4	4	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquinol-cytochrome c reductase iron-sulfur subunit

Chain E:

- Molecule 1: Ubiquinol-cytochrome c reductase iron-sulfur subunit

Chain R:

- Molecule 2: Cytochrome b

Chain C:





- Molecule 2: Cytochrome b

Chain P:



- Molecule 3: Cytochrome c1

Chain D:



- Molecule 3: Cytochrome c1

Chain Q:


MET
LYS
LYS
LEU
LEU
ILE
SER
ALA
VAL
SER
ALA
LEU
VAL
LEU
GLY
SER
GLY
ALA
ALA
PHE
ALA
ASN
SER
ASN
VAL
P5
D6
H7
I14
Y30
N31
E32
V33
C37
M40
K41
F42
V43
D50
Q55
L56
D57
Y64
I71
I72
D73
K74
D75
S76
G77
E78
E79
R80
D81
R82
M87
F88
P89
T90
R91
V92
G93
D94
S101
V102
M103
R107
ALA
GLY
PHE
SER
GLY
PRO
ALA
GLY
SER
GLY
MET
ASN
GLN
LEU
PHE
LYS
GLY
MET
G126
Y134
E139
E143
C144
A145
P146
E147
D150
K156
T157
F158
Q159
I160
D165
T166
C167
K168
D169
A170
A171
G172
V173
K174
I175
T176
H177
R182
M183
P186
L187
V188
D189
D190
Q191
V192
T193
Y194
E195
D196
G197
M205
V209
L213
A216
M241
G255
H256
K257
ALA

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	37997	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.266	Depositor
Minimum map value	-0.181	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.05	Depositor
Map size (Å)	326.4, 326.4, 326.4	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.36, 1.36, 1.36	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	E	0.35	0/1395	0.66	1/1895 (0.1%)
1	R	0.34	0/1395	0.66	1/1895 (0.1%)
2	C	0.37	0/3363	0.61	7/4617 (0.2%)
2	P	0.37	0/3363	0.62	8/4617 (0.2%)
3	D	0.28	0/1846	0.60	6/2500 (0.2%)
3	Q	0.27	0/1839	0.59	5/2492 (0.2%)
All	All	0.34	0/13201	0.62	28/18016 (0.2%)

There are no bond length outliers.

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	P	222	ASN	N-CA-C	9.89	121.65	111.07
1	E	49	SER	N-CA-C	8.95	122.07	110.53
1	R	49	SER	N-CA-C	8.95	122.07	110.53
3	D	183	MET	CA-C-N	-8.10	112.03	120.38
3	D	183	MET	C-N-CA	-8.10	112.03	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1365	0	1340	139	0
1	R	1365	0	1340	140	0
2	C	3244	0	3072	94	0
2	P	3244	0	3072	101	0
3	D	1801	0	1722	79	0
3	Q	1794	0	1709	78	0
4	E	4	0	0	0	0
4	R	4	0	0	0	0
5	C	86	0	64	15	0
5	D	43	0	30	5	0
5	P	86	0	64	17	0
5	Q	43	0	30	5	0
All	All	13079	0	12443	597	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 597 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:40:ASN:O	2:C:224:PRO:HG2	1.44	1.18
3:D:188:VAL:HG12	3:D:189:ASP:H	0.98	1.13
1:R:183:VAL:HB	1:R:187:THR:HB	1.31	1.13
2:C:199:TYR:CE1	2:P:199:TYR:HD1	1.69	1.10
2:P:199:TYR:CE2	5:P:501:HEC:HBC1	1.86	1.10

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	179/191 (94%)	141 (79%)	36 (20%)	2 (1%)	11 39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	R	179/191 (94%)	141 (79%)	36 (20%)	2 (1%)	11	39
2	C	419/437 (96%)	352 (84%)	65 (16%)	2 (0%)	24	55
2	P	419/437 (96%)	352 (84%)	66 (16%)	1 (0%)	43	71
3	D	231/279 (83%)	197 (85%)	34 (15%)	0	100	100
3	Q	231/279 (83%)	197 (85%)	34 (15%)	0	100	100
All	All	1658/1814 (91%)	1380 (83%)	271 (16%)	7 (0%)	31	60

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	99	GLU
1	R	99	GLU
1	E	144	LYS
1	R	144	LYS
2	C	223	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	142/149 (95%)	142 (100%)	0	100	100
1	R	142/149 (95%)	142 (100%)	0	100	100
2	C	301/360 (84%)	300 (100%)	1 (0%)	86	86
2	P	300/360 (83%)	300 (100%)	0	100	100
3	D	184/220 (84%)	184 (100%)	0	100	100
3	Q	182/220 (83%)	182 (100%)	0	100	100
All	All	1251/1458 (86%)	1250 (100%)	1 (0%)	87	90

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	224	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
3	Q	177	HIS
1	R	122	ASN
1	R	100	ASN
3	D	31	ASN
3	Q	31	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	HEC	Q	501	3	50,50,50	1.79	7 (14%)	68,82,82	1.24	2 (2%)
5	HEC	C	501	2	50,50,50	1.57	9 (18%)	68,82,82	1.39	9 (13%)
4	FES	R	501	1	0,4,4	-	-	-		
5	HEC	P	501	2	50,50,50	1.57	9 (18%)	68,82,82	1.39	9 (13%)
5	HEC	C	502	2	50,50,50	1.66	9 (18%)	68,82,82	1.40	9 (13%)
4	FES	E	501	1	0,4,4	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	HEC	P	502	2	50,50,50	1.66	9 (18%)	68,82,82	1.41	9 (13%)
5	HEC	D	501	3	50,50,50	1.79	7 (14%)	68,82,82	1.24	2 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEC	Q	501	3	1/1/3/7	6/14/54/54	-
5	HEC	C	501	2	-	8/14/54/54	-
4	FES	R	501	1	-	-	0/1/1/1
5	HEC	P	501	2	-	8/14/54/54	-
5	HEC	C	502	2	1/1/3/7	9/14/54/54	-
4	FES	E	501	1	-	-	0/1/1/1
5	HEC	P	502	2	1/1/3/7	9/14/54/54	-
5	HEC	D	501	3	1/1/3/7	6/14/54/54	-

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	501	HEC	C3D-C2D	7.71	1.53	1.36
5	Q	501	HEC	C3D-C2D	7.71	1.53	1.36
5	C	502	HEC	FE-NB	5.94	2.13	1.94
5	P	502	HEC	FE-NB	5.93	2.13	1.94
5	Q	501	HEC	FE-ND	5.14	2.10	1.94

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	D	501	HEC	C1D-ND-C4D	5.04	111.18	105.21
5	Q	501	HEC	C1D-ND-C4D	5.02	111.16	105.21
5	P	502	HEC	CBB-CAB-C3B	4.57	124.81	112.42
5	C	502	HEC	CBB-CAB-C3B	4.55	124.77	112.42
5	C	502	HEC	CBC-CAC-C3C	4.24	123.91	112.42

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	C	502	HEC	NA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
5	D	501	HEC	NA
5	P	502	HEC	NA
5	Q	501	HEC	NA

5 of 46 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	501	HEC	C2C-C3C-CAC-CBC
5	P	501	HEC	C2C-C3C-CAC-CBC
5	C	501	HEC	C4B-C3B-CAB-CBB
5	P	501	HEC	C4B-C3B-CAB-CBB
5	C	501	HEC	C2B-C3B-CAB-CBB

There are no ring outliers.

6 monomers are involved in 42 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	Q	501	HEC	5	0
5	C	501	HEC	12	0
5	P	501	HEC	14	0
5	C	502	HEC	3	0
5	P	502	HEC	3	0
5	D	501	HEC	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEC Q 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEC C 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEC P 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEC C 502



Bond lengths



Bond angles



Torsions



Rings

Ligand HEC P 502



Bond lengths



Bond angles



Torsions



Rings



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-22189. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 120



Y Index: 120



Z Index: 120

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 106



Y Index: 118



Z Index: 121

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 59 nm^3; this corresponds to an approximate mass of 53 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.303 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.303 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.28	3.67	3.36
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-22189 and PDB model 6XI0. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.05).

9.4 Atom inclusion (i)



At the recommended contour level, 61% of all backbone atoms, 60% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6050	0.4800
C	0.7520	0.5190
D	0.6410	0.4930
E	0.1920	0.3710
P	0.7530	0.5180
Q	0.6430	0.4930
R	0.1960	0.3720