



wwPDB EM Validation Summary Report ⓘ

Aug 13, 2026 – 04:47 AM JST

PDB ID : 8XNP / pdb_00008xnp
EMDB ID : EMD-38510
Title : Respiratory complex Peripheral Arm of CI, close form B, focus-refined map of type IA, Wild type mouse under Cold Acclimation
Authors : Shin, Y.-C.; Liao, M.
Deposited on : 2023-12-30
Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	115	34% 21% 72% ...
2	B	224	6% 30% 35% 5% 30%
3	C	263	8% 38% 36% 25%
4	D	463	5% 34% 45% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	248	
6	F	464	
7	G	727	
8	H	318	
9	I	212	
10	P	377	
11	Q	175	
12	R	116	
13	S	99	
14	T	156	
15	V	116	
16	W	131	
17	X	172	
18	Z	144	
19	a	70	
20	b	84	
21	q	145	
22	r	113	
23	s	104	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
24	SF4	B	301	-	-	X	-
24	SF4	F	502	-	-	X	-
24	SF4	G	802	-	-	X	-
24	SF4	I	303	-	-	X	-
24	SF4	I	304	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
27	FES	E	301	-	-	X	-
27	FES	G	803	-	-	X	-
33	CDL	a	101	-	-	X	-

2 Entry composition

There are 33 unique types of molecules in this entry. The entry contains 34048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-ubiquinone oxidoreductase chain 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	113	Total	C	N	O	S	0	0
			915	623	131	154	7		

- Molecule 2 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	157	Total	C	N	O	S	0	0
			1258	802	227	215	14		

- Molecule 3 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 3, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	198	Total	C	N	O	S	0	0
			1641	1060	279	299	3		

- Molecule 4 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	385	Total	C	N	O	S	0	0
			3088	1970	533	562	23		

- Molecule 5 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	210	Total	C	N	O	S	0	0
			1635	1039	275	310	11		

- Molecule 6 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	426	Total	C	N	O	S	0	0
			3288	2073	588	605	22		

- Molecule 7 is a protein called NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	687	Total	C	N	O	S	0	0
			5287	3316	918	1012	41		

- Molecule 8 is a protein called NADH-ubiquinone oxidoreductase chain 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	318	Total	C	N	O	S	0	0
			2538	1705	384	427	22		

- Molecule 9 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	174	Total	C	N	O	S	0	0
			1398	880	240	266	12		

- Molecule 10 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	P	340	Total	C	N	O	S	0	0
			2730	1765	479	479	7		

- Molecule 11 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	Q	118	Total	C	N	O	S	0	0
			957	608	165	180	4		

- Molecule 12 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	R	83	Total	C	N	O	S	0	0
			660	411	120	126	3		

- Molecule 13 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	S	83	Total	C	N	O	S	0	0
			667	419	126	119	3		

- Molecule 14 is a protein called Acyl carrier protein, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	T	75	Total	C	N	O	S	0	0
			604	388	89	122	5		

- Molecule 15 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	V	112	Total	C	N	O	S	0	0
			915	596	152	164	3		

- Molecule 16 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	W	114	Total	C	N	O	S	0	0
			970	619	180	165	6		

- Molecule 17 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	X	142	Total	C	N	O	S	0	0
			1164	736	209	209	10		

- Molecule 18 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	Z	139	Total	C	N	O	S	0	0
			1152	741	204	199	8		

- Molecule 19 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	a	67	Total	C	N	O	S	0	0
			548	356	97	91	4		

- Molecule 20 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	b	80	Total	C	N	O	S	0	0
			628	414	99	111	4		

- Molecule 21 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	q	120	Total	C	N	O	S	0	0
			1004	645	178	177	4		


- Molecule 22 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	r	51	Total	C	N	O	S	0	0
			418	266	78	73	1		

- Molecule 23 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 3, mitochondrial.


Mol	Chain	Residues	Atoms				AltConf	Trace
23	s	13	Total	C	N	O	0	0
			107	69	15	23		

- Molecule 24 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			AltConf
24	B	1	Total	Fe	S	0
			8	4	4	
24	F	1	Total	Fe	S	0
			8	4	4	
24	G	1	Total	Fe	S	0
			8	4	4	
24	G	1	Total	Fe	S	0
			8	4	4	
24	I	1	Total	Fe	S	0
			8	4	4	
24	I	1	Total	Fe	S	0
			8	4	4	

- Molecule 25 is Ubiquinone-9 (CCD ID: UQ9) (formula: $C_{54}H_{82}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
25	B	1	Total	C	O	0
			30	26	4	

- Molecule 26 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: C₄₄H₈₈NO₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
26	B	1	Total	C	N	O	P	0
			35	25	1	8	1	
26	H	1	Total	C	N	O	P	0
			42	32	1	8	1	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf
26	I	1	Total	C	N	O	P	0
			47	37	1	8	1	

- Molecule 27 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			AltConf
27	E	1	Total	Fe	S	0
			4	2	2	
27	G	1	Total	Fe	S	0
			4	2	2	

- Molecule 28 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $\text{C}_{17}\text{H}_{21}\text{N}_4\text{O}_9\text{P}$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
28	F	1	Total	C	N	O	P	0
			31	17	4	9	1	

- Molecule 29 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula: $C_{41}H_{82}NO_8P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
29	I	1	Total	C	N	O	P	0
			51	41	1	8	1	
29	Z	1	Total	C	N	O	P	0
			46	36	1	8	1	

- Molecule 30 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
30	P	1	Total	C	N	O	P	0
			48	21	7	17	3	

- Molecule 31 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
31	R	1	Total	Zn	0
			1	1	

- Molecule 32 is {S}-[2-[3-[(2 {R})-3,3-dimethyl-2-oxidanyl-4-phosphonooxy-butanoyl]amino]propanoylamino]ethyl] (3 {S})-3-oxidanyltetradecanethioate (CCD ID: EHZ) (formula: $C_{25}H_{49}N_2O_9PS$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf	
32	W	1	Total	C	N	O	P	S	0
			32	19	2	9	1	1	

- Molecule 33 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
33	a	1	Total	C	O	P	0
			57	38	17	2	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADH-ubiquinone oxidoreductase chain 3




- Molecule 2: NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial



- Molecule 3: NADH dehydrogenase [ubiquinone] iron-sulfur protein 3, mitochondrial






• Molecule 6: NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial





• Molecule 7: NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial






● Molecule 8: NADH-ubiquinone oxidoreductase chain 1







• Molecule 9: NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial




• Molecule 10: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial




• Molecule 11: NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial




- Molecule 12: NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial




- Molecule 13: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2




- Molecule 14: Acyl carrier protein, mitochondrial




- Molecule 15: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5





- Molecule 16: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6



- Molecule 17: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8




- Molecule 18: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13




- Molecule 19: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1








- Molecule 20: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3




- Molecule 21: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12



- Molecule 22: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7



- Molecule 23: NADH dehydrogenase [ubiquinone] flavoprotein 3, mitochondrial



PRO	THR	ASP	THR	THR	THR	TYR	LYS	ASN	LEU	GLN	HIS	HIS	ASP	TYR																																																																																																																																																																																																																																																																																																																																																																																																																								
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	22658	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	46.1, 45.9	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k), GATAN K3 (6k x 4k)	Depositor
Maximum map value	4.385	Depositor
Minimum map value	-1.018	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.083	Depositor
Recommended contour level	0.55	Depositor
Map size ($\text{\AA}$)	422.40002, 422.40002, 422.40002	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.1, 1.1, 1.1	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FES, EH2, NDP, SF4, CDL, FMN, PC1, UQ9, 3PE, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/941	1.24	10/1285 (0.8%)
2	B	1.00	1/1289 (0.1%)	1.36	11/1744 (0.6%)
3	C	0.87	1/1687 (0.1%)	1.31	14/2297 (0.6%)
4	D	0.94	5/3162 (0.2%)	1.31	22/4276 (0.5%)
5	E	0.81	2/1675 (0.1%)	1.25	15/2282 (0.7%)
6	F	0.88	1/3363 (0.0%)	1.44	54/4543 (1.2%)
7	G	0.89	6/5374 (0.1%)	1.59	104/7281 (1.4%)
8	H	0.97	2/2615 (0.1%)	1.55	60/3574 (1.7%)
9	I	0.90	0/1427	1.54	19/1927 (1.0%)
10	P	0.79	1/2804 (0.0%)	1.26	34/3802 (0.9%)
11	Q	0.87	1/980 (0.1%)	1.28	12/1324 (0.9%)
12	R	0.63	0/671	1.19	9/903 (1.0%)
13	S	0.93	1/678 (0.1%)	1.65	17/915 (1.9%)
14	T	0.91	0/613	1.51	9/826 (1.1%)
15	V	0.87	0/937	1.47	14/1270 (1.1%)
16	W	0.80	0/993	1.33	15/1335 (1.1%)
17	X	0.84	1/1191 (0.1%)	1.44	21/1605 (1.3%)
18	Z	0.76	0/1183	1.10	7/1597 (0.4%)
19	a	0.82	0/561	1.36	5/755 (0.7%)
20	b	0.70	0/651	0.91	1/895 (0.1%)
21	q	0.96	2/1037 (0.2%)	1.44	15/1408 (1.1%)
22	r	0.98	1/426 (0.2%)	1.57	9/573 (1.6%)
23	s	0.76	0/108	1.08	0/147
All	All	0.88	25/34366 (0.1%)	1.41	477/46564 (1.0%)

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	H	60	PRO	N-CD	13.80	1.67	1.47
6	F	235	VAL	N-CA	-9.28	1.34	1.46
11	Q	53	ILE	C-O	7.78	1.33	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	G	203	ASP	CA-C	6.75	1.62	1.52
4	D	106	VAL	C-N	-6.55	1.24	1.33

The worst 5 of 477 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	174	THR	N-CA-C	-16.80	89.10	114.64
8	H	208	VAL	N-CA-C	-13.66	100.83	111.62
7	G	251	ILE	N-CA-C	13.31	126.81	108.17
7	G	77	MET	N-CA-C	-13.09	97.28	113.38
9	I	164	VAL	N-CA-C	-12.87	100.09	112.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	915	0	957	189	0
2	B	1258	0	1270	164	0
3	C	1641	0	1596	131	0
4	D	3088	0	3059	305	0
5	E	1635	0	1626	181	0
6	F	3288	0	3243	348	0
7	G	5287	0	5323	551	0
8	H	2538	0	2627	496	0
9	I	1398	0	1359	205	0
10	P	2730	0	2747	245	0
11	Q	957	0	949	73	0
12	R	660	0	639	74	0
13	S	667	0	685	72	0
14	T	604	0	596	88	0
15	V	915	0	954	76	0
16	W	970	0	991	89	0
17	X	1164	0	1150	83	0
18	Z	1152	0	1148	104	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
19	a	548	0	562	96	0
20	b	628	0	628	136	0
21	q	1004	0	962	83	0
22	r	418	0	434	46	0
23	s	107	0	100	9	0
24	B	8	0	0	6	0
24	F	8	0	0	10	0
24	G	16	0	0	5	0
24	I	16	0	0	16	0
25	B	30	0	35	17	0
26	B	35	0	44	6	0
26	H	42	0	58	13	0
26	I	47	0	71	6	0
27	E	4	0	0	5	0
27	G	4	0	0	4	0
28	F	31	0	19	2	0
29	I	51	0	82	17	0
29	Z	46	0	69	6	0
30	P	48	0	26	7	0
31	R	1	0	0	0	0
32	W	32	0	0	7	0
33	a	57	0	58	28	0
All	All	34048	0	34067	3216	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 47.

The worst 5 of 3216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:H:127:TYR:CE1	8:H:208:VAL:HG23	1.47	1.48
1:A:51:PHE:HD1	4:D:80:MET:SD	1.36	1.47
9:I:53:ALA:CB	20:b:14:ALA:HB2	1.48	1.42
1:A:51:PHE:CE1	4:D:80:MET:HE1	1.55	1.42
9:I:50:MET:SD	20:b:10:LYS:HG2	1.61	1.39

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/115 (96%)	100 (90%)	11 (10%)	0	100	100
2	B	155/224 (69%)	144 (93%)	10 (6%)	1 (1%)	21	54
3	C	196/263 (74%)	186 (95%)	10 (5%)	0	100	100
4	D	383/463 (83%)	357 (93%)	26 (7%)	0	100	100
5	E	208/248 (84%)	188 (90%)	20 (10%)	0	100	100
6	F	424/464 (91%)	404 (95%)	20 (5%)	0	100	100
7	G	685/727 (94%)	627 (92%)	58 (8%)	0	100	100
8	H	316/318 (99%)	296 (94%)	19 (6%)	1 (0%)	36	67
9	I	170/212 (80%)	169 (99%)	1 (1%)	0	100	100
10	P	338/377 (90%)	316 (94%)	22 (6%)	0	100	100
11	Q	116/175 (66%)	114 (98%)	2 (2%)	0	100	100
12	R	81/116 (70%)	76 (94%)	5 (6%)	0	100	100
13	S	81/99 (82%)	78 (96%)	3 (4%)	0	100	100
14	T	73/156 (47%)	72 (99%)	1 (1%)	0	100	100
15	V	110/116 (95%)	107 (97%)	3 (3%)	0	100	100
16	W	112/131 (86%)	106 (95%)	6 (5%)	0	100	100
17	X	140/172 (81%)	129 (92%)	11 (8%)	0	100	100
18	Z	137/144 (95%)	129 (94%)	8 (6%)	0	100	100
19	a	65/70 (93%)	57 (88%)	8 (12%)	0	100	100
20	b	78/84 (93%)	68 (87%)	10 (13%)	0	100	100
21	q	116/145 (80%)	114 (98%)	2 (2%)	0	100	100
22	r	47/113 (42%)	43 (92%)	4 (8%)	0	100	100
23	s	11/104 (11%)	11 (100%)	0	0	100	100
All	All	4153/5036 (82%)	3891 (94%)	260 (6%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	H	92	PRO
2	B	195	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/104 (98%)	101 (99%)	1 (1%)	68	75
2	B	133/185 (72%)	131 (98%)	2 (2%)	57	71
3	C	180/227 (79%)	180 (100%)	0	100	100
4	D	332/395 (84%)	330 (99%)	2 (1%)	78	79
5	E	182/206 (88%)	181 (100%)	1 (0%)	81	80
6	F	341/370 (92%)	341 (100%)	0	100	100
7	G	579/610 (95%)	578 (100%)	1 (0%)	87	85
8	H	280/280 (100%)	279 (100%)	1 (0%)	84	81
9	I	148/178 (83%)	148 (100%)	0	100	100
10	P	297/325 (91%)	296 (100%)	1 (0%)	86	83
11	Q	105/153 (69%)	104 (99%)	1 (1%)	68	75
12	R	70/96 (73%)	69 (99%)	1 (1%)	59	71
13	S	74/80 (92%)	73 (99%)	1 (1%)	59	71
14	T	69/135 (51%)	68 (99%)	1 (1%)	59	71
15	V	100/102 (98%)	100 (100%)	0	100	100
16	W	108/114 (95%)	108 (100%)	0	100	100
17	X	129/154 (84%)	129 (100%)	0	100	100
18	Z	120/123 (98%)	119 (99%)	1 (1%)	73	77
19	a	57/60 (95%)	57 (100%)	0	100	100
20	b	71/73 (97%)	71 (100%)	0	100	100
21	q	108/131 (82%)	107 (99%)	1 (1%)	70	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
22	r	44/96 (46%)	44 (100%)	0	100	100
23	s	13/95 (14%)	13 (100%)	0	100	100
All	All	3642/4292 (85%)	3627 (100%)	15 (0%)	81	81

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	H	229	THR
18	Z	7	LYS
10	P	316	LYS
21	q	138	VAL
13	S	68	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 96 such sidechains are listed below:

Mol	Chain	Res	Type
8	H	169	GLN
10	P	275	HIS
8	H	212	ASN
10	P	79	GLN
11	Q	88	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 1 is monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
24	SF4	I	303	9	0,12,12	-	-	-		
24	SF4	F	502	6	0,12,12	-	-	-		
26	PC1	I	302	-	46,46,53	1.00	2 (4%)	52,54,61	1.08	3 (5%)
27	FES	G	803	7	0,4,4	-	-	-		
33	CDL	a	101	-	56,56,99	1.19	4 (7%)	62,68,111	1.29	6 (9%)
26	PC1	H	401	-	41,41,53	1.04	2 (4%)	47,49,61	1.06	4 (8%)
27	FES	E	301	5	0,4,4	-	-	-		
29	3PE	Z	401	-	45,45,50	0.93	2 (4%)	48,50,55	1.01	2 (4%)
24	SF4	B	301	2	0,12,12	-	-	-		
24	SF4	G	801	7	0,12,12	-	-	-		
24	SF4	I	304	9	0,12,12	-	-	-		
25	UQ9	B	302	-	30,30,58	1.03	3 (10%)	36,39,73	0.77	1 (2%)
32	EHZ	W	201	-	27,31,37	1.89	7 (25%)	37,41,47	1.85	11 (29%)
30	NDP	P	401	-	49,52,52	1.20	5 (10%)	66,80,80	1.67	11 (16%)
28	FMN	F	501	-	33,33,33	1.40	5 (15%)	48,50,50	1.22	7 (14%)
29	3PE	I	301	-	50,50,50	0.90	2 (4%)	53,55,55	1.03	3 (5%)
24	SF4	G	802	7	0,12,12	-	-	-		
26	PC1	B	303	-	34,34,53	1.15	2 (5%)	40,42,61	1.15	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
24	SF4	I	303	9	-	-	0/6/5/5
24	SF4	F	502	6	-	-	0/6/5/5
26	PC1	I	302	-	-	13/50/50/57	-
27	FES	G	803	7	-	-	0/1/1/1
33	CDL	a	101	-	-	14/67/67/110	-
26	PC1	H	401	-	-	15/45/45/57	-
27	FES	E	301	5	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
29	3PE	Z	401	-	-	14/49/49/54	-
24	SF4	B	301	2	-	-	0/6/5/5
24	SF4	G	801	7	-	-	0/6/5/5
24	SF4	I	304	9	-	-	0/6/5/5
25	UQ9	B	302	-	-	14/24/48/81	0/1/1/1
32	EHZ	W	201	-	-	21/39/39/45	-
30	NDP	P	401	-	-	6/34/77/77	0/5/5/5
28	FMN	F	501	-	-	4/18/18/18	0/3/3/3
29	3PE	I	301	-	-	16/54/54/54	-
24	SF4	G	802	7	-	-	0/6/5/5
26	PC1	B	303	-	-	11/38/38/57	-

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
32	W	201	EHZ	C15-N2	5.37	1.45	1.33
32	W	201	EHZ	C12-N1	5.13	1.45	1.33
28	F	501	FMN	C9A-C5A	4.76	1.49	1.41
30	P	401	NDP	C5A-C4A	4.24	1.46	1.39
26	I	302	PC1	O31-C31	4.19	1.45	1.33

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
32	W	201	EHZ	C8-C9-S1	6.27	121.39	113.63
30	P	401	NDP	C5A-C4A-N3A	-6.03	118.88	126.75
30	P	401	NDP	N3A-C4A-N9A	4.73	134.87	127.08
33	a	101	CDL	OA6-CA5-C11	4.36	120.91	111.50
26	I	302	PC1	O21-C21-C22	4.22	120.59	111.50

There are no chirality outliers.

5 of 128 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
25	B	302	UQ9	C12-C13-C14-C16
25	B	302	UQ9	C12-C13-C14-C15
25	B	302	UQ9	C1-C6-C7-C8
25	B	302	UQ9	C5-C6-C7-C8
26	B	303	PC1	C1-O11-P-O12



There are no ring outliers.



17 monomers are involved in 155 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
24	I	303	SF4	11	0
24	F	502	SF4	10	0
26	I	302	PC1	6	0
27	G	803	FES	4	0
33	a	101	CDL	28	0
26	H	401	PC1	13	0
27	E	301	FES	5	0
29	Z	401	3PE	6	0
24	B	301	SF4	6	0
24	I	304	SF4	5	0
25	B	302	UQ9	17	0
32	W	201	EHZ	7	0
30	P	401	NDP	7	0
28	F	501	FMN	2	0
29	I	301	3PE	17	0
24	G	802	SF4	5	0
26	B	303	PC1	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-38510. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 235




Y Index: 175




Z Index: 198


6.3.2 Raw map



X Index: 236



Y Index: 175




Z Index: 3


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.55. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 173 nm³; this corresponds to an approximate mass of 156 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.286 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.56	4.06	3.68
Unmasked-calculated*	7.35	11.60	7.77

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.35 differs from the reported value 3.5 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-38510 and PDB model 8XNP. Per-residue inclusion information can be found in section 3 on page 15.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.55 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.55).

9.4 Atom inclusion [i](#)



At the recommended contour level, 73% of all backbone atoms, 51% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.55) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5130	 0.3890
A	 0.4770	 0.4190
B	 0.5980	 0.4540
C	 0.6130	 0.4550
D	 0.6340	 0.4570
E	 0.3230	 0.3040
F	 0.3990	 0.3280
G	 0.5880	 0.4140
H	 0.5230	 0.3910
I	 0.6130	 0.4340
P	 0.5230	 0.3740
Q	 0.4870	 0.4230
R	 0.2260	 0.3350
S	 0.5490	 0.3490
T	 0.3420	 0.2640
V	 0.5820	 0.3960
W	 0.5120	 0.4000
X	 0.6020	 0.3660
Z	 0.5710	 0.3710
a	 0.5580	 0.3970
b	 0.5610	 0.3770
q	 0.1460	 0.3440
r	 0.2560	 0.3110
s	 0.0660	 0.2200

