



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:06 AM UTC

PDB ID : 2XTA / pdb\_00002xta  
Title : Crystal structure of the SucA domain of Mycobacterium smegmatis alpha-ketoglutarate decarboxylase in complex with acetyl-CoA (triclinic form)  
Authors : Wagner, T.; Bellinzoni, M.; Wehenkel, A.M.; O'Hare, H.M.; Alzari, P.M.  
Deposited on : 2010-10-05  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	868	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div>• 5%</div> </div> </div>
1	B	868	<div> <div>6%</div> <div> <div></div> <div>87%</div> <div>6%</div> <div>6%</div> </div> </div>
1	C	868	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>• 6%</div> </div> </div>
1	D	868	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>6%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 26097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2-OXOGLUTARATE DECARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	826	Total	C	N	O	S	0	0	0
			6360	4007	1128	1202	23			
1	B	814	Total	C	N	O	S	0	0	0
			6223	3925	1105	1169	24			
1	C	817	Total	C	N	O	S	0	0	0
			6309	3980	1112	1193	24			
1	D	818	Total	C	N	O	S	0	0	0
			6252	3939	1105	1184	24			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	360	GLY	-	expression tag	UNP A0R2B1
B	360	GLY	-	expression tag	UNP A0R2B1
C	360	GLY	-	expression tag	UNP A0R2B1
D	360	GLY	-	expression tag	UNP A0R2B1

- Molecule 2 is THIAMINE DIPHOSPHATE (CCD ID: TPP) (formula: C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O<sub>7</sub>P<sub>2</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
2	B	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
2	C	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
2	D	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Mg 1	0	0
3	B	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0
3	D	1	Total 1	Mg 1	0	0

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		

- Molecule 5 is ACETYL COENZYME \*A (CCD ID: ACO) (formula: C<sub>23</sub>H<sub>38</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			33	12	5	13	3		
5	C	1	Total	C	N	O	P	0	0
			33	12	5	13	3		
5	D	1	Total	C	N	O	P	0	0
			33	12	5	13	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	190	Total	O	0	0
			190	190		
6	B	186	Total	O	0	0
			186	186		
6	C	183	Total	O	0	0
			183	183		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	183	Total 183	O 183	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: 2-OXOGLUTARATE DECARBOXYLASE




#### • Molecule 1: 2-OXOGLUTARATE DECARBOXYLASE



#### • Molecule 1: 2-OXOGLUTARATE DECARBOXYLASE



● Molecule 1: 2-OXOGLUTARATE DECARBOXYLASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.55Å 83.58Å 160.07Å 99.59° 98.94° 100.68°	Depositor
Resolution (Å)	78.11 – 2.20 78.11 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.1 (78.11-2.20) 93.1 (78.11-2.20)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.9.3	Depositor
R, $R_{free}$	0.188 , 0.214 0.202 , 0.229	Depositor DCC
$R_{free}$ test set	9391 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.4	Xtriage
Anisotropy	0.617	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.022 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	26097	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACO, MG, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.87	0/6492	1.24	11/8818 (0.1%)
1	B	0.86	1/6349 (0.0%)	1.24	11/8623 (0.1%)
1	C	0.88	1/6438 (0.0%)	1.24	10/8737 (0.1%)
1	D	0.87	1/6377 (0.0%)	1.24	19/8656 (0.2%)
All	All	0.87	3/25656 (0.0%)	1.24	51/34834 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	641	MET	SD-CE	-11.77	1.50	1.79
1	C	870	MET	SD-CE	-7.04	1.61	1.79
1	D	870	MET	SD-CE	-5.09	1.66	1.79

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	627	GLY	CA-C-N	7.14	129.73	120.44
1	A	627	GLY	C-N-CA	7.14	129.73	120.44
1	D	836	ASP	CA-CB-CG	6.42	119.02	112.60
1	A	399	SER	CA-C-N	6.25	127.78	122.28
1	A	399	SER	C-N-CA	6.25	127.78	122.28

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6360	0	6105	28	0
1	B	6223	0	5967	25	0
1	C	6309	0	6084	26	0
1	D	6252	0	5998	28	0
2	A	26	0	16	1	0
2	B	26	0	16	2	0
2	C	26	0	16	1	0
2	D	26	0	16	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	33	0	13	0	0
5	C	33	0	13	0	0
5	D	33	0	13	0	0
6	A	190	0	0	0	0
6	B	186	0	0	0	0
6	C	183	0	0	0	0
6	D	183	0	0	1	0
All	All	26097	0	24257	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:641:MET:HE3	1:B:643:HIS:HE1	1.19	1.06
1:B:641:MET:HE3	1:B:643:HIS:CE1	2.01	0.94
1:B:641:MET:CE	1:B:643:HIS:HE1	1.94	0.80
1:B:641:MET:CE	1:B:643:HIS:CE1	2.65	0.77
1:A:1044:THR:HG22	1:A:1047:ASN:H	1.54	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	818/868 (94%)	800 (98%)	18 (2%)	0	100	100
1	B	804/868 (93%)	786 (98%)	18 (2%)	0	100	100
1	C	805/868 (93%)	789 (98%)	16 (2%)	0	100	100
1	D	808/868 (93%)	792 (98%)	16 (2%)	0	100	100
All	All	3235/3472 (93%)	3167 (98%)	68 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	649/726 (89%)	634 (98%)	15 (2%)	44	59
1	B	628/726 (86%)	612 (98%)	16 (2%)	42	56
1	C	648/726 (89%)	629 (97%)	19 (3%)	37	51
1	D	634/726 (87%)	624 (98%)	10 (2%)	55	71
All	All	2559/2904 (88%)	2499 (98%)	60 (2%)	44	59

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1087	GLU
1	D	950	LEU
1	C	766	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	930	LEU
1	D	1187	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1030	GLN
1	D	524	GLN
1	D	503	GLN
1	D	748	ASN
1	B	933	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 8 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACO	D	2004	-	35,35,53	1.83	8 (22%)	51,54,79	2.07	13 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TPP	B	2001	3	26,27,27	1.32	2 (7%)	38,40,40	1.56	11 (28%)
5	ACO	C	2004	-	35,35,53	1.76	7 (20%)	51,54,79	2.19	13 (25%)
2	TPP	C	2001	3	26,27,27	1.45	3 (11%)	38,40,40	1.63	9 (23%)
2	TPP	D	2001	3	26,27,27	1.36	2 (7%)	38,40,40	1.58	8 (21%)
5	ACO	A	2004	-	35,35,53	1.94	8 (22%)	51,54,79	2.17	13 (25%)
2	TPP	A	2001	3	26,27,27	1.36	3 (11%)	38,40,40	1.62	10 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ACO	D	2004	-	-	5/25/41/67	0/3/3/3
2	TPP	B	2001	3	-	3/17/17/17	0/2/2/2
5	ACO	C	2004	-	-	5/25/41/67	0/3/3/3
2	TPP	C	2001	3	-	3/17/17/17	0/2/2/2
2	TPP	D	2001	3	-	3/17/17/17	0/2/2/2
5	ACO	A	2004	-	-	5/25/41/67	0/3/3/3
2	TPP	A	2001	3	-	3/17/17/17	0/2/2/2

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	2004	ACO	P2A-O3A	5.74	1.65	1.59
5	A	2004	ACO	P2A-O3A	5.47	1.65	1.59
5	C	2004	ACO	P2A-O3A	4.66	1.64	1.59
5	C	2004	ACO	P1A-O3A	4.53	1.64	1.59
5	A	2004	ACO	P3B-O7A	4.18	1.63	1.50

The worst 5 of 77 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2004	ACO	N3A-C2A-N1A	-5.87	119.69	128.58
5	C	2004	ACO	C5A-C4A-N3A	-5.60	119.01	126.72
5	D	2004	ACO	N3A-C2A-N1A	-5.57	120.15	128.58
5	C	2004	ACO	N3A-C2A-N1A	-5.53	120.21	128.58
5	D	2004	ACO	C5A-C4A-N3A	-5.19	119.58	126.72

There are no chirality outliers.

5 of 27 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	2001	TPP	PA-O3A-PB-O3B
2	C	2001	TPP	PA-O3A-PB-O3B
2	D	2001	TPP	PA-O3A-PB-O2B
2	D	2001	TPP	PA-O3A-PB-O3B
5	A	2004	ACO	C5B-O5B-P1A-O1A



There are no ring outliers.



4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2001	TPP	2	0
2	C	2001	TPP	1	0
2	D	2001	TPP	1	0
2	A	2001	TPP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	826/868 (95%)	0.35	40 (4%)	35	32	23, 40, 81, 111	0
1	B	814/868 (93%)	0.41	54 (6%)	24	21	21, 41, 80, 108	0
1	C	817/868 (94%)	0.30	42 (5%)	33	30	20, 39, 78, 105	0
1	D	818/868 (94%)	0.32	39 (4%)	35	32	18, 39, 78, 107	0
All	All	3275/3472 (94%)	0.35	175 (5%)	32	29	18, 40, 79, 111	0

The worst 5 of 175 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	413	TRP	6.1
1	D	397	PHE	5.4
1	D	472	ASP	5.2
1	A	412	LEU	5.1
1	B	560	GLU	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ACO	D	2004	33/51	0.72	0.15	53,75,91,92	0
5	ACO	A	2004	33/51	0.86	0.12	47,60,77,77	0
5	ACO	C	2004	33/51	0.87	0.11	43,54,78,82	0
4	CA	A	2003	1/1	0.94	0.09	75,75,75,75	0
4	CA	B	2003	1/1	0.95	0.15	74,74,74,74	0
4	CA	C	2003	1/1	0.95	0.14	55,55,55,55	0
4	CA	D	2003	1/1	0.95	0.16	57,57,57,57	0
2	TPP	D	2001	26/26	0.97	0.08	18,27,40,55	0
2	TPP	B	2001	26/26	0.97	0.08	18,30,44,50	0
2	TPP	A	2001	26/26	0.98	0.06	19,28,38,50	0
3	MG	C	2002	1/1	0.98	0.04	18,18,18,18	0
2	TPP	C	2001	26/26	0.98	0.07	18,30,43,49	0
3	MG	A	2002	1/1	0.99	0.05	18,18,18,18	0
3	MG	D	2002	1/1	0.99	0.03	23,23,23,23	0
3	MG	B	2002	1/1	0.99	0.04	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ACO D 2004:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ACO A 2004:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ACO C 2004:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around TPP D 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around TPP B 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around TPP A 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TPP C 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.