



wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 05:09 AM UTC

PDB ID : 7YI4 / pdb_00007yi4
EMDB ID : EMD-33851
Title : Cryo-EM structure of Rpd3S complex bound to H3K36me3 nucleosome in close state
Authors : Li, H.T.; Yan, C.Y.; Guan, H.P.; Wang, P.
Deposited on : 2022-07-14
Resolution : 3.96 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	7646 (3.46 - 4.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1536	12% 25% 8% 67%
2	B	433	60% 24% 15%
3	C	401	21% 46% 26% 29%
3	E	401	24% 18% 8% 74%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	684	
4	F	684	
5	H	102	
5	L	102	
6	I	129	
6	M	129	
7	J	122	
7	N	122	
8	O	167	
9	P	167	
10	G	135	
10	K	135	

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 25351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulatory protein SIN3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	506	Total	C	N	O	S	0	0
			4258	2737	712	796	13		

- Molecule 2 is a protein called Histone deacetylase RPD3.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	368	Total	C	N	O	S	0	0
			2923	1858	495	545	25		

- Molecule 3 is a protein called Chromatin modification-related protein EAF3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	286	Total	C	N	O	S	0	0
			2326	1495	379	438	14		
3	E	104	Total	C	N	O	S	0	0
			853	551	142	155	5		

- Molecule 4 is a protein called Transcriptional regulatory protein RCO1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	316	Total	C	N	O	S	0	0
			2588	1644	445	481	18		
4	F	20	Total	C	N	O	S	0	0
			178	117	30	30	1		

- Molecule 5 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	H	79	Total	C	N	O	S	0	0
			633	399	124	109	1		
5	L	81	Total	C	N	O	S	0	0
			646	407	126	112	1		

- Molecule 6 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	I	107	Total	C	N	O	0	0
			823	519	161	143		
6	M	107	Total	C	N	O	0	0
			823	519	161	143		

- Molecule 7 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	J	93	Total	C	N	O	S	0	0
			726	457	130	137	2		
7	N	93	Total	C	N	O	S	0	0
			726	457	130	137	2		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	29	THR	SER	engineered mutation	UNP P02281
N	29	THR	SER	engineered mutation	UNP P02281

- Molecule 8 is a DNA chain called Wisdom 601 DNA (167-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
8	O	151	Total	C	N	O	P	0	0
			3076	1462	554	909	151		

- Molecule 9 is a DNA chain called Wisdom 601 DNA (167-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
9	P	151	Total	C	N	O	P	0	0
			3115	1474	587	903	151		


- Molecule 10 is a protein called Histone H3.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	G	100	Total	C	N	O	S	0	0
			830	526	160	141	3		
10	K	99	Total	C	N	O	S	0	0
			822	521	159	140	2		



- Molecule 11 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
11	B	1	Total 1	Zn 1	0
11	D	4	Total 4	Zn 4	0






- Molecule 4: Transcriptional regulatory protein RCO1




- Molecule 4: Transcriptional regulatory protein RCO1







- Molecule 6: Histone H2A




- Molecule 7: Histone H2B 1.1



- Molecule 7: Histone H2B 1.1



- Molecule 8: Wisdom 601 DNA (167-MER)



- Molecule 9: Wisdom 601 DNA (167-MER)

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	87742	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.036	Depositor
Minimum map value	-0.004	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.011	Depositor
Map size (Å)	324.75, 324.75, 324.75	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0825, 1.0825, 1.0825	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M3L, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/4346	0.33	1/5848 (0.0%)
2	B	0.14	0/2997	0.37	1/4051 (0.0%)
3	C	0.14	0/2374	0.37	0/3202
3	E	0.13	0/875	0.35	0/1174
4	D	0.11	0/2649	0.33	0/3566
4	F	0.09	0/181	0.23	0/237
5	H	0.17	0/640	0.30	0/857
5	L	0.16	0/653	0.38	0/873
6	I	0.15	0/833	0.26	0/1124
6	M	0.16	0/833	0.27	0/1124
7	J	0.16	0/737	0.32	0/993
7	N	0.17	0/737	0.35	0/993
8	O	0.24	0/3445	0.41	0/5310
9	P	0.23	0/3499	0.36	0/5403
10	G	0.17	0/830	0.29	0/1114
10	K	0.16	0/823	0.36	0/1106
All	All	0.17	0/26452	0.35	2/36975 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1164	MET	CA-CB-CG	5.54	125.18	114.10
2	B	47	MET	CA-CB-CG	5.18	124.47	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4258	0	4211	91	0
2	B	2923	0	2805	86	0
3	C	2326	0	2337	92	0
3	E	853	0	840	25	0
4	D	2588	0	2534	91	0
4	F	178	0	185	5	0
5	H	633	0	673	15	0
5	L	646	0	687	15	0
6	I	823	0	882	19	0
6	M	823	0	882	20	0
7	J	726	0	747	13	0
7	N	726	0	747	19	0
8	O	3076	0	1697	74	0
9	P	3115	0	1694	95	0
10	G	830	0	881	25	0
10	K	822	0	871	27	0
11	B	1	0	0	0	0
11	D	4	0	0	0	0
All	All	25351	0	22673	619	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 619 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:O:-50:DC:N3	9:P:50:DG:N1	2.23	0.86
3:C:365:TRP:HA	3:C:368:MET:HE2	1.59	0.85
4:D:420:CYS:SG	4:D:448:HIS:CE1	2.70	0.84
3:C:347:MET:SD	3:C:351:SER:OG	2.38	0.82
8:O:66:DC:H2'	8:O:67:DA:C8	2.15	0.81

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	494/1536 (32%)	474 (96%)	20 (4%)	0	100	100
2	B	366/433 (84%)	353 (96%)	13 (4%)	0	100	100
3	C	280/401 (70%)	270 (96%)	10 (4%)	0	100	100
3	E	100/401 (25%)	97 (97%)	3 (3%)	0	100	100
4	D	304/684 (44%)	278 (91%)	26 (9%)	0	100	100
4	F	18/684 (3%)	18 (100%)	0	0	100	100
5	H	77/102 (76%)	77 (100%)	0	0	100	100
5	L	79/102 (78%)	76 (96%)	3 (4%)	0	100	100
6	I	105/129 (81%)	102 (97%)	3 (3%)	0	100	100
6	M	105/129 (81%)	102 (97%)	3 (3%)	0	100	100
7	J	91/122 (75%)	88 (97%)	3 (3%)	0	100	100
7	N	91/122 (75%)	90 (99%)	1 (1%)	0	100	100
10	G	97/135 (72%)	94 (97%)	3 (3%)	0	100	100
10	K	97/135 (72%)	96 (99%)	1 (1%)	0	100	100
All	All	2304/5115 (45%)	2215 (96%)	89 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	472/1391 (34%)	472 (100%)	0	100	100
2	B	311/367 (85%)	310 (100%)	1 (0%)	86	85
3	C	260/359 (72%)	259 (100%)	1 (0%)	84	84
3	E	89/359 (25%)	89 (100%)	0	100	100
4	D	299/653 (46%)	299 (100%)	0	100	100
4	F	20/653 (3%)	20 (100%)	0	100	100
5	H	65/78 (83%)	65 (100%)	0	100	100
5	L	66/78 (85%)	66 (100%)	0	100	100
6	I	84/101 (83%)	84 (100%)	0	100	100
6	M	84/101 (83%)	84 (100%)	0	100	100
7	J	79/102 (78%)	79 (100%)	0	100	100
7	N	79/102 (78%)	79 (100%)	0	100	100
10	G	87/109 (80%)	87 (100%)	0	100	100
10	K	85/109 (78%)	85 (100%)	0	100	100
All	All	2080/4562 (46%)	2078 (100%)	2 (0%)	87	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	288	MET
3	C	350	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
6	I	68	ASN
3	E	104	ASN
10	K	108	ASN
10	K	39	HIS
2	B	188	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	M3L	K	36	10	10,11,12	0.51	0	9,14,16	0.47	0
10	M3L	G	36	10	10,11,12	0.52	0	9,14,16	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	M3L	K	36	10	-	4/9/10/12	-
10	M3L	G	36	10	-	2/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	G	36	M3L	N-CA-CB-CG
10	G	36	M3L	C-CA-CB-CG
10	K	36	M3L	C-CA-CB-CG
10	K	36	M3L	CG-CD-CE-NZ
10	K	36	M3L	CE-CD-CG-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	K	36	M3L	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	G	36	M3L	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33851. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 112




Y Index: 143




Z Index: 135


6.3.2 Raw map



X Index: 112



Y Index: 143




Z Index: 135


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.011. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 326 nm^3 ; this corresponds to an approximate mass of 295 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.253 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.253 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.96	-	-
Author-provided FSC curve	3.94	5.92	4.13
Unmasked-calculated*	6.03	8.10	6.63

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.03 differs from the reported value 3.96 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-33851 and PDB model 7YI4. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay [i](#)



X



Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.011 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.011).

9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 77% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.011) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7660	 0.2300
A	 0.5460	 0.1200
B	 0.8890	 0.1270
C	 0.5940	 0.1580
D	 0.6000	 0.1340
E	 0.0600	 0.0970
F	 0.0860	 0.0400
G	 0.9120	 0.3530
H	 0.9230	 0.3650
I	 0.9080	 0.3590
J	 0.9250	 0.3520
K	 0.8700	 0.3210
L	 0.8550	 0.3350
M	 0.8870	 0.3600
N	 0.9030	 0.3340
O	 0.9820	 0.3320
P	 0.9800	 0.3330

