Full wwPDB X-ray Structure Validation Report (i)

Oct 27, 2026 – 10:07 AM EDT

PDB ID	:	9ZF4 / pdb_00009zf4
Title	:	The structure of human Vacuolar Protein Sorting 34 catalytic domain bound to RD-II-83
Authors	:	Litchfield, C.M.; Abiodun, W.O.; Peterson, M.A.; Moody, J.D.
Deposited on	:	2025-12-01
Resolution	:	2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	594	7% 83% 6% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	A1C15	A	913	-	X	-	-

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 8900 atoms, of which 4379 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 3-kinase catalytic subunit type 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	528	Total	C	H	N	O	S	0	18	0
			8601	2740	4311	724	801	25			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	278	MET	-	expression tag	UNP Q8NEB9
A	279	GLY	-	expression tag	UNP Q8NEB9
A	280	HIS	-	expression tag	UNP Q8NEB9
A	281	HIS	-	expression tag	UNP Q8NEB9
A	282	HIS	-	expression tag	UNP Q8NEB9
A	283	HIS	-	expression tag	UNP Q8NEB9
A	284	HIS	-	expression tag	UNP Q8NEB9
A	285	HIS	-	expression tag	UNP Q8NEB9
A	286	HIS	-	expression tag	UNP Q8NEB9
A	287	HIS	-	expression tag	UNP Q8NEB9
A	288	HIS	-	expression tag	UNP Q8NEB9
A	289	HIS	-	expression tag	UNP Q8NEB9

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			13	3	7	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	3	6	1		
3	A	1	Total	C	H	O	0	0
			15	4	9	2		
3	A	1	Total	C	H	O	0	0
			17	4	10	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 17 4 10 3	0	0
3	A	1	Total C H O 17 4 10 3	0	0

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O S 10 2 6 1 1	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Mg 2 2	0	0


- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Cl 2 2	0	0

- Molecule 7 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total K 1 1	0	0

- Molecule 8 is methyl 2-[(8R)-pyrazolo[1,5-a]pyrimidin-3-yl]-1,3-benzothiazole-6-carboxylate (CCD ID: A1C15) (formula: $C_{15}H_{10}N_4O_2S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C H N O S 32 15 10 4 2 1	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	163	Total O 163 163	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphatidylinositol 3-kinase catalytic subunit type 3

Chain A:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α, β, γ	113.07Å 113.07Å 145.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.17 – 2.09 61.17 – 2.09	Depositor EDS
% Data completeness (in resolution range)	100.0 (61.17-2.09) 100.0 (61.17-2.09)	Depositor EDS
R_{merge}	0.02	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.29 (at 2.08Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.200 , 0.231 0.202 , 0.232	Depositor DCC
R_{free} test set	2861 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å^2)	46.6	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.42 , 51.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	8900	wwPDB-VP
Average B, all atoms (Å^2)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1C15, CL, K, GOL, PEG, MG, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.39	0/4460	0.56	0/6033

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4290	4311	4202	20	0
2	A	6	7	8	0	0
3	A	31	45	39	1	0
4	A	4	6	6	0	0
5	A	2	0	0	0	0
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	22	10	0	0	0
9	A	163	0	0	0	0
All	All	4521	4379	4255	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:MET:HE2	1:A:656:LYS:HD3	1.64	0.79
1:A:733[A]:CYS:SG	1:A:777:MET:SD	2.89	0.71
1:A:694:LEU:HD11	1:A:782:GLU:HG2	1.75	0.68
1:A:749:LEU:HD11	1:A:757:LEU:HD21	1.84	0.59
1:A:529:LEU:HD21	1:A:539:ARG:HD3	1.93	0.51
1:A:514:HIS:CE1	1:A:518:LEU:HD11	2.47	0.49
1:A:366:LEU:HB2	1:A:380:ALA:HB2	1.93	0.49
1:A:594:PRO:O	1:A:673:LEU:HD22	2.14	0.47
1:A:305:THR:HG21	1:A:839:VAL:HG11	1.99	0.45
1:A:443:SER:HB2	1:A:546:GLN:HG2	1.99	0.44
1:A:824:VAL:O	1:A:835:PRO:HB3	2.17	0.44
1:A:555:LEU:HD21	1:A:573:LEU:HD12	1.99	0.44
1:A:777:MET:HB3	1:A:866[B]:SER:OG	2.18	0.44
1:A:296:LEU:O	1:A:300:VAL:HG23	2.17	0.44
1:A:514:HIS:NE2	1:A:518:LEU:HD11	2.33	0.44
1:A:385:ARG:HG2	1:A:478:PHE:CE1	2.53	0.43
1:A:374:PRO:O	1:A:378[A]:ARG:HG2	2.19	0.42
1:A:849:ASP:OD1	1:A:849:ASP:N	2.52	0.42
1:A:757:LEU:HD23	1:A:757:LEU:C	2.45	0.41
1:A:446:ILE:HG13	1:A:494:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	539/594 (91%)	531 (98%)	8 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	481/538 (89%)	481 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	473	GLN
1	A	504	GLN
1	A	519	ASN
1	A	545	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	PEG	A	903	-	5,5,6	0.26	0	4,4,5	0.15	0
3	PEG	A	902	-	3,3,6	0.36	0	2,2,5	0.28	0
8	A1C15	A	913	-	25,25,25	3.33	16 (64%)	32,36,36	4.66	23 (71%)
3	PEG	A	904	-	6,6,6	0.20	0	5,5,5	0.14	0
4	DMS	A	905	-	3,3,3	0.76	0	3,3,3	1.15	0
3	PEG	A	906	-	6,6,6	0.23	0	5,5,5	0.09	0
2	GOL	A	901	-	5,5,5	0.48	0	5,5,5	0.60	0
3	PEG	A	907	-	6,6,6	0.10	0	5,5,5	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	903	-	-	1/3/3/4	-
3	PEG	A	902	-	-	0/0/1/4	-
8	A1C15	A	913	-	-	6/8/10/10	0/4/4/4
3	PEG	A	904	-	-	4/4/4/4	-
3	PEG	A	906	-	-	2/4/4/4	-
2	GOL	A	901	-	-	0/4/4/4	-
3	PEG	A	907	-	-	0/4/4/4	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	913	A1C15	C15-N12	7.22	1.54	1.39
8	A	913	A1C15	C01-N06	5.55	1.48	1.38
8	A	913	A1C15	C02-S11	5.15	1.82	1.75
8	A	913	A1C15	C17-C16	5.05	1.61	1.50
8	A	913	A1C15	O21-C16	4.80	1.44	1.33
8	A	913	A1C15	C10-C02	4.67	1.53	1.46
8	A	913	A1C15	C19-C15	4.43	1.46	1.40
8	A	913	A1C15	C18-C17	3.74	1.45	1.39
8	A	913	A1C15	C14-C17	3.21	1.44	1.39
8	A	913	A1C15	C19-C18	3.19	1.44	1.38
8	A	913	A1C15	O21-C22	-2.82	1.39	1.45
8	A	913	A1C15	C01-N07	2.35	1.40	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	913	A1C15	C09-N03	2.34	1.36	1.32
8	A	913	A1C15	C15-C13	2.32	1.43	1.40
8	A	913	A1C15	C05-N06	2.17	1.41	1.37
8	A	913	A1C15	C05-C04	2.06	1.39	1.35

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
8	A	913	A1C15	C10-C02-N12	9.79	140.25	122.82
8	A	913	A1C15	C08-N07-C01	9.61	123.02	115.55
8	A	913	A1C15	C13-S11-C02	8.84	94.77	88.92
8	A	913	A1C15	C09-N03-N06	6.63	110.30	103.81
8	A	913	A1C15	S11-C02-N12	-6.56	107.79	116.17
8	A	913	A1C15	C05-N06-N03	6.37	136.48	127.01
8	A	913	A1C15	C14-C13-C15	6.07	126.49	121.99
8	A	913	A1C15	C19-C15-C13	-5.69	111.44	119.59
8	A	913	A1C15	C10-C02-S11	-5.17	111.94	121.53
8	A	913	A1C15	O21-C16-C17	4.99	119.78	112.31
8	A	913	A1C15	C18-C17-C14	-4.96	113.51	119.25
8	A	913	A1C15	O21-C16-O20	-4.42	114.86	123.46
8	A	913	A1C15	C04-C05-N06	4.24	124.07	117.44
8	A	913	A1C15	N06-C01-N07	-3.98	117.17	121.96
8	A	913	A1C15	C19-C15-N12	3.87	139.21	128.21
8	A	913	A1C15	C10-C01-N07	3.87	135.08	132.07
8	A	913	A1C15	C15-N12-C02	3.54	117.73	109.84
8	A	913	A1C15	C18-C17-C16	3.39	127.88	120.40
8	A	913	A1C15	C13-C15-N12	-3.35	109.35	115.20
8	A	913	A1C15	C05-N06-C01	-2.89	117.08	122.70
8	A	913	A1C15	C05-C04-C08	-2.89	115.53	118.87
8	A	913	A1C15	C19-C18-C17	2.73	123.72	120.80
8	A	913	A1C15	C09-C10-C01	-2.07	103.02	104.57

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	913	A1C15	O21-C16-C17-C18
8	A	913	A1C15	C17-C16-O21-C22
8	A	913	A1C15	O20-C16-O21-C22
8	A	913	A1C15	O21-C16-C17-C14
8	A	913	A1C15	O20-C16-C17-C14
8	A	913	A1C15	O20-C16-C17-C18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	904	PEG	O2-C3-C4-O4
3	A	906	PEG	O2-C3-C4-O4
3	A	903	PEG	C4-C3-O2-C2
3	A	904	PEG	C4-C3-O2-C2
3	A	904	PEG	O1-C1-C2-O2
3	A	906	PEG	O1-C1-C2-O2
3	A	904	PEG	C1-C2-O2-C3


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	904	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1C15 A 913




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	528/594 (88%)	0.31	41 (7%) 19 20	19, 60, 91, 114	8 (1%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	712	ASN	5.3
1	A	867[A]	VAL	5.0
1	A	293	ARG	4.7
1	A	441	ASP	4.5
1	A	417	PRO	4.5
1	A	833	LEU	4.0
1	A	307	GLN	4.0
1	A	294[A]	ASP	3.7
1	A	866[A]	SER	3.6
1	A	711	GLU	3.4
1	A	448	SER	3.2
1	A	378[A]	ARG	3.2
1	A	778	LYS	3.0
1	A	509	ARG	2.9
1	A	295	GLN	2.9
1	A	444	GLN	2.8
1	A	502	GLU	2.7
1	A	773	LEU	2.7
1	A	581	GLU	2.6
1	A	340	ASP	2.6
1	A	580	ASN	2.6
1	A	790	THR	2.6
1	A	779	LEU	2.5
1	A	803	TYR	2.3
1	A	584	ASN	2.3
1	A	473	GLN	2.3
1	A	311	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	510	ASP	2.2
1	A	522	ARG	2.2
1	A	771	LYS	2.2
1	A	529	LEU	2.2
1	A	662[A]	ASN	2.2
1	A	776	PRO	2.1
1	A	562	GLU	2.1
1	A	370	HIS	2.1
1	A	347	GLN	2.1
1	A	325	ASN	2.1
1	A	583	MET	2.0
1	A	318	LYS	2.0
1	A	582	LYS	2.0
1	A	517	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	PEG	A	902	4/7	0.63	0.35	50,66,81,81	0
3	PEG	A	906	7/7	0.67	0.24	71,85,105,105	0
7	K	A	912	1/1	0.72	0.20	113,113,113,113	0
3	PEG	A	903	6/7	0.74	0.27	66,80,94,94	0
3	PEG	A	907	7/7	0.75	0.26	68,82,97,103	0
3	PEG	A	904	7/7	0.80	0.24	56,70,83,83	17
5	MG	A	909	1/1	0.81	0.36	76,76,76,76	0
6	CL	A	910	1/1	0.84	0.34	99,99,99,99	0
4	DMS	A	905	4/4	0.84	0.23	61,82,83,94	0

Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	MG	A	908	1/1	0.87	0.19	71,71,71,71	0
8	A1C15	A	913	22/22	0.91	0.14	44,62,91,96	0
2	GOL	A	901	6/6	0.95	0.11	55,63,72,76	0
6	CL	A	911	1/1	0.95	0.17	63,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1C15 A 913:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.