



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 23, 2026 – 05:05 am BST

PDB ID : 6ZI5 / pdb_00006zi5
Title : Ultrafast Structural Response to Charge Redistribution Within a Photosynthetic Reaction Centre - 300 ps (a) structure
Authors : Baath, P.; Dods, R.; Braenden, G.; Neutze, R.
Deposited on : 2020-06-24
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	336	% 84% 15% .
2	H	258	2% 86% 14% .
3	L	273	% 78% 22%
4	M	323	% 88% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	BCB	L	301[A]	X	-	-	-
10	BCB	L	301[B]	X	-	-	-
10	BCB	L	302[A]	X	-	-	-
10	BCB	L	302[B]	X	-	-	-
10	BCB	M	403	X	-	-	-
10	BCB	M	404[A]	X	-	-	-
10	BCB	M	404[B]	X	-	-	-
5	HEC	C	401	X	-	-	-
5	HEC	C	402	X	-	-	-
5	HEC	C	403	X	-	-	-
5	HEC	C	404	X	-	-	-

2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 11266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Photosynthetic reaction center cytochrome c subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	332	Total	C	N	O	S	0	0	0
			2602	1640	466	478	18			

- Molecule 2 is a protein called Reaction center protein H chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	258	Total	C	N	O	S	0	0	0
			2018	1292	344	380	2			


- Molecule 3 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	273	Total	C	N	O	S	0	41	0
			2508	1678	409	413	8			

- Molecule 4 is a protein called Reaction center protein M chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	M	323	Total	C	N	O	S	0	51	0
			2977	1983	491	490	13			

- Molecule 5 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{36}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 6 is DIACYL GLYCEROL (CCD ID: DGA) (formula: C₃₉H₇₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			37	33	4		

- Molecule 7 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	O	S	0	0
			5	4	1		
7	H	1	Total	O	S	0	0
			5	4	1		
7	H	1	Total	O	S	0	0
			5	4	1		
7	H	1	Total	O	S	0	0
			5	4	1		
7	H	1	Total	O	S	0	0
			5	4	1		
7	M	1	Total	O	S	0	0
			5	4	1		
7	M	1	Total	O	S	0	0
			5	4	1		
7	M	1	Total	O	S	0	0
			5	4	1		
7	M	1	Total	O	S	0	0
			5	4	1		
7	M	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	M	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: C₁₄H₃₁NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	H	1	Total	C	N	O	0	0
			16	14	1	1		
8	H	1	Total	C	N	O	0	0
			16	14	1	1		
8	H	1	Total	C	N	O	0	0
			16	14	1	1		
8	L	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 9 is HEPTANE-1,2,3-TRIOL (CCD ID: HTO) (formula: C₇H₁₆O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	H	1	Total	C	O	0	0
			10	7	3		
9	L	1	Total	C	O	0	0
			10	7	3		

- Molecule 10 is BACTERIOCHLOROPHYLL B (CCD ID: BCB) (formula: $C_{55}H_{72}MgN_4O_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	L	1	Total	C	Mg	N	O	0	1
			132	110	2	8	12		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	L	1	Total	C	Mg	N	O	0	1
			132	110	2	8	12		
10	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
10	M	1	Total	C	Mg	N	O	0	1
			132	110	2	8	12		

- Molecule 11 is BACTERIOPHEOPHYTIN B (CCD ID: BPB) (formula: $C_{55}H_{74}N_4O_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	L	1	Total	C	N	O	0	1
			130	110	8	12		
11	M	1	Total	C	N	O	0	0
			65	55	4	6		

- Molecule 12 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	M	1	Total	Fe	0	1
			2	2		

- Molecule 13 is MENAQUINONE-7 (CCD ID: MQ7) (formula: $C_{46}H_{64}O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	M	1	Total	C	O	0	1
			96	92	4		

- Molecule 14 is 15-cis-1,2-dihydroneurosporene (CCD ID: NS5) (formula: C₄₀H₆₀).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	M	1	Total	C	0	0
			40	40		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	C	3	Total 3	O 3	0	0
15	H	2	Total 2	O 2	0	0
15	L	4	Total 4	O 4	0	0
15	M	4	Total 4	O 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Photosynthetic reaction center cytochrome c subunit




- Molecule 2: Reaction center protein H chain




- Molecule 3: Reaction center protein L chain



- Molecule 4: Reaction center protein M chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	226.40 Å 226.40 Å 113.70 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.75 – 2.80 37.75 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (37.75-2.80) 99.9 (37.75-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.31 Å)	Xtriage
Refinement program	REFMAC 5.8.01158, PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.176 , 0.202 0.184 , 0.209	Depositor DCC
R_{free} test set	5696 reflections (4.49%)	wwPDB-VP
Wilson B-factor (Å ²)	74.7	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 65.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11266	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DGA, FE, SO4, HEC, BPB, NS5, MQ7, LDA, BCB, HTO, FME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.38	0/2669	0.60	0/3637
2	H	0.37	0/2055	0.58	0/2807
3	L	0.34	0/2612	0.54	0/3568
4	M	0.35	0/3101	0.52	0/4242
All	All	0.36	0/10437	0.56	0/14254

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2602	0	2578	51	0
2	H	2018	0	2020	39	0
3	L	2508	0	2390	51	0
4	M	2977	0	2832	35	0
5	C	172	0	120	3	0
6	C	37	0	58	0	0
7	C	5	0	0	0	0
7	H	20	0	0	1	0
7	M	35	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	H	48	0	93	2	0
8	L	16	0	31	1	0
9	H	10	0	16	0	0
9	L	10	0	16	0	0
10	L	264	0	288	14	0
10	M	198	0	216	7	0
11	L	130	0	148	8	0
11	M	65	0	74	5	0
12	M	2	0	0	0	0
13	M	96	0	128	3	0
14	M	40	0	60	4	0
15	C	3	0	0	0	0
15	H	2	0	0	0	0
15	L	4	0	0	0	0
15	M	4	0	0	0	0
All	All	11266	0	11068	175	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:13:GLY:CA	1:C:19:MET:HE2	1.69	1.21
1:C:13:GLY:HA3	1:C:19:MET:HE2	1.17	1.08
2:H:89:LYS:HE2	2:H:110:ASP:CB	1.96	0.95
4:M:262[B]:SER:HA	4:M:265[B]:ARG:HG3	1.49	0.92
2:H:89:LYS:HE2	2:H:110:ASP:HB2	1.55	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	330/336 (98%)	319 (97%)	11 (3%)	0	100	100
2	H	256/258 (99%)	245 (96%)	8 (3%)	3 (1%)	10	34
3	L	312/273 (114%)	297 (95%)	12 (4%)	3 (1%)	12	38
4	M	372/323 (115%)	355 (95%)	16 (4%)	1 (0%)	36	66
All	All	1270/1190 (107%)	1216 (96%)	47 (4%)	7 (1%)	24	51

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	47	LEU
2	H	50	VAL
2	H	46	PRO
3	L	165[A]	LEU
3	L	165[B]	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	281/282 (100%)	279 (99%)	2 (1%)	76	91
2	H	212/212 (100%)	212 (100%)	0	100	100
3	L	253/218 (116%)	253 (100%)	0	100	100
4	M	288/249 (116%)	288 (100%)	0	100	100
All	All	1034/961 (108%)	1032 (100%)	2 (0%)	87	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	123	GLN
1	C	316	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
4	M	26	ASN
4	M	87	GLN
2	H	8	GLN
2	H	72	HIS
2	H	102	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FME	H	1	2	8,9,10	1.01	0	7,9,11	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FME	H	1	2	-	1/7/9/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	1	FME	CB-CG-SD-CE

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 38 ligands modelled in this entry, 2 are monoatomic - leaving 36 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	BPB	L	303[A]	-	57,70,70	2.66	14 (24%)	56,101,101	2.89	20 (35%)
9	HTO	L	305	-	9,9,9	0.75	0	10,10,10	0.89	1 (10%)
7	SO4	H	704	-	4,4,4	0.12	0	6,6,6	0.14	0
10	BCB	L	301[A]	3	68,74,74	2.08	16 (23%)	79,115,115	3.25	28 (35%)
8	LDA	L	304	-	12,15,15	0.37	0	14,17,17	0.73	0
10	BCB	M	404[B]	4	68,74,74	2.05	15 (22%)	79,115,115	3.20	30 (37%)
8	LDA	H	707	-	12,15,15	0.31	0	14,17,17	1.17	1 (7%)
7	SO4	M	411	-	4,4,4	0.27	0	6,6,6	0.24	0
7	SO4	C	406	-	4,4,4	0.24	0	6,6,6	0.18	0
7	SO4	M	408	-	4,4,4	0.11	0	6,6,6	0.14	0
7	SO4	M	410	-	4,4,4	0.26	0	6,6,6	0.21	0
5	HEC	C	401	1	50,50,50	1.39	6 (12%)	68,82,82	1.64	15 (22%)
7	SO4	M	409	-	4,4,4	0.20	0	6,6,6	0.26	0
10	BCB	M	403	4	68,74,74	1.96	16 (23%)	79,115,115	3.28	25 (31%)
11	BPB	L	303[B]	-	57,70,70	2.74	13 (22%)	56,101,101	2.76	18 (32%)
14	NS5	M	406	-	39,39,39	1.38	1 (2%)	44,46,46	2.04	12 (27%)
10	BCB	L	301[B]	3	68,74,74	2.05	16 (23%)	79,115,115	3.20	29 (36%)
7	SO4	H	702	-	4,4,4	0.21	0	6,6,6	0.28	0
7	SO4	M	412	-	4,4,4	0.32	0	6,6,6	0.33	0
7	SO4	H	705	-	4,4,4	0.15	0	6,6,6	0.15	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
13	MQ7	M	402[A]	-	49,49,49	1.54	2 (4%)	60,63,63	1.49	10 (16%)
8	LDA	H	706	-	12,15,15	0.38	0	14,17,17	0.79	0
10	BCB	L	302[A]	3	68,74,74	1.95	16 (23%)	79,115,115	3.14	29 (36%)
9	HTO	H	708	-	9,9,9	0.79	0	10,10,10	0.79	0
7	SO4	M	407	-	4,4,4	0.13	0	6,6,6	0.41	0
5	HEC	C	403	1	50,50,50	1.39	6 (12%)	68,82,82	1.74	13 (19%)
7	SO4	M	413	-	4,4,4	0.22	0	6,6,6	0.14	0
5	HEC	C	404	1	50,50,50	1.45	6 (12%)	68,82,82	1.57	14 (20%)
6	DGA	C	405	1	36,36,43	1.17	3 (8%)	38,38,45	1.22	4 (10%)
8	LDA	H	701	-	12,15,15	0.36	0	14,17,17	0.68	0
13	MQ7	M	402[B]	-	49,49,49	1.43	2 (4%)	60,63,63	1.45	10 (16%)
10	BCB	M	404[A]	4	68,74,74	2.04	16 (23%)	79,115,115	3.27	26 (32%)
10	BCB	L	302[B]	3	68,74,74	2.01	16 (23%)	79,115,115	3.01	28 (35%)
5	HEC	C	402	1	50,50,50	1.36	6 (12%)	68,82,82	1.59	10 (14%)
11	BPB	M	405	-	57,70,70	2.68	13 (22%)	56,101,101	2.83	20 (35%)
7	SO4	H	703	-	4,4,4	0.22	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	BPB	L	303[A]	-	-	8/37/105/105	0/5/6/6
9	HTO	L	305	-	-	4/10/10/10	-
10	BCB	L	301[A]	3	3/3/21/26	7/41/137/137	-
10	BCB	M	404[B]	4	3/3/21/26	12/41/137/137	-
8	LDA	L	304	-	-	6/13/13/13	-
8	LDA	H	707	-	-	11/13/13/13	-
5	HEC	C	401	1	1/1/3/7	7/14/54/54	-
10	BCB	M	403	4	3/3/21/26	18/41/137/137	-
11	BPB	L	303[B]	-	-	7/37/105/105	0/5/6/6
14	NS5	M	406	-	-	10/43/43/43	-
10	BCB	L	301[B]	3	3/3/21/26	11/41/137/137	-
13	MQ7	M	402[A]	-	-	0/41/61/61	0/2/2/2
8	LDA	H	706	-	-	6/13/13/13	-
10	BCB	L	302[A]	3	2/2/21/26	6/41/137/137	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	HTO	H	708	-	-	1/10/10/10	-
5	HEC	C	403	1	1/1/3/7	4/14/54/54	-
5	HEC	C	404	1	1/1/3/7	7/14/54/54	-
6	DGA	C	405	1	-	15/37/37/45	-
8	LDA	H	701	-	-	4/13/13/13	-
13	MQ7	M	402[B]	-	-	2/41/61/61	0/2/2/2
10	BCB	M	404[A]	4	3/3/21/26	13/41/137/137	-
10	BCB	L	302[B]	3	2/2/21/26	7/41/137/137	-
5	HEC	C	402	1	1/1/3/7	7/14/54/54	-
11	BPB	M	405	-	-	8/37/105/105	0/5/6/6

The worst 5 of 183 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	L	303[B]	BPB	C1B-C2B	9.67	1.50	1.39
11	L	303[A]	BPB	C1B-C2B	9.36	1.50	1.39
11	M	405	BPB	C1B-C2B	8.85	1.49	1.39
13	M	402[A]	MQ7	C3-C2	8.07	1.49	1.35
11	L	303[B]	BPB	C1D-C2D	7.89	1.48	1.39

The worst 5 of 343 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	M	403	BCB	C1C-NC-C4C	-20.38	97.55	106.71
10	M	404[A]	BCB	C1C-NC-C4C	-19.43	97.97	106.71
10	L	301[A]	BCB	C1C-NC-C4C	-19.03	98.15	106.71
10	L	301[B]	BCB	C1C-NC-C4C	-18.18	98.53	106.71
10	L	302[A]	BCB	C1C-NC-C4C	-16.85	99.13	106.71

5 of 23 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	C	401	HEC	NA
5	C	402	HEC	NA
5	C	403	HEC	NA
5	C	404	HEC	NA
10	L	301[A]	BCB	NA

5 of 181 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	C	405	DGA	OG1-CG1-CG2-OG2
6	C	405	DGA	OG1-CG1-CG2-CG3
8	H	706	LDA	C2-C1-N1-O1
8	H	706	LDA	C2-C1-N1-CM1
8	H	706	LDA	C2-C1-N1-CM2


There are no ring outliers.


17 monomers are involved in 44 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	L	303[A]	BPB	4	0
10	L	301[A]	BCB	3	0
8	L	304	LDA	1	0
10	M	404[B]	BCB	3	0
8	H	707	LDA	1	0
5	C	401	HEC	1	0
10	M	403	BCB	4	0
11	L	303[B]	BPB	4	0
14	M	406	NS5	4	0
10	L	301[B]	BCB	7	0
7	H	702	SO4	1	0
13	M	402[A]	MQ7	3	0
8	H	701	LDA	1	0
10	M	404[A]	BCB	1	0
10	L	302[B]	BCB	6	0
5	C	402	HEC	2	0
11	M	405	BPB	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























Ligand MQ7 M 402 (B)





Ligand BCB M 404 (A)



Ligand BCB L 302 (B)







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	332/336 (98%)	-0.40	2 (0%) 85 80	74, 93, 120, 146	0
2	H	257/258 (99%)	-0.07	6 (2%) 61 51	78, 103, 147, 193	0
3	L	273/273 (100%)	-0.39	2 (0%) 84 77	38, 90, 115, 134	41 (15%)
4	M	323/323 (100%)	-0.34	2 (0%) 85 80	38, 90, 115, 139	51 (15%)
All	All	1185/1190 (99%)	-0.31	12 (1%) 79 72	38, 94, 125, 193	92 (7%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	50	VAL	4.0
2	H	52	LEU	4.0
2	H	53	ALA	3.9
3	L	1	ALA	3.5
4	M	89	PHE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FME	H	1	10/11	0.95	0.12	90,102,112,127	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	SO4	M	413	5/5	0.51	0.14	145,160,166,179	0
7	SO4	M	411	5/5	0.61	0.11	130,156,170,175	0
7	SO4	M	410	5/5	0.63	0.11	127,133,147,161	0
9	HTO	L	305	10/10	0.63	0.31	96,120,130,131	0
7	SO4	H	703	5/5	0.68	0.11	149,152,158,170	0
7	SO4	M	412	5/5	0.69	0.13	125,128,152,179	0
7	SO4	C	406	5/5	0.75	0.10	129,143,150,163	0
7	SO4	M	408	5/5	0.79	0.11	114,115,124,146	0
9	HTO	H	708	10/10	0.85	0.23	102,115,129,131	0
6	DGA	C	405	37/44	0.85	0.22	97,131,165,170	0
8	LDA	L	304	16/16	0.89	0.27	102,125,150,154	0
8	LDA	H	706	16/16	0.94	0.18	83,99,167,169	0
11	BPB	M	405	65/65	0.94	0.15	78,91,172,180	0
7	SO4	H	702	5/5	0.96	0.09	104,114,121,125	0
7	SO4	H	705	5/5	0.96	0.06	107,107,119,119	5
7	SO4	M	407	5/5	0.96	0.06	99,112,123,124	0
8	LDA	H	707	16/16	0.96	0.13	97,115,132,136	0
14	NS5	M	406	40/40	0.96	0.16	80,96,136,138	0
7	SO4	H	704	5/5	0.97	0.17	97,101,115,115	5
13	MQ7	M	402[A]	48/48	0.97	0.13	78,86,123,146	48
13	MQ7	M	402[B]	48/48	0.97	0.13	78,86,123,146	48
10	BCB	M	403	66/66	0.97	0.11	75,84,166,177	0
10	BCB	L	302[B]	66/66	0.98	0.10	72,80,112,115	66
8	LDA	H	701	16/16	0.98	0.11	81,96,111,111	0
10	BCB	M	404[A]	66/66	0.98	0.10	60,75,106,112	66
10	BCB	M	404[B]	66/66	0.98	0.10	60,75,106,112	66
11	BPB	L	303[A]	65/65	0.98	0.09	77,83,92,97	65
11	BPB	L	303[B]	65/65	0.98	0.09	77,83,92,97	65
7	SO4	M	409	5/5	0.98	0.05	89,97,103,104	0
10	BCB	L	301[A]	66/66	0.98	0.10	65,75,91,109	66
10	BCB	L	301[B]	66/66	0.98	0.10	65,75,91,109	66
10	BCB	L	302[A]	66/66	0.98	0.10	72,80,112,115	66
5	HEC	C	401	43/43	0.99	0.08	90,98,112,127	0
5	HEC	C	402	43/43	0.99	0.07	82,94,105,131	0
5	HEC	C	403	43/43	0.99	0.07	73,82,93,96	0
5	HEC	C	404	43/43	0.99	0.06	75,84,101,123	0
12	FE	M	401[B]	1/1	1.00	0.02	88,88,88,88	1

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
12	FE	M	401[A]	1/1	1.00	0.02	88,88,88,88	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around BPB M 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NS5 M 406:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MQ7 M 402 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MQ7 M 402 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB M 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BCB L 302 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB M 404 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BCB M 404 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around BPB L 303 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BPB L 303 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB L 301 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BCB L 301 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCB L 302 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC C 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.