



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:47 PM UTC

PDB ID : 6ZZI / pdb_00006zzi
Title : Crystal structure of the catalytic domain of *Corynebacterium glutamicum* acetyltransferase AceF (E2p).
Authors : Bruch, E.M.; Lexa-Sapart, N.; Bellinzoni, M.
Deposited on : 2020-08-04
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	3% 98% .
1	B	241	5% 94% 5% .
1	C	241	% 95% 5%
1	D	241	4% 95% 5%
1	E	241	5% 97% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	241	3% 93% 7%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	241	Total	C	N	O	S	0	1	0
			1839	1162	324	349	4			
1	B	239	Total	C	N	O	S	0	2	0
			1833	1159	324	346	4			
1	C	240	Total	C	N	O	S	0	1	0
			1834	1159	319	352	4			
1	D	241	Total	C	N	O	S	0	0	0
			1831	1157	321	349	4			
1	E	239	Total	C	N	O	S	0	1	0
			1822	1151	321	346	4			
1	F	239	Total	C	N	O	S	0	0	0
			1821	1150	319	348	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	435	GLY	-	expression tag	UNP Q8NNJ2
A	436	SER	-	expression tag	UNP Q8NNJ2
B	435	GLY	-	expression tag	UNP Q8NNJ2
B	436	SER	-	expression tag	UNP Q8NNJ2
C	435	GLY	-	expression tag	UNP Q8NNJ2
C	436	SER	-	expression tag	UNP Q8NNJ2
D	435	GLY	-	expression tag	UNP Q8NNJ2
D	436	SER	-	expression tag	UNP Q8NNJ2
E	435	GLY	-	expression tag	UNP Q8NNJ2
E	436	SER	-	expression tag	UNP Q8NNJ2
F	435	GLY	-	expression tag	UNP Q8NNJ2
F	436	SER	-	expression tag	UNP Q8NNJ2

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	242	Total 242	O 242	0	0
2	B	200	Total 200	O 200	0	0
2	C	242	Total 242	O 242	0	0
2	D	176	Total 176	O 176	0	0
2	E	177	Total 177	O 177	0	0
2	F	220	Total 220	O 220	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex




- Molecule 1: Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex



- Molecule 1: Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex




- Molecule 1: Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex




- Molecule 1: Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex





- Molecule 1: Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex

Chain F: 3% 93% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	158.28Å 158.28Å 63.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	79.14 – 1.93 79.14 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.2 (79.14-1.93) 99.3 (79.14-1.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 1.94Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.166 , 0.185 (Not available) , 0.181	Depositor DCC
R_{free} test set	6556 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l 0.015 for h,-h-k,-l 0.009 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12237	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	0/1870	0.92	0/2541
1	B	0.68	0/1867	0.92	0/2538
1	C	0.71	0/1865	0.92	0/2537
1	D	0.68	0/1859	0.96	0/2527
1	E	0.67	0/1853	0.92	0/2520
1	F	0.69	0/1849	0.91	0/2514
All	All	0.68	0/11163	0.92	0/15177

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1839	0	1897	4	0
1	B	1833	0	1891	5	0
1	C	1834	0	1885	5	0
1	D	1831	0	1884	6	0
1	E	1822	0	1869	5	0
1	F	1821	0	1869	7	0
2	A	242	0	0	0	0
2	B	200	0	0	0	0
2	C	242	0	0	0	0
2	D	176	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	177	0	0	0	0
2	F	220	0	0	0	0
All	All	12237	0	11295	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:573:ASN:HB3	1:E:575:LYS:HZ2	1.72	0.55
1:F:534:LEU:HD11	1:F:588:ILE:HG13	1.89	0.55
1:E:612:GLY:O	1:E:640:PRO:HD2	2.10	0.52
1:C:612:GLY:O	1:C:640:PRO:HD2	2.11	0.51
1:B:612:GLY:O	1:B:640:PRO:HD2	2.11	0.51
1:F:612:GLY:O	1:F:640:PRO:HD2	2.11	0.50
1:A:612:GLY:O	1:A:640:PRO:HD2	2.11	0.50
1:D:611:LEU:HD11	1:D:639:LEU:HB3	1.92	0.50
1:D:570:ARG:NH2	1:D:580:ASP:OD2	2.47	0.48
1:D:612:GLY:O	1:D:640:PRO:HD2	2.13	0.48
1:F:570:ARG:NH2	1:F:580:ASP:OD2	2.49	0.46
1:D:569:ASP:OD1	1:D:573:ASN:OD1	2.34	0.44
1:C:534:LEU:HD21	1:C:588:ILE:HD12	2.00	0.44
1:B:488:ILE:HG12	1:B:494:ASN:HA	2.00	0.44
1:A:569:ASP:CG	1:E:562:LYS:NZ	2.77	0.43
1:A:469:HIS:CD2	1:A:657:LEU:HD13	2.54	0.43
1:F:480:ARG:HE	1:F:496:THR:HA	1.84	0.42
1:F:469:HIS:CD2	1:F:657:LEU:HD13	2.54	0.42
1:A:569:ASP:CG	1:E:562:LYS:HZ3	2.28	0.42
1:C:469:HIS:CD2	1:C:657:LEU:HD13	2.55	0.41
1:B:468:LEU:HD21	1:B:638:PHE:CD1	2.55	0.41
1:B:534:LEU:HD21	1:B:588:ILE:HD12	2.01	0.41
1:F:603:LEU:HB3	1:F:610:ILE:HG22	2.02	0.41
1:C:468:LEU:HD21	1:C:638:PHE:CD1	2.55	0.41
1:D:469:HIS:CD2	1:D:657:LEU:HD13	2.56	0.41
1:F:468:LEU:HD21	1:F:638:PHE:CD1	2.56	0.41
1:C:480:ARG:HE	1:C:496:THR:HA	1.86	0.41
1:D:468:LEU:HD21	1:D:638:PHE:CD1	2.55	0.41
1:E:573:ASN:HB3	1:E:575:LYS:NZ	2.35	0.41
1:B:469:HIS:CD2	1:B:657:LEU:HD13	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	240/241 (100%)	240 (100%)	0	0	100	100
1	B	239/241 (99%)	239 (100%)	0	0	100	100
1	C	239/241 (99%)	238 (100%)	1 (0%)	0	100	100
1	D	239/241 (99%)	239 (100%)	0	0	100	100
1	E	238/241 (99%)	238 (100%)	0	0	100	100
1	F	237/241 (98%)	237 (100%)	0	0	100	100
All	All	1432/1446 (99%)	1431 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/204 (100%)	203 (100%)	0	100	100
1	B	202/204 (99%)	200 (99%)	2 (1%)	68	64
1	C	204/204 (100%)	202 (99%)	2 (1%)	68	64
1	D	202/204 (99%)	201 (100%)	1 (0%)	81	82
1	E	200/204 (98%)	199 (100%)	1 (0%)	81	82
1	F	201/204 (98%)	199 (99%)	2 (1%)	68	64

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	1212/1224 (99%)	1204 (99%)	8 (1%)	76	74

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	530	SER
1	B	591	ILE
1	C	574	ASN
1	C	635	GLN
1	D	635	GLN
1	E	536	ILE
1	F	555	LEU
1	F	626	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	483	ASN
1	A	550	HIS
1	B	635	GLN
1	C	483	ASN
1	C	550	HIS
1	C	674	GLN
1	D	483	ASN
1	D	550	HIS
1	D	573	ASN
1	D	635	GLN
1	E	483	ASN
1	E	635	GLN
1	F	483	ASN
1	F	579	ASN
1	F	607	GLN
1	F	646	GLN
1	F	674	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	241/241 (100%)	-0.16	8 (3%)	49	55	14, 24, 46, 58	1 (0%)
1	B	239/241 (99%)	0.05	12 (5%)	34	38	13, 26, 50, 65	2 (0%)
1	C	240/241 (99%)	-0.15	3 (1%)	75	80	13, 22, 45, 56	1 (0%)
1	D	241/241 (100%)	0.16	9 (3%)	45	51	14, 31, 57, 70	0
1	E	239/241 (99%)	0.10	11 (4%)	37	42	16, 29, 54, 62	1 (0%)
1	F	239/241 (99%)	-0.14	7 (2%)	53	60	13, 23, 47, 68	0
All	All	1439/1446 (99%)	-0.02	50 (3%)	47	53	13, 26, 52, 70	5 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	626	ASP	4.5
1	F	437	LEU	4.0
1	D	626	ASP	3.7
1	B	572	ARG	3.3
1	D	488	ILE	3.2
1	F	438	ARG	3.0
1	F	628	ILE	2.9
1	E	625	GLU	2.9
1	D	576	LEU	2.9
1	A	626	ASP	2.8
1	B	579	ASN	2.7
1	F	439	GLY	2.7
1	B	544	LEU	2.7
1	C	572	ARG	2.7
1	A	573	ASN	2.6
1	A	628	ILE	2.6
1	E	437	LEU	2.6
1	B	565	VAL	2.6
1	D	675	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	579	ASN	2.5
1	E	543	GLY	2.4
1	A	569	ASP	2.4
1	B	437	LEU	2.4
1	E	441	THR	2.4
1	E	444	VAL	2.4
1	B	576	LEU	2.4
1	B	539	ASP	2.4
1	D	627	GLY	2.4
1	C	567	LEU	2.3
1	B	493	VAL	2.3
1	D	498	LEU	2.3
1	E	447	ILE	2.3
1	D	628	ILE	2.3
1	E	541	PRO	2.2
1	B	578	PRO	2.2
1	E	453	MET	2.2
1	E	446	ARG	2.2
1	A	625	GLU	2.2
1	F	578	PRO	2.2
1	D	579	ASN	2.1
1	A	627	GLY	2.1
1	B	626	ASP	2.1
1	B	577	LYS	2.1
1	C	447	ILE	2.0
1	D	625	GLU	2.0
1	F	442	GLN	2.0
1	B	625	GLU	2.0
1	A	575	LYS	2.0
1	E	628	ILE	2.0
1	A	579	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.